

Fourier analysis of Whittaker functions on a real reductive group

Erik van den Ban

University of Utrecht

Reductive Groups 2020

Organizers: B. Krötz, J. Kuit

On Zoom, June 29, 2020

Whittaker functions

Setting

- ▶ G real reductive group
- ▶ K maximal compact, $G = KAN_0$ Iwasawa decomposition
- ▶ $\chi : N_0 \rightarrow U(1)$ unitary character, **regular (!)**

i.e.: $\forall \alpha \in \Sigma(\mathfrak{n}_0, \mathfrak{a})$ simple: $d\chi(e)|_{\mathfrak{g}_\alpha} \neq 0$.

Whittaker functions

$$\mathcal{M}(G/N_0, \chi) := \{f : G \xrightarrow{\text{meas}} \mathbb{C} \mid f(xn) = \chi(n)^{-1}f(x) \quad (x \in G, n \in N_0)\}$$

$$L^2(G/N_0, \chi) := \{f \in \mathcal{M}(G/N_0, \chi) \mid |f| \in L^2(G/N_0)\}$$

- ▶ Left reg^r repⁿ: $L = \text{Ind}_{N_0}^G(\chi)$ is unitary

Whittaker Plancherel formula

Abstractly

$$\text{▶ } \text{Ind}_{N_0}^G(\chi) = \int_G^{\oplus} m_{\pi} \pi d\mu(\pi).$$

Concrete realization

- ▶ Harish-Chandra, Announcement 1982.

Details in Collected Papers Vol 5 (posthumous), 141- 307,
eds. R. Gangolli, V.S. Varadarajan, Springer 2018.

Final step not clear.

- ▶ N.R. Wallach, Independent treatment.

Real reductive groups II, Acad. Press 1992.

Erroneous estimate. Repair addressed in arXiv:1705.06787.

- ▶ **Today: final step in HC through new inversion theorem.**

Discrete part of decomposition

Discrete part

$\pi \in \widehat{G}$ (unitary dual) is said to appear discretely in $L^2(G/N_0, \chi)$ if it can be realized as a closed subrepresentation.

Theorem (HC, W)

If $\pi \in \widehat{G}$ appears discretely in $L^2(G/N_0, \chi)$, then it appears discretely in $L^2(G)$, i.e., it belongs to the discrete series of G .

Corollary

If $\pi \in \widehat{G}$ appears discretely in $L^2(G/N_0, \chi)$, then its infinitesimal character is real and regular, while $\text{rk}(\mathfrak{k}) = \text{rk}(\mathfrak{g})$.

This result is crucial for the separation of tempered spectra in the Whittaker Plancherel decomposition.

Schwartz functions

Define $\rho \in \mathfrak{a}^*$ by $\rho(X) = \frac{1}{2} \operatorname{tr}(\operatorname{ad}(X)|_{\mathfrak{N}_0})$. Let $\mathfrak{Z} := \operatorname{center} U(\mathfrak{g})$

Definition (Schwartz space)

$\mathcal{C}(G/N_0, \chi)$: the space of $f \in C^\infty(G/N_0, \chi)$ s.t. $\forall u \in U(\mathfrak{g}), N \in \mathbb{N}$,

$$|L_u f(kan)| \leq C_{u,N} (1 + |\log(a)|)^{-N} a^{-\rho} \quad (kan \in KAN_0).$$

For (τ, V_τ) a finite dimensional unitary representation of K ,

$$\mathcal{C}(\tau, G/N_0, \chi) := (\mathcal{C}(G/N_0, \chi) \otimes V_\tau)^K$$

$$\mathcal{A}_2(\tau, G/N_0, \chi) := \{f \in \mathcal{C}(\tau, G/N_0, \chi) \mid \dim \mathfrak{Z}f < \infty\}.$$

Theorem (HC, W)

$$\mathcal{A}_2(\tau, G/N_0, \chi) = L_d^2(\tau, G/N_0, \chi).$$

The space is finite dimensional.

Parabolic subgroups

- ▶ $\Sigma = \text{Roots}(\mathfrak{g}, \mathfrak{a})$, $\Sigma^+ := \{\alpha \in \Sigma \mid \mathfrak{g}_\alpha \subset \mathfrak{n}_0\}$, $\Delta \subset \Sigma^+$ simple roots,
- ▶ $W(\mathfrak{a}) = N_K(\mathfrak{a})/Z_K(\mathfrak{a})$.
- ▶ $P_0 := Z_K(A)AN_0$, minimal psg.
- ▶ $\mathcal{P}(A)$: (finite) set of psg's $P \supset A$.
- ▶ $\mathcal{P}_{st} := \{P \in \mathcal{P}(A) \mid P \supset P_0\}$ (standard psg's).
- ▶ For P a psg: Langlands deco: $P = M_P A_P N_P$.

Associated parabolics

For $P, Q \in \mathcal{P}(A)$ define: $P \sim Q$ iff \mathfrak{a}_P and \mathfrak{a}_Q are $W(\mathfrak{a})$ -conjugate. If so,

$$W(\mathfrak{a}_Q \mid \mathfrak{a}_P) := \{T \in \text{Hom}(\mathfrak{a}_P, \mathfrak{a}_Q) \mid \exists w \in W(\mathfrak{a}) : T = w|_{\mathfrak{a}_P}\}$$

Parabolic induction and Whittaker integrals

For $P = M_P A_P N_P \in \mathcal{P}_{st}$, put $\mathcal{A}_{2,P} := \mathcal{A}_2(\tau|_{K_P}, M_P/M_P \cap N_0, \chi|_{M_P \cap N_0})$.

For $\lambda \in \mathfrak{a}_{P\mathbb{C}}^*$, $\operatorname{Re} \lambda >_P 0$,

$$\operatorname{Ind}_P^G(\cdot \otimes -\lambda) : \mathcal{A}_{2,P} \ni \psi \mapsto \mathbf{Wh}(P, \lambda, \cdot, \psi) \in \mathcal{A}_{\text{temp}}(\tau, G/N_0, \chi)$$

Remark

The above **Whittaker integral** is essentially a finite sum of generalized matrix coefficients (defined by Jacquet integrals) of $\operatorname{Ind}_P^G(\sigma \otimes -\lambda \otimes 1)$, with $\sigma \in \widehat{G}_{\text{ds}}$ appearing in $\mathcal{A}_{2,P}$. (Analogue of Eisenstein integral.)

Viewpoint

The Whittaker integral $\mathbf{Wh}(P, \lambda)$ is viewed as a (K -fixed) element of

$$\mathcal{A}_{\text{temp}}(G/N_0, \chi) \otimes \operatorname{Hom}(\mathcal{A}_{2,P}, V_\tau)$$

depending holomorphically on $\lambda \in \mathfrak{a}_{P\mathbb{C}}^*$ in the region $\operatorname{Re} \lambda >_P 0$.

Classical Whittaker functions

Example

- ▶ $G = \mathrm{SL}(2, \mathbb{R})$, $\tau \in \mathrm{SO}(2)^\wedge$
- ▶ $\mathrm{Wh}(P, \lambda, \psi)$ is essentially a classical Whittaker function in the variable $a^{-\alpha} \in (0, \infty)$.
- ▶ satisfies 2nd order ODE on $(0, \infty)$ with regular singularity at 0
- ▶ this ODE has irregular singularity at ∞ ;

For $a^{-\alpha} \rightarrow \infty$

- ▶ generic solution W of ODE:

$$\forall k \geq 0 : |W(a)| \geq a^{-k\alpha} \quad (\text{very fast growth}).$$

- ▶ representation theory selects the special solution

$$\forall k \geq 0 : \mathrm{Wh}(P, \lambda, \psi)(a) = \mathcal{O}(a^{k\alpha}) \quad (\text{very fast decay}).$$

Holomorphic extension

Theorem (W)

$Wh(P, \lambda)$, initially defined for $\operatorname{Re}\lambda >_\rho 0$, extends to entire holom^c function of $\lambda \in \mathfrak{a}_{\mathbb{P}\mathbb{C}}^*$ with values in $C^\infty(G/N_0, \chi) \otimes \operatorname{Hom}(\mathcal{A}_{2,P}, V_\tau)$.

Remark: HC: there exists a merom^c extension, regular on $i\mathfrak{a}_P^*$.

Theorem (\sim): Uniformly tempered estimates

Let $\varepsilon > 0$ be sufftly small. If $u \in U(\mathfrak{g})$ then $\exists C, N, r > 0$ s.t.


$$|Wh(P, \lambda, u; ka)| \leq C(1 + |\lambda|)^N(1 + |\log a|)^N e^{r|\operatorname{Re}\lambda||\log a|} a^{-\rho},$$

for all $k \in K$, $a \in A$, $\lambda \in \mathfrak{a}_{\mathbb{P}\mathbb{C}}^*$ with $|\operatorname{Re}\lambda| < \varepsilon$.

- ▶ Bernstein-Sato type functional equation for Jacquet integrals.
- ▶ Uniformly moderate estimates.
- ▶ Wallach's method of improving estimates along max psg's, with parameters.

C-function, Normalized Whittaker function

- ▶ $W(P, \lambda)$ is \mathfrak{J} -finite,
- ▶ top order asymptotic behavior of \exp^l type along $\text{cl}(A^+)$,
- ▶ rapid decay outside $\text{cl}(A^+)$.



Lemma

Let $P \in \mathcal{P}_{st}$. For $\psi \in \mathcal{A}_{2,P}$, $\text{Re} \lambda \in \mathfrak{a}_P^{*+}$, $m \in M_P$, $a \rightarrow \infty$ in A_P^+ ,

$$\text{Wh}(P, \lambda)(ma)\psi \sim a^{\lambda - \rho_P} [C_P(\lambda)\psi](m),$$

with $C_P(\lambda) \in \text{End}(\mathcal{A}_{2,P})$, merom^c in $\lambda \in \mathfrak{a}_{P\mathbb{C}}^*$ (reg^r for $\text{Re} \lambda \in \mathfrak{a}_P^{*+}$).

Definition (HC) $\text{Wh}^\circ(P, \lambda) := \text{Wh}(P, \lambda) \circ C_P(\lambda)^{-1}$.

Functional equations, Maass-Selberg relations

Lemma (Functional equations: HC)

Let $P, Q \in \mathcal{P}_{st}$, $P \sim Q$. Then for all $s \in W(\mathfrak{a}_Q | \mathfrak{a}_P)$,

$$\mathrm{Wh}^\circ(Q, s\lambda) \circ C_{Q|P}^\circ(s, \lambda) = \mathrm{Wh}^\circ(P, \lambda), \quad (\lambda \in \mathfrak{a}_{P\mathbb{C}}^*),$$

with $C_{Q|P}^\circ(s, \lambda) \in \mathrm{Hom}(\mathcal{A}_{2,P}, \mathcal{A}_{2,Q})$ a uniquely determined meromorphic function of $\lambda \in \mathfrak{a}_{P\mathbb{C}}^*$.

Thm (Maass-Selberg relations, HC)

For all $s \in W(\mathfrak{a}_Q | \mathfrak{a}_P)$, $\lambda \in \mathfrak{a}_{P\mathbb{C}}^*$,

$$C_{Q|P}^\circ(s, -\bar{\lambda})^* \circ C_{Q|P}^\circ(s, \lambda) = I_{\mathcal{A}_{2,P}}$$

In particular, for $\lambda \in i\mathfrak{a}_P^*$, the map $C_{Q|P}^\circ(s, \lambda)$ is **unitary**.

Theorem (HC) $\lambda \mapsto \mathrm{Wh}^\circ(P, \lambda)$ is regular on $i\mathfrak{a}_P^*$.

Fourier transform

Dualized Whittaker function (\sim)

$$\mathrm{Wh}^*(P, \lambda, x) := \mathrm{Wh}^\circ(P, -\bar{\lambda}, x)^* \in \mathrm{Hom}(V_\tau, \mathcal{A}_{2,P}).$$

Fourier transform

For $f \in \mathcal{C}(\tau, G/N_0, \chi)$, $P \in \mathcal{P}_{st}$, $\lambda \in i\mathfrak{a}_P^*$,

$$\mathcal{F}_P f(\lambda) := \int_{G/N_0} \mathrm{Wh}^*(P, \lambda, x) f(x) dx \in \mathcal{A}_{2,P}.$$

Theorem (\sim)

$$\mathcal{F}_P : \mathcal{C}(\tau, G/N_0, \chi) \rightarrow \mathcal{S}(i\mathfrak{a}_P^*) \otimes \mathcal{A}_{2,P},$$

continuous linearly.

Remark: HC proves this for \mathcal{F}_P restricted to $C_c^\infty(\tau, G/N_0, \chi)$.

Proof this follows from the uniformly tempered estimates.

Wave packets

Definition

For $P \in \mathcal{P}_{st}$, $\psi \in \mathcal{S}(i\mathfrak{a}_P^*) \otimes \mathcal{A}_{2,P}$, $x \in G$,

$$\mathcal{W}_P \psi(x) := \int_{i\mathfrak{a}_P^*} \text{Wh}^\circ(P, \lambda, x) \psi(\lambda) d\lambda.$$

Theorem (\sim)

$$\mathcal{W}_P : \mathcal{S}(i\mathfrak{a}_P^*) \otimes \mathcal{A}_{2,P} \rightarrow \mathcal{C}(\tau, G/N_0, \chi)$$

is continuous linear.

Remark: HC proves this for \mathcal{W}_P restricted to $C_c^\infty(i\mathfrak{a}_P^*) \otimes \mathcal{A}_{2,P}$.

Proof requires

- ▶ the uniformly tempered estimates
- ▶ theory of constant term with parameter
- ▶ families of type $\text{II}_{\text{hol}}(\Lambda)$ (as in previous joint work with Carmona and Delorme for reductive symmetric space G/H).

Plancherel formula

If $P, Q \in \mathcal{P}_{st}$, $P \sim Q$ then from the MS rel^s: $\|\mathcal{F}_P f(\lambda)\| = \|\mathcal{F}_Q f(\lambda)\|$.

Plancherel identity (HC)

With suitable normalization of the Lebesgue measures on $i\mathfrak{a}_P^*$,

$$\|f\|_{L^2(\tau, G/N_0, \chi)}^2 = \sum_{P \in \mathcal{P}_{st}/\sim} \|\mathcal{F}_P f\|_{L^2(i\mathfrak{a}_P^*) \otimes \mathcal{A}_{2,P}}^2,$$

for f in the linear span $\mathcal{W} \subset \mathcal{C}(\tau, G/N_0, \chi)$ of the wavepackets $\mathcal{W}_Q(\psi)$, for $Q \in \mathcal{P}_{st}$, and $\psi \in C_c^\infty(i\mathfrak{a}_Q^*) \otimes \mathcal{A}_{2,Q}$.

Problem of the final step: Is \mathcal{W} dense in $L^2(\tau, G/N_0, \chi)$?

Theorem (\sim)

Yes! More precisely, for $f \in \mathcal{C}(\tau, G/N_0, \chi)$ we have

$$f = \sum_{P \in \mathcal{P}_{st}/\sim} \mathcal{W}_P \circ \mathcal{F}_P(f).$$

Series expansion

Strategy for the final step: use Paley-Wiener shift argument and residue calculus as known from the theory of symmetric spaces (previous joint work with Schlichtkrull).

Let $P = P_0$ be minimal. Then $\text{Wh}(P, \lambda) \in C^\infty(\tau, \mathbf{G}, \chi) \otimes \mathcal{A}_{2,P}^*$ is holomorphic in $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$. The function is \mathfrak{Z} -finite, hence satisfies a cofinite system of differential equations, which has regular singularities at infinity in the direction of A^+ .

Expansion at infinity

$$\text{Wh}(P, \lambda) = \sum_{s \in W(\mathfrak{a})} \text{Wh}_+(P, s\lambda) C_{P|P}(s, \lambda)$$

where $\text{Wh}_+(P, \lambda) \in C^\infty(\tau, \mathbf{G}, \chi) \otimes \mathcal{A}_{2,P}^*$ is merom^c in $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$, and

$$\text{Wh}_+(P, \lambda)(a) = a^{\lambda - \rho} \sum_{\mu \in \mathbb{N}\Delta} a^{-\mu} \Gamma_\mu(\lambda), \quad (a \in A),$$

with $\Gamma_\mu(\lambda) \in \text{Hom}(\mathcal{A}_{2,P}, V_\tau)$ meromorphic, $\Gamma_0(\lambda)(\psi) = \psi(e)$.

Fourier inversion

Key theorem (\sim)

$$f(x) = \mathcal{T}_\eta(f)(x) := |W(\mathfrak{a})| \int_{i\mathfrak{a}^* + \eta} \text{Wh}_+(P, \lambda, x) \mathcal{F}_P f(\lambda) d\lambda,$$

$\forall f \in C_c^\infty(\tau, G/N_0, \chi), \forall x \in G$, provided $\eta \in \mathfrak{a}^*$, $\eta \gg_{\bar{P}} 0$.

NB: For generic $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$, the function $\text{Wh}_+(P, \lambda)$ is globally def^d on X , but may exhibit super exp^l growth in directions diff^t from $\text{cl}(A^+)$.

Ideas of proof

- ▶ $\mathcal{T}_\eta : C_c^\infty(\tau, G/N_0, \chi) \rightarrow C^\infty(\tau, G/N_0, \chi)$.
- ▶ $\exists D \in \mathfrak{Z} : DT_\eta = DT_0 = D \circ \mathcal{W}_P \circ \mathcal{F}_P$.
- ▶ By PW shift $\eta \rightarrow \infty$ in \bar{P} -dominant direction: $\text{supp} DT_\eta f \subset K \text{supp} f$.
 $\implies \text{rad}(DT_\eta)$ is a differential operator on A

Proof of Fourier inversion

Ideas of proof:

- ▶ $\text{rad}(DT_\eta)$ differential operator commuting with $\text{rad}(3)$
- ▶ By asymptotic analysis along A_P^+ : $DT_\eta = D$ on $C_c^\infty(\tau, G/N_0, \chi)$.
- ▶ By Holmgren's uniqueness theorem for analytic PDO:

$$DT_\eta f = Df \implies D(\mathcal{T}_\eta f - f) = 0 \implies \mathcal{T}_\eta f - f = 0.$$

Residual kernels

By Fourier inversion, if $f \in C_c^\infty(\tau, G/N_0, \chi)$, $x \in G$,

$$f(x) = |W(\mathfrak{a})| \int_{i\mathfrak{a}^* + \eta} \text{Wh}_+(P, \lambda, x) \mathcal{F}_P f(\lambda) d\lambda.$$

Shifting η towards zero and **organizing residues**, one gets

$$f(x) = \sum_{Q \in \mathcal{P}_{st}} [W : N_W(\mathfrak{a}_Q)] t(Q) T_Q^t f(x),$$

where

$$T_Q^t f(x) = \int_{i\mathfrak{a}_Q^* + \varepsilon_Q} \int_{G/N_0} K_Q^t(\lambda, x, y) f(y) dy d\lambda_Q.$$

- ▶ $\varepsilon_Q \in \mathfrak{a}_Q^{*+}$ sufficiently close to 0.
- ▶ $t : \mathcal{P}_{st} \rightarrow [0, 1]$ is a weight function describing a certain organization of residue shifts.

Conclusion

Theorem (\sim)

$$K_Q^t(\lambda, x, y) = \text{Wh}^\circ(Q, \lambda)(x) \circ \text{Wh}^*(Q, \lambda)(y) = K_{\mathcal{W}_Q \circ \mathcal{F}_Q}.$$

This identification relies on the Maass-Selberg relations. These also imply that the functions $\lambda \mapsto K_Q^t(\lambda, x, y)$ are regular on $i\mathfrak{a}_Q^*$, hence we may let $\varepsilon_Q \rightarrow 0$ and then:

Plancherel formula

$$f(x) = \sum_{Q \in \mathcal{P}_{st}} [W : N_W(\mathfrak{a}_Q)] t(Q) \mathcal{W}_Q \mathcal{F}_Q f(x).$$

- ▶ $[W : N_W(\mathfrak{a}_Q)] t(Q)$ gives the weight by which Q contributes to its class in \mathcal{P}_{st} / \sim .


Bonus:

Bonus: Paley-Wiener theorem

Definition

A function $f \in \mathcal{C}(\tau, G/N_0, \chi)$ is said to be *cone supported* (notation \mathcal{C}_{cs}) if $\exists H_0 \in \mathfrak{a}$ s.t.

$$\text{supp} f \subset K \exp(H_0 - \mathfrak{a}^{++}) N_0.$$



Lemma

If $f \in \mathcal{C}_{cs}(\tau, G/N_0, \chi)$, then $\forall u \in U(\mathfrak{g}) \forall m > 0 \exists C > 0$:

$$\|L_u f(ka)\| < C e^{-m|\log a|} \quad (\forall k \in K, a \in A).$$

Paley-Wiener theorem

Let $P = P_0$ (minimal). Then ${}^u\mathcal{F}_P$ (unnormalized) is injective on $\mathcal{C}_{cs}(\tau, G/N_0, \chi)$. The image of this space under ${}^u\mathcal{F}_P$ equals the space $\text{PW}(\chi, \tau)$ of holomorphic functions $\varphi : \mathfrak{a}_{\mathbb{C}}^* \rightarrow \mathcal{A}_{2,P}$ satisfying

- ▶ certain estimates of Paley–Wiener type;
- ▶ relations of Arthur–Campoli type.

Thank you

Arthur–Campoli type relations

More precisely, the definition of the PW space is as follows.

Definition Paley–Wiener space

$\text{PW}(\chi, \tau)$ is the space of holomorphic functions $\varphi : \mathfrak{a}_{\mathbb{C}}^* \rightarrow \mathcal{A}_{2,P}$ satisfying

- ▶ $\exists R > 0: \forall \lambda_0 \in \mathfrak{a}_{\mathbb{C}}^*, \forall N \in \mathbb{N}, \exists C > 0$ s.t.

$$|\varphi(\lambda)| \leq C(1 + \|\lambda\|)^{-N} e^{R\|Re\lambda\|} \quad (\lambda \in \lambda_0 - \mathfrak{a}_{\mathbb{C}}^{*+}).$$

- ▶ For all finite collections $\lambda_i \in \mathfrak{a}_{\mathbb{C}}^*, u_i \in \mathcal{S}(\mathfrak{a}^*), \xi_i \in \text{Hom}(V_{\tau}, \mathcal{A}_{2,P})^*, 1 \leq i \leq N,$

$$\sum_{i=1}^N \langle \xi_i, \partial_{u_i} \text{Wh}^*(P, \cdot)(\lambda_i) \rangle = 0 \quad \implies \quad \sum_{i=1}^N \langle \xi_i, \partial_{u_i} \varphi(\lambda_i) \rangle = 0.$$

C-functions, Maass-Selberg relations

Thm (asymptotic behavior, HC)

Let $P \in \mathcal{P}_{st}$, $\psi \in \mathcal{A}_{2,P}$. If $Q \in \mathcal{P}_{st}$, $Q \sim P$, then for $\lambda \in i\mathfrak{a}_P^*$ generic, $m \in M_Q$, $a \rightarrow \infty$ in A_Q^+ ,

$$Wh(P, \lambda)(ma)\psi \sim \sum_{s \in W(\mathfrak{a}_Q | \mathfrak{a}_P)} a^{s\lambda - \rho_Q} [C_{Q|P}(s, \lambda)\psi](m),$$

with $C_{Q|P}(s, \lambda) \in \text{Hom}(\mathcal{A}_P, \mathcal{A}_Q)$ meromorphic in $\lambda \in \mathfrak{a}_{PC}^*$.

If $R \in \mathcal{P}(A) \setminus \mathcal{P}_{st}$ then $Wh(P, \lambda)(ma) = o(a^{-\rho_R})$ for $a \rightarrow \infty$ in A_R^+ .

Thm (Maass-Selberg relations, HC)

For all $s \in W(\mathfrak{a}_Q | \mathfrak{a}_P)$, $\lambda \in i\mathfrak{a}_P^*$,

$$C_{Q|P}^\circ(s, \lambda) := C_{Q|P}(s, \lambda)C_{P|P}(1, \lambda)^{-1}$$

is *unitary* $\mathcal{A}_{2,P} \rightarrow \mathcal{A}_{2,Q}$.

Discrete part of Fourier transform

Discrete part of Fourier transform

If $\text{rk}(\mathfrak{k}) = \text{rk}(\mathfrak{g})$ one has $\mathfrak{a}_G = \{0\}$ and \mathcal{F}_G is given by the (finite rank) L^2 -orthogonal projection

$$\mathcal{C}(\tau, \mathbf{G}/\mathbf{N}_0, \chi) \rightarrow \mathcal{A}_{2,\mathbf{G}} = L_d^2(\tau, \mathbf{G}/\mathbf{N}_0, \chi).$$

HC's fundamental theorem

- ▶ $\Sigma = \text{Roots}(\mathfrak{g}, \mathfrak{a})$, $\Sigma^+ := \{\alpha \in \Sigma \mid \mathfrak{g}_\alpha \subset \mathfrak{n}_0\}$, $\Delta \subset \Sigma^+$ simple roots.
- ▶ $P_0 := Z_K(A)AN_0$, minimal psg; $\mathcal{P}(A)$: (finite) set of psg's $P \supset A$.
- ▶ $\mathcal{P}_{st} := \{P \in \mathcal{P}(A) \mid P \supset P_0\}$ (standard psgs).

Let $P \in \mathcal{P}_{st}$, Langlands deco: $P = M_P A_P N_P$.

Then $\bar{P}N_0$ is open dense in G .

Harish-Chandra's Thm 1

Let $u \in \mathcal{D}'(G)$ be such that

$$L_{\bar{n}}u = u, \quad R_n u = \chi^{-1}(n)u, \quad (\bar{n} \in \bar{N}_P, n \in N_0)$$

If χ is regular and $u|_{\bar{P}N_0} = 0$ then $u = 0$.

Ref. for proof also: J.A.C. Kolk, V.S. Varadarajan, Indag. Math. 1996.