

# Anomalies and the Atiyah-Singer index theorem

Author:  
Andries Salm

Supervisor  
Gil Cavalcanti

Supervisor  
Eric Laenen

June 11, 2018



Universiteit Utrecht

Nikhef

*This line will be changed later.*

## Introduction

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

## Contents

<b>1</b>	<b>Clifford Algebras</b>	<b>6</b>
1.1	Definitions . . . . .	6
1.2	Graded Clifford bundles . . . . .	10
1.3	The adjoint of a Dirac operator . . . . .	14
1.4	The Weitzenböck formula . . . . .	16
<b>2</b>	<b>Calculating anomalies using Feynman diagrams</b>	<b>22</b>
2.1	Revisit of Noethers theorem . . . . .	22
2.2	Chiral currents in triangle diagrams . . . . .	24
2.2.1	Dimensional regularization . . . . .	26
2.2.2	Regularization using Pauli-Villars method . . . . .	30
2.3	Other triangle diagrams . . . . .	32
2.4	Anomalies in box- and pentagon-diagrams . . . . .	33
2.5	Non-Abelian Anomalies in Feynman diagrams . . . . .	38
<b>3</b>	<b>Calculating anomalies using path integrals</b>	<b>41</b>
3.1	The Fujikawa method . . . . .	41
3.2	The covariant anomaly . . . . .	46
3.3	The Bardeen anomaly . . . . .	48
3.4	The consistent anomaly . . . . .	52
3.5	Mathematical interpretation of the Fujikawa method . . . . .	55
<b>4</b>	<b>Smoothing operators and Heat kernels</b>	<b>58</b>
4.1	Definitions . . . . .	58
4.2	Examples of heat kernels . . . . .	60
4.3	Uniqueness of the heat kernel . . . . .	64
4.4	The formal solution of the heat kernel . . . . .	66
4.5	The existence of the heat kernel . . . . .	70
<b>5</b>	<b>Traces and the Index of a Dirac operator</b>	<b>79</b>
5.1	Traceclass operators . . . . .	79
5.2	Spectral theory of generalized Laplacians . . . . .	86
5.3	The McKean-Singer formula . . . . .	90
5.4	The trace and the Clifford action . . . . .	92

<b>6</b>	<b>Characteristic classes</b>	<b>96</b>
6.1	Invariant polynomials . . . . .	96
6.2	Examples of characteristic classes . . . . .	102
6.3	Characteristic classes of Clifford bundles . . . . .	103
<b>7</b>	<b>Symbol Calculus</b>	<b>105</b>
7.1	Definitions . . . . .	105
7.2	Getzler filtration . . . . .	109
7.3	The symbol of the heat kernel . . . . .	115
7.4	Atiyah-Singer Index theorem . . . . .	118
<b>8</b>	<b>Summary and Outlook</b>	<b>122</b>
<b>9</b>	<b>Glossary: Differential geometry for physicists</b>	<b>123</b>
9.1	Sections and fields . . . . .	123
9.2	Constructing vector bundles . . . . .	123
9.3	Derivatives and curvature . . . . .	123
9.4	Topology of vector bundles . . . . .	128
<b>10</b>	<b>Field theory in terms of differential geometry</b>	<b>131</b>
<b>11</b>	<b>Appendix: Source codes</b>	<b>132</b>
11.1	Calculation of Bardeen Anomaly . . . . .	132

# 1 Clifford Algebras

In the study of spin- $\frac{1}{2}$  particles, physicists use gamma matrices. They are defined as

$$\begin{aligned}\gamma_0 &= \begin{pmatrix} -i & 0 & & \\ 0 & -i & & \\ & & i & 0 \\ & & 0 & i \end{pmatrix} & \gamma_1 &= \begin{pmatrix} & 0 & -i & \\ & -i & 0 & \\ 0 & i & & \\ i & 0 & & \end{pmatrix} \\ \gamma_2 &= \begin{pmatrix} & & 0 & -1 \\ & & 1 & 0 \\ 0 & 1 & & \\ -1 & 0 & & \end{pmatrix} & \gamma_3 &= \begin{pmatrix} & & -i & 0 \\ & & 0 & i \\ i & 0 & & \\ 0 & -i & & \end{pmatrix}\end{aligned}\quad (1.1)$$

and they satisfy the commutation relation

$$\{\gamma_\mu, \gamma_\nu\} := \gamma_\mu \gamma_\nu + \gamma_\nu \gamma_\mu = \begin{cases} 2 & \text{if } \mu = \nu = 0 \\ -2 & \text{if } \mu = \nu \text{ and } \mu, \nu \in \{1, 2, 3\} \\ 0 & \text{else} \end{cases} \quad (1.2)$$

The algebra these matrices is one of the first examples of an object now called Clifford algebra. In this chapter we give a rigorous definition of this algebra and we derive its basic properties. Using this algebra we define a differential operator called the Dirac operator. We also introduce the physics notation for dealing with Clifford algebras.

In the end, we prove the Weitzenböck formula. This equation relates the square of a Dirac operator with the Laplacian and we show that the difference can be given in terms of the curvature and the gamma matrices. For this we need an explicit expression for the adjoint of the Dirac operator and the connection. In the third paragraph we perform these calculations and we show that the Dirac operator is self-adjoint.

The material covered in this section is standard. For more information see Roe [1998]

## 1.1 Definitions

**Definition 1.1.** Let  $V$  be a vector space with a symmetric 2-form  $g$ . A **Clifford algebra** for  $V$  is an unital algebra  $A$  such that the following holds:

1. There exists a linear map  $\phi: V \rightarrow A$  such that  $\phi(v)^2 = -g(v, v) \text{Id}$  for all  $v \in V$ .

2.  $A$  satisfies the universal property. That is, if there exists another map  $\phi': V \rightarrow A'$  that have the same property of  $\phi$ , then there exists a unique algebra homomorphism  $A \rightarrow A'$  such that the following diagram commutes:

$$\begin{array}{ccc} V & \xrightarrow{\phi} & A \\ & \searrow \phi' & \downarrow \\ & & A' \end{array}$$

Given a vector space  $V$  and a bilinear symmetric map  $g$  we can construct a Clifford algebra. Indeed, consider the tensor algebra

$$T(V) = \bigoplus_{k=0}^{\infty} V \otimes V \otimes \dots \otimes V \quad (1.3)$$

and take the quotient by the ideal that is generated by  $v \otimes v + g(v, v) \text{Id}$ . Denote this quotient as  $\text{Cl}(V)$  or  $\text{Cl}(V, g)$ . By construction it satisfies the first part of Definition 1.1. The universal property follows from the universal property of tensor algebras. It states that for all linear maps  $\phi: V \rightarrow A$  there exists a unique algebra homomorphism  $\phi': T(V) \rightarrow A$  such that

$$\begin{array}{ccc} V & \longrightarrow & T(V) \\ & \searrow \phi & \downarrow \phi' \\ & & A \end{array}$$

commutes. Now assume that  $\phi(v)^2 = -g(v, v) \text{Id}$  for all  $v \in V$ . This relation extends on  $\phi'$  to

$$\phi'(v \otimes v + g(v, v) \text{Id}) = \phi'(v)^2 + g(v, v) \phi'(\text{Id}) = 0 \quad (1.4)$$

So  $\phi'$  factors over the ideal generated by  $v \otimes v + g(v, v) \text{Id}$  and so  $\phi'$  is a unique algebra homomorphism between  $\text{Cl}(V)$  and  $A$ . Hence,

**Proposition 1.2.** *For any vector space  $V$  with a bilinear symmetric map, there exists a Clifford algebra and it is unique.*

The requirement in Definition 1.1 that  $\phi(v)^2 = -g(v, v) \cdot \text{Id}$  for all  $v \in V$  is equivalent to the requirement that  $\phi(u)\phi(v) + \phi(v)\phi(u) = -2g(u, v) \cdot \text{Id}$  for all  $u, v \in V$ .

This can be easily seen if we consider  $\phi(u+v)^2$ . By linearity this equals  $\phi(u)^2 + \phi(u)\phi(v) + \phi(v)\phi(u) + \phi(v)^2$  and so the anti-commutator between  $\phi(u)$  and  $\phi(v)$  is  $\phi(u+v)^2 - \phi(u)^2 - \phi(v)^2$ . However, by the commutation relation it follows that

$$\phi(u)\phi(v) + \phi(v)\phi(u) = -(g(u+v, u+v) - g(u, u) - g(v, v)) \cdot \text{Id} \quad (1.5)$$

$$= -2g(u, v) \cdot \text{Id} \quad (1.6)$$

Hence these two conditions are equivalent.

**Example 1.3** (Gamma matrices). Consider the vector space  $V = \mathbb{R}^4$  and equip it with the Minkowski metric  $\eta$ . In the basis  $(t, x, y, z)$  the Minkowski metric is given by,  $\eta_{tt} = -1$ ,  $\eta_{xx} = \eta_{yy} = \eta_{zz} = 1$  and  $\eta_{\mu\nu} = 0$  else. Hence, the gamma matrices satisfy the relation  $\gamma_\mu \gamma_\nu + \gamma_\nu \gamma_\mu = -2\eta_{\mu\nu}$ . and so  $\gamma_\mu^2 = -\eta_{\mu\mu} \cdot \text{Id}_{\mathbb{C}^4}$ . The unital matrix subalgebra  $A$  generated by the gamma matrices is a Clifford algebra. Indeed, the map  $\text{Cl}(V, \eta) \rightarrow A$  that is defined by  $t^i \cdot x^j \cdot y^k \cdot z^l \mapsto \gamma_t^i \circ \gamma_x^j \circ \gamma_y^k \circ \gamma_z^l$  for all  $i, j, k, l \in \mathbb{N}$  is an isomorphism between two algebras. It is a well-defined linear map and by the definition of  $A$  it is surjective. It is a homomorphism, because both sides satisfy the same commutation relations. We only need to show that it is injective. For this we need to work out all matrix multiplications and this is left for the reader.

In field theory the gamma matrices act on  $\text{spin-}\frac{1}{2}$  fermions which are represented by fields. Mathematically particle fields are sections of a certain vector bundle. So to formalize  $\text{spin-}\frac{1}{2}$  fermions we need to let the Clifford action act on section

**Definition 1.4.** Let  $(M, g)$  be a (pseudo)-Riemannian manifold. The vector bundle  $S \rightarrow M$  is a **bundle of Clifford modules** if there exists a smooth bundle map  $\text{Cl}(TM, g) \otimes \mathbb{C} \times S \rightarrow S$  which makes each fiber  $S_p$  a left-module over  $\text{Cl}(T_p M, g) \otimes \mathbb{C}$ .

**Example 1.5** (Spin- $\frac{1}{2}$  particles). A  $\text{spin-}\frac{1}{2}$  particle is represented by a section over the trivial bundle  $\mathbb{R}^4 \times \mathbb{C}^4 \rightarrow \mathbb{R}^4$ . Clearly, the gamma matrices forms a left-module on  $\mathbb{C}^4$ . So  $\mathbb{R}^4 \otimes \mathbb{C}^4$  is a bundle of Clifford modules.

For a bundle of Clifford modules  $S \rightarrow M$ , we define  $\gamma: \Gamma(TM) \rightarrow \Gamma(\text{End}(S))$  as the composition of the left-module action and the map  $\phi$  from Definition 1.1. In local coordinates  $e_\mu \in \Gamma(TM)$ , this map is given by  $\gamma(e_\mu)s = \phi(e_\mu) \cdot s$  for all sections  $s \in \Gamma(S)$ . If we use the shorthand  $\gamma_\mu = \gamma(e_\mu)$ , we see that  $\gamma_\mu$  extends the gamma matrices from Equation 1.1 to any bundle of Clifford modules. We call  $\gamma: \Gamma(TM) \rightarrow \Gamma(\text{End}(S))$  the **Clifford action**.

Given a bundle of Clifford modules  $S \rightarrow M$ , a vector bundle  $E \rightarrow M$  and a section  $A \in \Gamma(TM \otimes S \otimes E)$  we formally define the **Feynman slash**  $\not{A}$  of  $A$  as follows: It is the element  $\not{A} \in \Gamma(S \otimes E)$  that is given by the composition of the following maps:

$$\Gamma(TM \otimes S \otimes E) \xrightarrow{\phi} \Gamma(\text{Cl}(TM) \otimes S \otimes E) \longrightarrow \Gamma(S \otimes E) \quad (1.7)$$

$$A = \sum_{\mu} A^{\mu} \otimes e_{\mu} \longmapsto \sum_{\mu} A^{\mu} \otimes \gamma_{\mu} \longmapsto \not{A} = \gamma_{\mu} A^{\mu}.$$

The second map denotes the multiplication defined by the left-module. We used  $\{e_{\mu}\}$  as a local basis on  $TM$ . We only need it for comparison with the definition physicists use and we see that they indeed coincide.

In physics there is an operator called the Dirac operator. For the generalization of this operator we need to consider the interplay between a connection and the Clifford action. In the next definition we give explicit requirements:

**Definition 1.6.** A **Clifford bundle** is a bundle of Clifford modules  $S$  over a (pseudo)-Riemannian manifold  $(M, g)$  equipped with a Hermitian metric and compatible connection such that

1. The Clifford action is skew adjoint. That is, for all  $p \in M$ ,  $v \in T_p M$  and  $s_1, s_2 \in S_p$ , we have

$$\langle \gamma(v) \cdot s_1, s_2 \rangle + \langle s_1, \gamma(v) s_2 \rangle = 0 \quad (1.8)$$

2. The connection on  $S$  is compatible with the Levi-Civita connection on  $M$ . So for all  $u, v \in \Gamma(TM)$  and  $s \in \Gamma(S)$ , we have

$$\nabla_u \gamma(v) s = \gamma(\nabla_u v) s + \gamma(v) \nabla_u s \quad (1.9)$$

**Example 1.7** (Spin- $\frac{1}{2}$  particles). Note that the gamma matrices from Equation 1.1 are skew adjoint under the Euclidean metric. Also the compatibility condition reduces to the Leibniz rule for a flat connection. Hence, if we equip  $\mathbb{R}^4 \times \mathbb{C}^4 \rightarrow \mathbb{R}^4$  with the Euclidean metric of  $\mathbb{C}^4$  and the flat connection, we get that  $\mathbb{R}^4 \times \mathbb{C}^4$  is a Clifford bundle.

**Definition 1.8.** Let  $S \rightarrow (M, g)$  be a Clifford bundle with compatible connection  $\nabla$  and the Clifford action  $\gamma$ . Interpret  $g$  as the isomorphism  $g: \Gamma(TM) \rightarrow \Gamma(T^*M)$  which is given by  $v \mapsto g(v, \cdot)$ . The differential operator  $D$  is a **Dirac operator** if it is the composition of the following maps:

$$\Gamma(S) \xrightarrow{\nabla} \Gamma(T^*M \otimes S) \xrightarrow{g^{-1}} \Gamma(TM \otimes S) \xrightarrow{\gamma} \Gamma(S) \quad (1.10)$$

Following the steps above we can compute the Dirac operator in local coordinates. Denote  $\{e_\mu\}$  as a local basis of  $TM$  and  $s$  as a section of  $S$ . Let  $e_\mu^b$  be the dual of  $e^\mu$ , i.e.  $e_\mu^b(e^\nu) = \delta_{\mu\nu}$ . By orthonormality it follows that  $\sum_\mu e_\mu e_\mu^b = \text{Id}$  and

$$\nabla s = \sum_\mu \nabla_\mu s \otimes e_\mu^b \quad (1.11)$$

Under  $g^{-1}$  this maps to  $\sum_\mu \nabla_\mu s \otimes e_\mu$  and so the Dirac operator is locally given by

$$\gamma^\mu \nabla_\mu s \quad (1.12)$$

In the example of the  $\text{spin-}\frac{1}{2}$  particles we use the flat connection  $\nabla_\mu = \partial_\mu$ . Then the Dirac operator is given by  $\partial_\mu \gamma^\mu$  which is the expression normally used in quantum field theory.

## 1.2 Graded Clifford bundles

For  $\text{spin-}\frac{1}{2}$  Dirac fermions, we distinguish left-handed and right-handed particles. This chirality is related to the  $\pm 1$  eigenvalues of the operator

$$\gamma_5 := -\gamma_0 \gamma_1 \gamma_2 \gamma_3, \quad (1.13)$$

That is, if  $\gamma_5 \psi = \psi$ , then we call  $\psi$  right handed and if  $\gamma_5 \psi = -\psi$  we call  $\psi$  left-handed. We see that  $\gamma_5$  defines a grading on  $\mathbb{R}^4 \times \mathbb{C}^4$ . In general we define the grading as follows.

**Definition 1.9.** Let  $S$  be a Clifford bundle. We say that  $S$  is a **graded Clifford bundle** if  $S$  can be decomposed into  $S^+ \oplus S^-$  such that

1. The Clifford action  $\gamma$  maps  $\Gamma(TM \otimes S^\pm)$  to  $\Gamma(S^\mp)$ .
2. The metric and the connection of  $S$  respects the grading.

In the example of  $\mathbb{R}^4 \times \mathbb{C}^4$  the operator  $\gamma_5$  is fully determined by how it acts on left- resp. right-handed particles. It acts by  $+1$  on right-handed fields and by  $-1$  on left-handed field. This can be generalized to any graded Clifford bundle.

**Definition 1.10.** *Let  $S$  be a graded Clifford bundle. We define the **grading operator**  $\gamma_5: \Gamma(S) \rightarrow \Gamma(S)$  by*

$$\begin{aligned}\gamma_5|_{S^+} &= \text{Id} \\ \gamma_5|_{S^-} &= -\text{Id}\end{aligned}\tag{1.14}$$

The grading operator anti-commutes with the Clifford action. Indeed if  $S = S^+ \oplus S^-$  is a graded Clifford bundle and  $s \in \Gamma(S^\pm)$ , then for all  $v \in \Gamma(TM)$  the section  $\gamma(v)s$  is an element of  $\Gamma(S^\mp)$  and hence

$$\gamma_5 \gamma(v)s = \mp \gamma(v)s = -\gamma(v)\gamma_5 s\tag{1.15}$$

Even more, the Dirac operator anti-commutes with the grading operator. This can be easily seen if we use local coordinates and write  $D = \sum_\mu \gamma_\mu \nabla_\mu$ .

Gradings on Clifford bundles are not unique. However, there is a canonical method to induce a grading.

**Definition 1.11.** *Let  $S \rightarrow (M, g)$  be a Clifford bundle on a  $2n$  dimensional manifold and let  $\{e_\mu\}$  be a local positively oriented orthonormal basis of  $TM$ . The element*

$$\omega = i^n \gamma_1 \cdot \dots \cdot \gamma_{2n} \in \Gamma(\text{End}(S))\tag{1.16}$$

*is called the **canonical grading operator**.*

**Lemma 1.12** (Roe [1998], Remark 4.4). *The canonical grading operator  $\omega$  from Definition 1.11 does not depend on the choice of local orthonormal frame and hence it is globally defined.*

*Proof.* Let  $\{e^\mu\}$  and  $\{\tilde{e}^\mu\}$  be two orthonormal local frames of  $TM$  with identical orientation. Denote  $\gamma_\mu$  and  $\tilde{\gamma}_\mu$  as the Clifford action on  $e_\mu$  resp.  $\tilde{e}_\mu$ . We expand  $\tilde{e}^\mu$

as  $A_\nu^\mu e^\nu$ . In local coordinates the canonical grading operator  $\tilde{\omega}$  w.r.t. the basis  $\{\tilde{e}^\mu\}$  equals

$$\tilde{\omega} := i^n \tilde{\gamma}_1 \cdot \dots \cdot \tilde{\gamma}_{2n} = i^n A_1^{\mu_1} \dots A_{2n}^{\mu_{2n}} \gamma_{\mu_1} \dots \gamma_{\mu_{2n}} \quad (1.17)$$

We view  $A$  as an orthogonal matrix on  $\{\tilde{e}^\mu\}$ . In the case that  $\mu_1 = \mu_2$  it follows

$$\sum_{\mu_1} A_1^{\mu_1} A_2^{\mu_2} = (AA^T)_{12} = \text{Id}_{12} = 0 \quad (1.18)$$

This shows that all  $\mu_i$ 's in Equation 1.17 must be unique. If  $S_{2n}$  is the set of all permutations of  $2n$ -elements, then  $\tilde{\omega}$  equals

$$\tilde{\omega} = i^n \sum_{\sigma \in S_{2n}} A_{\sigma(1)}^{\mu_{\sigma(1)}} \dots A_{\sigma(2n)}^{\mu_{\sigma(2n)}} \gamma_{\mu_{\sigma(1)}} \dots \gamma_{\mu_{\sigma(2n)}} \quad (1.19)$$

$$= i^n \sum_{\sigma \in S_{2n}} \text{sgn}(\sigma) A_{\sigma(1)}^{\mu_{\sigma(1)}} \dots A_{\sigma(2n)}^{\mu_{\sigma(2n)}} \gamma_1 \dots \gamma_{2n} \quad (1.20)$$

$$= \det(A) \cdot \omega \quad (1.21)$$

Because  $A \in SO(2n)$  the determinant of  $A$  equals one and  $\tilde{\omega}$  equals the canonical grading operator w.r.t. the basis  $\{e^\mu\}$ .  $\square$

In the next proposition we show that the canonical grading operator  $\omega$  indeed defines a graded Clifford bundle and so we see  $\omega$  as the generalization of  $\gamma_5$ .

**Proposition 1.13** (Roe [1998], Page 142). *Let  $S \rightarrow (M, g)$  be a Clifford bundle on a  $2n$  dimensional manifold and let  $\omega$  be the canonical grading operator. Then  $\omega^2 = \text{Id}$  and the  $\pm 1$  eigenspaces of  $\omega$ ,  $S^\pm$ , are subbundles of  $S$ , therefore decomposing it as graded Clifford bundle  $S = S^+ \oplus S^-$ .*

*Proof.* The square of  $\omega$  can be easily calculated and is indeed equal to one. Therefore we can split  $S$  into  $S^+ \oplus S^-$  where  $S^\pm$  are the  $\pm 1$  eigenspaces of  $\omega$ . Also, the commutation identity

$$\omega \gamma + \gamma \omega = 0 \quad (1.22)$$

can be easily calculated in a suitable basis. This proves the first part of Definition 1.9.

To show that  $g$  respects the grading we assume that  $v_+ \in \Gamma(S^+)$  and  $v_- \in \Gamma(S^-)$ . By the skew-adjointness of the Clifford action and  $\omega v_{\pm} = \pm v_{\pm}$  we can calculate  $g(v_+, v_-)$ .

$$\begin{aligned} g(v_+, v_-) &= g(\omega v_+, v_-) = i^n g(\gamma^1 \dots \gamma^{2n} v_+, v_-) \\ &= (-i)^n g(v_+, \gamma^{2n} \dots \gamma^1 v_-) = (i)^n g(v_+, \gamma^1 \dots \gamma^{2n} v_-) \\ &= g(v_+, \omega v_-) = -g(v_+, v_-) \end{aligned} \quad (1.23)$$

This shows  $g(S^+, S^-) = 0$  and hence  $g$  respects the grading.

To show that the connection on  $S$  respects the grading we need to pick a suitable coordinate system on  $M$ . Fix  $p \in M$  and pick the local coordinate frame  $\{\frac{\partial}{\partial x^\mu}\}$  that follows from the exponential map  $\exp_p: T_p M \rightarrow M$ . This frame is called the **Riemannian normal coordinate system** and it has the property that at  $p$

$$\nabla \frac{\partial}{\partial x^\mu} = 0$$

where  $\nabla$  denote the Levi-Civita connection and  $\frac{\partial}{\partial x^\mu}$  is a basis element in  $\Gamma(TM)$ . Let  $s \in \Gamma(S)$ ,  $v \in \Gamma(TM)$  and use  $\nabla$  also for the connection on  $S$ . Using the compatibility property of the connection we get

$$\nabla_v(\omega s) = i^n \nabla(\gamma^1 \dots \gamma^{2n} s) \quad (1.24)$$

$$= i^n \gamma(\nabla_v(\partial/\partial x^1)) \cdot \gamma^2 \dots \gamma^{2n} s + i^n \gamma^1 \nabla_v(\gamma^2 \dots \gamma^{2n} s) \quad (1.25)$$

$$= i^n \gamma^1 \nabla_v(\gamma^2 \dots \gamma^{2n} s) \quad (1.26)$$

$$= \dots = \omega \nabla_v s \quad (1.27)$$

This shows that  $\nabla$  also respects the grading.  $\square$

**Remark 1.14.** Unless not stated otherwise we assume that a Clifford bundle is canonically graded.

**Remark 1.15.** If  $\gamma$  is the Clifford action and  $\{e^\mu\}$  a basis of  $TM$  we simplify  $\gamma(e^\mu)$  into  $\gamma_\mu$ . However, if we explicitly<sup>1</sup> write  $\gamma_5$ , then it does not mean  $\gamma(e_5)$ , but it means the grading operator. This might be confusing, but it is standard notation in physics.

**Remark 1.16.** In higher dimensions physicists mostly use the notation  $\gamma_{n+1}$  where  $n$  is the dimension of the spacetime. However, when we consider higher dimensions in physics, we will use the Veltman-'t Hooft regularization. There it is custom to use  $\gamma_5$  instead of  $\gamma_{n+1}$ . To prevent confusion we never use  $\gamma_{n+1}$ , but use  $\gamma_5$ .

<sup>1</sup>In  $\gamma_0 \dots \gamma_k$  the operator  $\gamma_5$  is not explicitly written so we don't refer here to the grading operator.

### 1.3 The adjoint of a Dirac operator

In this paragraph we will compute the formal adjoint of a Dirac operator  $D$ . Since  $D$  is the composition of the Clifford action  $\gamma$  and the covariant derivative  $\nabla$ . Therefore we compute the adjoint of  $\gamma$  and  $\nabla$  separately. These will be calculated in the next two lemmas. We conclude this paragraph by showing that  $D$  is self-adjoint.

**Lemma 1.17.** *Let  $E \rightarrow M$  be a vector bundle with Hermitian metric  $g$  and compatible connection  $\nabla$  on an oriented Riemannian manifold  $M$ . Let  $v \in \Gamma(TM)$  such that the Lie derivative along  $v$  of the Riemannian volume form  $\text{Vol}(M)$  on  $M$  is zero. Then, for all  $s \in \Gamma(E)$  with compact support  $\nabla^*(v^\flat \otimes s) = -\nabla_v(s)$ .*

*Proof.* Let  $s, t \in \Gamma_c(E)$  be two sections on  $E$  with compact support. We use  $\langle \cdot, \cdot \rangle = \int_M g(\cdot, \cdot) \text{Vol}(M)$  as the inner product on  $\Gamma_c(E)$ . Using Cartan's magic formula,  $\mathcal{L}_v = d\iota_v + \iota_v d$ , and Stokes theorem we have

$$\int_M \mathcal{L}_v(g(s, t) \wedge \text{Vol}(M)) = \int_M (d\iota_v + \iota_v d)(g(s, t) \wedge \text{Vol}(M)) = 0 \quad (1.28)$$

We expand  $\mathcal{L}_v(g(s, t) \wedge \text{Vol}(M))$ . It follows from  $\mathcal{L}_v \text{Vol}(M) = 0$  that

$$\mathcal{L}_v(g(s, t) \wedge \text{Vol}(M)) = \mathcal{L}_v g(s, t) \wedge \text{Vol}(M) \quad (1.29)$$

Because  $\nabla$  is a compatible connection, we can write  $\mathcal{L}_v(s, t)$  in terms of  $\nabla_v$ .

$$\mathcal{L}_v(g(s, t) \wedge \text{Vol}(M)) = g(\nabla_v s, t) \wedge \text{Vol}(M) + g(s, \nabla_v t) \wedge \text{Vol}(M) \quad (1.30)$$

This concludes

$$\langle \nabla_v s, t \rangle = -\langle s, \nabla_v t \rangle \quad (1.31)$$

Finally, denote  $(\cdot, \cdot)$  as the inner product on  $\Gamma_c(T^*M \otimes E)$  and let  $\{e_\mu\}$  be an orthonormal frame on  $TM$ . Using the identity  $\sum_\mu e_\mu e_\mu^\flat = \text{Id}$ , we calculate

$$\langle s, \nabla^*(v^\flat \otimes t) \rangle = \langle \nabla s, v^\flat \otimes t \rangle \quad (1.32)$$

$$= \sum_{\mu, \nu} v(e_\nu) \cdot (e_\mu^\flat \otimes \nabla e_\mu s, e_\nu^\flat \otimes t) \quad (1.33)$$

$$= \sum_\mu \langle \nabla_v s, t \rangle \quad (1.34)$$

Combining Equation 1.31 and 1.34 we conclude

$$\langle s, \nabla^*(v^\flat \otimes t) \rangle = -\langle \nabla_v s, t \rangle \quad (1.35)$$

□

Recall that for a vector bundle  $E \rightarrow M$  with a metric and compatible connection the **Laplacian** is defined as

$$\nabla^* \nabla: \Gamma(E) \rightarrow \Gamma(E). \quad (1.36)$$

Using Lemma 1.17 we can calculate the Laplacian in local coordinates.

**Corollary 1.18.** *Let  $E \rightarrow M$  be a vector bundle with Hermitian metric  $g$  and compatible connection  $\nabla$  on an oriented Riemannian manifold  $M$ . Let  $x \in M$  and let  $\{\frac{\partial}{\partial x^\mu}\}$  be the Riemannian normal coordinate frame centered at  $y$ . Then at the origin of the chart the Laplacian  $\nabla^* \nabla$  satisfies*

$$\nabla^* \nabla = - \sum_{\mu} \nabla_{\mu} \nabla_{\mu} \quad (1.37)$$

where  $\nabla_{\mu} = \nabla_{\frac{\partial}{\partial x^\mu}}$ .

*Proof.* Let  $n$  be the dimension of the manifold  $M$ . By Cartan's magic formula we can easily check that  $\mathcal{L}_{\frac{\partial}{\partial x^\mu}} \text{Vol}(M) = 0$ . Indeed, it equals

$$\begin{aligned} \mathcal{L}_{\frac{\partial}{\partial x^\mu}} \text{Vol}(M) &= d \iota_{\frac{\partial}{\partial x^\mu}} dx^1 \wedge \dots \wedge dx^n \\ &= (-1)^\mu d(dx^1 \wedge \dots \wedge dx^{\mu-1} \wedge dx^{\mu+1} \wedge \dots \wedge dx^n) \\ &= 0 \end{aligned}$$

Thus we can use Lemma 1.17 and by the identity  $\sum_{\mu} \frac{\partial}{\partial x^\mu} dx^\mu = \text{Id}$  we have for all  $s \in \Gamma_c(E)$

$$\nabla^* \nabla s = \sum_{\mu} \nabla^* (dx^\mu \otimes \nabla_{\mu} s) = - \sum_{\mu} \nabla_{\mu} \nabla_{\mu} s \quad (1.38)$$

□

**Lemma 1.19.** *Let  $S \rightarrow (M, g)$  be a Clifford bundle and let  $\gamma$  be the Clifford action. In the local orthonormal basis  $\{e^\mu\}$  of  $TM$  the formal adjoint of  $\gamma: \Gamma(TM \otimes S) \rightarrow S$  is given by*

$$\gamma^*(\cdot) = -e^\mu \otimes \gamma_{\mu}(\cdot) \quad (1.39)$$

*Proof.* Let  $s, t \in \Gamma(S)$  and let  $v \in \Gamma(TM)$ . Denote  $\langle \cdot, \cdot \rangle$  as the metric on  $\Gamma(TM \otimes S)$  and  $(\cdot, \cdot)$  as the metric on  $S$ . We write down the definition of the formal adjoint and use the skew symmetry of the Clifford action.

$$\langle v \otimes s, \gamma^* t \rangle = g(\gamma(v)s, t) = -g(s, \gamma(v)t) \quad (1.40)$$

We compare this to  $\langle v \otimes s, -e_\mu \otimes (\gamma^\mu t) \rangle$ . Expand  $v$  as  $v = v_\nu e^\nu$ . Then,

$$\langle v \otimes s, -e^\mu \otimes (\gamma_\mu t) \rangle = \langle v_\nu e^\nu \otimes s, -e^\mu \otimes (\gamma_\mu t) \rangle = v_\nu \cdot g(e^\nu, e^\mu) \cdot (s, \gamma_\nu t) \quad (1.41)$$

By orthonormality  $v_\nu \cdot g(e^\nu, e^\mu)$  simplifies to  $v_\mu$  and hence

$$\langle v \otimes s, -e^\mu \otimes (\gamma_\mu t) \rangle = v_\mu \cdot \langle s, \gamma_\mu t \rangle = -(s, \gamma(v) t) \quad (1.42)$$

Comparing Equation 1.40 and 1.42 we conclude the result.  $\square$

**Proposition 1.20** (Roe [1998], Proposition 3.11). *Let  $S \rightarrow M$  be a Clifford bundle and let  $D$  be a Dirac operator. Then  $D$  is formally self-adjoint.*

*Proof.* Recall that  $D$  is defined as the composition of the following maps

$$\Gamma(S) \xrightarrow{\nabla} \Gamma(T^*M \otimes S) \xrightarrow{g^{-1}} \Gamma(TM \otimes S) \xrightarrow{\gamma} \Gamma(S) \quad (1.43)$$

So the formal adjoint of  $D$  is given by

$$\Gamma(S) \xrightarrow{\gamma^*} \Gamma(TM \otimes S) \xrightarrow{g} \Gamma(T^*M \otimes S) \xrightarrow{\nabla^*} \Gamma(S) \quad (1.44)$$

Let  $x \in M$  and consider the Riemannian normal coordinates centered at  $y$ . At  $x \in M$  we get for any  $s \in \Gamma(S)$  that

$$s \mapsto -\frac{\partial}{\partial x^\mu} \otimes \gamma^\mu s \mapsto -\sum_\mu dx^\mu \otimes \gamma_\mu s \mapsto \nabla_\mu \gamma_\mu s \quad (1.45)$$

At the origin of our chart, we have  $\nabla_\mu e^\mu = 0$ . So by the compatibility condition we get

$$D^*s = \nabla_\mu \gamma^\mu s = \gamma^\mu \nabla_\mu s = Ds \quad (1.46)$$

Therefore,  $D$  is formally self-adjoint.  $\square$

## 1.4 The Weitzenböck formula

In this paragraph we compare the difference between the square of an Dirac operator and the Laplacian. The result will be the so called Weitzenböck formula, which gives the difference in terms of the Clifford action and the curvature. Before we do this, we need to generalize the Feynman-slash operator for 2-forms.

**Definition 1.21.** Let  $S \rightarrow (M, g)$  be a Clifford bundle with the Clifford action  $\gamma$  and let  $K \in \Omega^2(M, \text{End}(S))$  be a 2-form with values in  $\text{End}(S)$ . Define the map  $\Omega^2(M) \rightarrow \Gamma(T^*M \otimes T^*M)$  as  $\alpha \wedge \beta \mapsto \frac{1}{2}(\alpha \otimes \beta - \beta \otimes \alpha) \in \Gamma(T^*M \otimes T^*M)$ . Interpret  $g$  as the isomorphism  $g: \Gamma(TM) \rightarrow \Gamma(T^*M)$  which is given by  $v \mapsto g(v, \cdot)$ . The **Clifford contraction of  $K$**  is the composition of the maps

$$\Gamma(S) \xrightarrow{K} \Omega^2(M, S) \hookrightarrow \Gamma(T^*M \otimes T^*M \otimes S) \xrightarrow{g^{-1}} \Gamma(TM \otimes TM \otimes S) \xrightarrow{\gamma} \Gamma(S). \quad (1.47)$$

The Clifford contraction of  $K$  is denoted as  $K$ .

In a local orthonormal frame  $\{e^\mu\}$  on  $TM$ ,  $K$  can be written as  $K = \sum_{\mu < \nu} e^{\mu\nu} \wedge e^{\nu\mu} \otimes K(e_\mu, e_\nu)$ . Therefore, the Clifford contraction of  $K$  equals It can be shown that the Clifford contraction of  $K$  equals

$$K = \sum_{\mu < \nu} \gamma_\mu \gamma_\nu K(e^\mu, e^\nu) \quad (1.48)$$

We are now able to show the Weitzenbock formula.

**Theorem 1.22** (Weitzenbock). Let  $S \rightarrow M$  be a Clifford bundle with compatible connection  $\nabla$ . Let  $D$  be the Dirac operator and  $K$  be the curvature w.r.t.  $\nabla$ . Denote  $K$  as the Clifford contraction of the curvature. Then

$$D^2 = \nabla^* \nabla + K \quad (1.49)$$

*Proof.* Let  $x \in M$  and let  $\{\frac{\partial}{\partial x^\mu}\}$  be the Riemannian normal coordinate frame centered at  $y$ . Let  $\{\gamma^\mu, \gamma^\nu\}$  and  $[\gamma^\mu, \gamma^\nu]$  be the (anti)-commutator. That is,

$$\{\gamma^\mu, \gamma^\nu\} = \gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu \quad (1.50)$$

$$[\gamma^\mu, \gamma^\nu] = \gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu \quad \forall \mu, \nu \quad (1.51)$$

Using Equation 1.12 we calculate the square of the Dirac operator.

$$\begin{aligned} D^2 &= \sum_{\mu\nu} \gamma^\mu \nabla_\mu \gamma^\nu \nabla_\nu = \sum_{\mu\nu} \gamma^\mu \gamma (\nabla_\mu e^\nu) \nabla_\nu + \gamma^\mu \gamma^\nu \nabla_\mu \nabla_\nu \\ &= \sum_{\mu\nu} \gamma^\mu \gamma^\nu \nabla_\mu \nabla_\nu \end{aligned} \quad (1.52)$$

This can be written in terms of the commutator and the anti-commutator.

$$D^2 = \frac{1}{2} \sum_{\mu\nu} \{\gamma_\mu, \gamma_\nu\} \nabla_\mu \nabla_\nu + [\gamma_\mu, \gamma_\nu] \nabla_\mu \nabla_\nu \quad (1.53)$$

$$= \sum_{\mu\nu} -g \left( \frac{\partial}{\partial x^\mu}, \frac{\partial}{\partial x^\nu} \right) \nabla_\mu \nabla_\nu + \frac{1}{2} \gamma_\mu \gamma_\nu [\nabla_\mu, \nabla_\nu] \quad (1.54)$$

$$= - \sum_{\mu} \nabla_\mu \nabla_\mu + \sum_{\mu < \nu} \gamma^\mu \gamma^\nu K \left( \frac{\partial}{\partial x^\mu}, \frac{\partial}{\partial x^\nu} \right) \quad (1.55)$$

From Corollary 1.18 and Equation 1.48 we conclude that  $D^2 = \nabla^* \nabla + K$ .  $\square$

Next, we will prove a refined version of the Weitzenböck formula, namely the Lichnerowicz formula. We show this by considering the commutation relations between the the curvature  $K$  and the Clifford action. Then  $K$  will split into the scalar curvature and a the Clifford contraction of a 2-form called the Riemann endomorphism.

**Definition 1.23.** Let  $S \rightarrow (M, g)$  be a Clifford bundle, let  $\gamma$  the Clifford action and let  $R$  be the Riemann curvature tensor. The **Riemann endomorphism**  $R^S$  of  $S$  is the following composition of maps applied to the Riemann curvature tensor  $R$ :

$$\begin{array}{ccc}
 \Gamma(\Omega^2(M) \otimes TM \otimes T^*M) & \xrightarrow{g^{-1}} & \Gamma(\Omega^2(M) \otimes TM \otimes TM) \\
 \downarrow \gamma & & \downarrow \gamma \\
 \Gamma(\Omega^2(M) \otimes \text{End}(S) \otimes \text{End}(S)) & \xrightarrow{\circ} & \Gamma(\Omega^2(M) \otimes \text{End}(S)) \\
 \downarrow 1/4 & & \downarrow 1/4 \\
 \Gamma(\Omega^2(M) \otimes \text{End}(S)) & & \Gamma(\Omega^2(M) \otimes \text{End}(S))
 \end{array}$$

$R \mapsto R^S$  (indicated by a dashed arrow from the first to the second row)

Here  $\circ$  denotes the point-wise composition of endomorphisms and  $\frac{1}{4}$  denotes deviation by four.

Given a local orthonormal frame  $\{e^\mu\}$  we calculate the Riemann endomorphism by

following the diagram:

$$\begin{array}{ccc}
 R = \sum_{\mu\nu} \langle Re^\mu, e^\nu \rangle \otimes e^\mu \otimes e_\nu^b & \xrightarrow{g^{-1}} & \sum_{\mu\nu} \langle Re^\mu, e^\nu \rangle \otimes e^\mu \otimes e^\nu \\
 & \searrow \gamma & \\
 \sum_{\mu\nu} \langle Re^\mu, e^\nu \rangle \otimes \gamma_\mu \otimes \gamma_\nu & \xrightarrow{\circ} & \sum_{\mu\nu} \langle Re^\mu, e^\nu \rangle \gamma_\mu \gamma_\nu \\
 & \nwarrow 1/4 & \\
 \frac{1}{4} \sum_{\mu\nu} \langle Re^\mu, e^\nu \rangle \gamma_\mu \gamma_\nu & & 
 \end{array} \tag{1.56}$$

Hence, in a local orthonormal frame  $\{e^\mu\}$  on  $TM$ , the Riemann endomorphism is given by

$$R^S(X, Y) = \frac{1}{4} \gamma_\mu \gamma_\nu \langle R(X, Y) e^\mu, e^\nu \rangle \quad \forall X, Y \in \Gamma(TM) \tag{1.57}$$

**Lemma 1.24** (Roe [1998], Lemma 3.13 and 3.15). *Let  $S \rightarrow M$  be a Clifford bundle, let  $K$  be the curvature on  $S$  and  $R$  be the Riemann curvature on  $M$ . Use  $\gamma$  for the Clifford action and let  $R^S$  be the Riemann endomorphism. Then for all  $u, v, w \in \Gamma(TM)$*

$$[K(u, v), \gamma(w)] = [R^S(u, v), \gamma(w)] \tag{1.58}$$

where  $[\cdot, \cdot]$  is the commutator.

*Proof.* By the definition of the curvature, the left hand side of Equation 1.58 equals

$$[K(u, v), \gamma(w)] = [\nabla_u \nabla_v - \nabla_v \nabla_u, \gamma(w)] \tag{1.59}$$

According to the Jacobi identity this is

$$[K(u, v), \gamma(w)] = [\nabla_u \nabla_v - \nabla_v \nabla_u - \nabla_{[u, v]}, \gamma(w)] \tag{1.60}$$

$$\begin{aligned}
 &= \nabla_u [\nabla_v, \gamma(w)] + [\nabla_u, \gamma(w)] \nabla_v - \\
 &\quad - \nabla_v [\nabla_u, \gamma(w)] - [\nabla_v, \gamma(w)] \nabla_u - [\nabla_{[u, v]}, \gamma(w)]
 \end{aligned} \tag{1.61}$$

$$= [\nabla_u, [\nabla_v, \gamma(w)]] - [\nabla_v, [\nabla_u, \gamma(w)]] - [\nabla_{[u, v]}, \gamma(w)] \tag{1.62}$$

From Definition 1.6 follows that the commutator  $[\nabla_v, \gamma(w)] = \gamma(\nabla_u w)$  and hence

$$[K(u, v), \gamma(w)] = [\nabla_u, \gamma(\nabla_v w)] - [\nabla_v, \gamma(\nabla_u w)] - \gamma(\nabla_{[u, v]} w) \tag{1.63}$$

$$= \gamma(\nabla_u \nabla_v w - \nabla_v \nabla_u w - \nabla_{[u, v]} w) \tag{1.64}$$

$$= \gamma(R(u, v)w) \tag{1.65}$$

This gives an explicit result for the left hand side. To calculate the right hand side of Equation 1.58 we need a local orthonormal frame  $\{e^\mu\}$  of  $TM$ . Using the identity

$$[\gamma(u)\gamma(v), \gamma(w)] = 2g(u, w)\gamma(v) - 2g(v, w)\gamma(u) \quad (1.66)$$

we get that the right hand side equals

$$[R^S(u, v), \gamma(w)] = \frac{1}{4} \langle R(u, v)e^\mu, e^\nu \rangle \cdot [\gamma_\mu \gamma_\nu, \gamma(w)] \quad (1.67)$$

$$= \frac{1}{2} \langle R(u, v)w, e^\nu \rangle \gamma_\nu - \frac{1}{2} \langle R(u, v)e^\mu, w \rangle \gamma_\mu \quad (1.68)$$

By the anti-symmetry property  $\langle R(u, v)e^\mu, w \rangle = -\langle R(u, v)w, e^\mu \rangle$  it follows that

$$[R^S(u, v), \gamma(w)] = \langle R(u, v)w, e^\nu \rangle \gamma_\nu = \gamma(R(u, v)w) \quad (1.69)$$

Comparing Equation 1.65 and 1.69 we conclude the result.  $\square$

**Definition 1.25.** Let  $S \rightarrow (M, g)$  be a Clifford bundle over a Riemannian manifold and let  $E \rightarrow M$  be a vector bundle. A **Clifford endomorphism** is a section  $F$  of  $\text{End}_{\mathbb{C}}(S) \otimes E$  such that for all  $v \in TM$ ,  $F \circ \gamma(v)$  equals  $\gamma(v) \circ F$ .

By Lemma 1.24 we conclude that  $K - R^S \in \Omega^2(\text{End}(S))$  is a Clifford endomorphism. This difference will play an important role in the index theorem and so we give it a name.

**Definition 1.26.** Let  $S \rightarrow M$  be a Clifford bundle and let  $K$  be its curvature. The **twisting curvature**  $F^S$  is the Clifford endomorphism  $K - R^S$ .

Using the twisting curvature and the Weitzenböck formula we can write

$$D^2 = \nabla^* \nabla + F^S + R^S \quad (1.70)$$

However, this can be further reduced.

**Proposition 1.27** (Lichnerowicz [1963]). Let  $S \rightarrow (M, g)$  be a Clifford bundle. Let  $D$  be the Dirac operator and  $F^S$  be the Clifford contraction of the twisting curvature. Denote  $\kappa$  as the scalar curvature. Then,

$$D^2 = \nabla^* \nabla + F^S + \frac{1}{4} \kappa \quad (1.71)$$

*Proof.* It is sufficient to show that  $R^S = \frac{1}{4}\kappa$ . In a local orthonormal frame  $\{e_\mu\}$  of  $TM$ , the Clifford contraction of the Riemann endomorphism is given by

$$R^S = \frac{1}{8}\gamma^\mu\gamma^\nu\gamma^\rho\gamma^\sigma\langle R(e_\mu, e_\nu)e_\rho, e_\sigma\rangle \quad (1.72)$$

It follows from the Bianchi identity that

$$R^S = -\frac{1}{8}\langle R(e_\mu, e_\nu)e_\rho, e_\sigma\rangle(\gamma^\rho\gamma^\mu\gamma^\nu\gamma^\sigma + \gamma^\nu\gamma^\rho\gamma^\mu\gamma^\sigma) \quad (1.73)$$

Using the commutation relation of the Clifford action, we can reorder the gammas back into  $\gamma^\mu\gamma^\nu\gamma^\rho\gamma^\sigma$ .

$$R^S = -\frac{1}{8}\langle R(e_\mu, e_\nu)e_\rho, e_\sigma\rangle(\gamma^\mu\gamma^\nu\gamma^\rho\gamma^\sigma - \gamma^\nu\gamma^\mu\gamma^\rho\gamma^\sigma - 4\delta^{\mu\rho}\gamma^\nu\gamma^\sigma + 2\delta^{\nu\rho}\gamma^\mu\gamma^\sigma) \quad (1.74)$$

By the antisymmetry of 2-forms, this simplifies into

$$R^S = -\frac{1}{8}\langle R(e_\mu, e_\nu)e_\rho, e_\sigma\rangle(2\gamma^\mu\gamma^\nu\gamma^\rho\gamma^\sigma - 6\delta^{\mu\rho}\gamma^\nu\gamma^\sigma) \quad (1.75)$$

$$= -2R^S + \frac{3}{4}\langle R(e_\mu, e_\nu)e_\mu, e_\sigma\rangle\gamma^\nu\gamma^\sigma \quad (1.76)$$

Because  $\langle R(e_\mu, e_\nu)e_\rho, e_\sigma\rangle = \langle R(e_\rho, e_\sigma)e_\mu, e_\nu\rangle$ , the Clifford contraction of the Riemann endomorphism becomes

$$R^S = \frac{1}{4}\langle R(e_\mu, e_\nu)e_\mu, e_\sigma\rangle\gamma^\nu\gamma^\sigma \quad (1.77)$$

$$= \frac{1}{8}\langle R(e_\mu, e_\nu)e_\mu, e_\sigma\rangle(\gamma^\nu\gamma^\sigma + \gamma^\sigma\gamma^\nu) \quad (1.78)$$

$$= -\frac{1}{4}\langle R(e_\mu, e_\nu)e_\mu, e_\nu\rangle \quad (1.79)$$

We conclude that  $R^S = \frac{1}{4}\kappa$ . □

## 2 Calculating anomalies using Feynman diagrams

In this chapter we introduce the notion of anomalies. For this we revisit Noethers theorem and we investigate if there are obstructions when we generalize it to quantum field theory. These obstructions are called anomalies:

**Definition 2.1.** *We call classical conservation law that is not satisfied in quantum field theory an **anomaly**.*

In this chapter we use the perturbative approach to quantum field theory. That is, we calculate the amplitude of Feynman diagrams that relate to classical conservation currents. We work out examples where the classical currents are not conserved.

Most calculations are done by van Nieuwenhuizen [1989], but without much detail.

### 2.1 Revisit of Noethers theorem

Informally, Noethers theorem can be stated as follows:

If a classical system has a continuous symmetry, then there are corresponding quantities whose values are conserved.

To elaborate this, we work out three examples. In each example we define a physical system by an action and we assume that the system is invariant under a symmetry. We then calculate what happens to the action when we apply this symmetry and we find that there must exist conserved quantities.

The first example is a massless complex scalar field  $\phi \in \Gamma(\mathbb{R}^4 \otimes \mathbb{C})$ . We describe the physics by the action

$$S = \int_{\mathbb{R}^4} dx \partial_\mu \phi^*(x) \partial^\mu \phi(x) \quad (2.1)$$

Assume that this system is invariant under the variation

$$\phi \mapsto \tilde{\phi} = e^{i\alpha} \phi \quad (2.2)$$

$$\phi^* \mapsto \tilde{\phi}^* = e^{-i\alpha} \phi^* \quad (2.3)$$

where  $\alpha$  is a smooth but infinitely small real valued function. This means that the action must not change under this transformation. We calculate  $\tilde{S}$  explicitly:

$$\tilde{S} := \int_{\mathbb{R}^4} dx \partial_\mu \tilde{\phi}^* \partial^\mu \tilde{\phi} \quad (2.4)$$

$$= \int_{\mathbb{R}^4} dx \partial_\mu (e^{-i\alpha(x)} \phi^*) \partial^\mu (e^{i\alpha(x)} \phi) \quad (2.5)$$

$$= S - i \int_{\mathbb{R}^4} dx \partial_\mu (\alpha(x)) \phi^* \partial^\mu \phi + i \partial^\mu (\alpha(x)) \partial_\mu (\phi^*) \phi + \mathcal{O}(\alpha^2) \quad (2.6)$$

$$= S + i \int_{\mathbb{R}^4} dx \alpha(x) \cdot (\partial_\mu (\phi^* \cdot \partial^\mu \phi - \partial^\mu \phi^* \cdot \phi)) + \mathcal{O}(\alpha^2) \quad (2.7)$$

Invariance implies that  $\int_{\mathbb{R}^4} dx \alpha(x) \cdot (\partial_\mu (\phi^* \cdot \partial^\mu \phi - \partial^\mu \phi^* \cdot \phi)) = 0$ . This is only possible when

$$\partial_\mu (\phi^* \cdot \partial^\mu \phi - \partial^\mu \phi^* \cdot \phi) = 0 \quad (2.8)$$

This equation implies that  $j^\mu := \phi^* \cdot \partial^\mu \phi - \partial^\mu \phi^* \cdot \phi$  is conserved. This result is predicted by Noethers theorem. It states that for all symmetries, there exists a **current**  $j^\mu$  such that  $\partial_\mu j^\mu = 0$ .

In this thesis we mainly look at the chiral symmetry. We study the fermions  $\psi \in \Gamma(\mathbb{R}^4 \otimes \mathbb{C}^4)$  with mass  $m$  which are coupled to external vector field  $V^\mu \in \Gamma(T\mathbb{R}^4)$  and axial-vector field  $A^\mu \in \Gamma(T\mathbb{R}^4)$ . The external fields are not necessary Abelian. So let  $\mathfrak{g}$  be a lie algebra and let  $\{\lambda_a\}$  be a set of generators of  $\mathfrak{g}$ . We write  $V^\mu = V_a^\mu \lambda^a \in \Gamma(T\mathbb{R}^4 \otimes \mathfrak{g})$  and  $A^\mu = A_a^\mu \lambda^a \in \Gamma(T\mathbb{R}^4 \otimes \mathfrak{g})$ . The physics is described by the action

$$S = \int_{\mathbb{R}^4} -\bar{\psi}(\not{\partial} + m)\psi + i\bar{\psi}(V + A\gamma_5)\psi dx \quad (2.9)$$

The **Abelian chiral symmetry** is given by

$$\psi \mapsto e^{i\alpha(x)\gamma_5} \psi \quad (2.10)$$

$$\bar{\psi} \mapsto \bar{\psi} e^{i\alpha(x)\gamma_5} \quad (2.11)$$

where  $\alpha$  is an infinitely small real valued function. Under this symmetry the action transforms into

$$S \mapsto \int_{\mathbb{R}^4} dx -\bar{\psi} e^{i\alpha(x)\gamma_5} (\not{\partial} + m) e^{i\alpha(x)\gamma_5} \psi + i\bar{\psi} e^{i\alpha(x)\gamma_5} (V + A\gamma_5) e^{i\alpha(x)\gamma_5} \psi \quad (2.12)$$

$$S \mapsto S + \int_{\mathbb{R}^4} dx -\bar{\psi} \gamma^\mu \gamma_5 \psi \cdot \partial_\mu \alpha(x) - m\bar{\psi} (e^{2i\alpha(x)\gamma_5} - 1) \psi \quad (2.13)$$

$$S \mapsto S + \int_{\mathbb{R}^4} dx \alpha(x) (\partial_\mu (\bar{\psi} \gamma^\mu \gamma_5 \psi) - 2im \bar{\psi} \gamma_5 \psi) + \mathcal{O}(\alpha^2) \quad (2.14)$$

The invariance of the action implies that

$$\partial_\mu (\bar{\psi} \gamma^\mu \gamma_5 \psi) = 2im \bar{\psi} \gamma_5 \psi \quad (2.15)$$

If  $m = 0$ , we notice that  $j^\mu = \bar{\psi} \gamma^\mu \gamma_5 \psi$  is conserved. This is Noethers theorem for the Abelian chiral symmetry.

There is also the **non-Abelian chiral symmetry**. Again we consider the action given in Equation 2.9 but now we assume that  $S$  is invariant under

$$\psi \mapsto e^{i\alpha(x)\lambda_a \gamma_5} \psi \quad (2.16)$$

$$\bar{\psi} \mapsto \bar{\psi} e^{i\alpha(x)\lambda_a \gamma_5}. \quad (2.17)$$

In the same manner we can calculate the variation of  $S$ . Note that  $\lambda_a$  does not commute with  $A$  and  $V$ . Therefore, the variation of  $S$  is not equal to Equation 2.14, but is actually equal to

$$S + \int_{\mathbb{R}^4} dx \alpha(x) (\partial_\mu (\bar{\psi} \gamma^\mu \gamma_5 \lambda_a \psi) + \bar{\psi} \gamma^\mu \gamma_5 (iV_\mu^b + iA_\mu^b \gamma_5) [\lambda_a, \lambda_b] \psi - 2im \bar{\psi} \gamma_5 \lambda_a \psi) \quad (2.18)$$

By the invariance of the action we conclude

$$\partial_\mu (\bar{\psi} \gamma^\mu \gamma_5 \lambda_a \psi) = -\bar{\psi} \gamma^\mu \gamma_5 (iV_\mu^b + iA_\mu^b \gamma_5) [\lambda_a, \lambda_b] \psi + 2im \bar{\psi} \gamma_5 \lambda_a \psi \quad (2.19)$$

To simplify the notation, let  $S \in \{\text{Id}\} \cup \mathfrak{g}$ . We define the **chiral current** as

$$j_S^\mu = \bar{\psi} \gamma^\mu \gamma_5 S \psi \quad (2.20)$$

If  $S = \text{Id}$ , then  $j_S^\mu$  is the current for the Abelian chiral symmetry. If  $S = \lambda_a$ , then  $j_a^\mu$  is the current for the non-Abelian chiral symmetry. In both cases, the chiral current satisfies

$$\partial_\mu (\bar{\psi} \gamma^\mu \gamma_5 S \psi) = -\bar{\psi} \gamma^\mu \gamma_5 (iV_\mu^b + iA_\mu^b \gamma_5) [S, \lambda_b] \psi + 2im \bar{\psi} \gamma_5 S \psi \quad (2.21)$$

## 2.2 Chiral currents in triangle diagrams

In the previous section, we applied Noethers theorem to chiral currents. The result we found was based on classical mechanics. Now we calculate the same chiral currents in quantum field theory. In this chapter we use one-loop approximations to calculate the amplitudes of scattering processes that are linearly dependent on the chiral current.




Figure 1: *Example of an anomalous scattering process. On top a particle of an axial-vector field couples to a massive fermion/anti-fermion which then decays into two vector fields. The labels  $p, q$  and  $l$  are used to denote the momentum,  $\mu, \nu, \rho$  are used to denote the momentum index and  $a$  and  $b$  are used to denote the gauge index. In case of the Abelian anomaly we set  $S = \text{Id}$ , else  $S$  is an element of the gauge group.*

We compare these amplitudes with the classically expected results and we will notice that they do not coincide.

Consider the  $AVV$  diagram depicted in Figure 1. Using the following Feynman rules, we can calculate the scattering amplitude  $M_{abS}^{\mu\nu\rho}$ :

1. For each loop, add an integral  $\int \frac{d^4 l}{(2\pi)^4}$ .
2. For each vertex between  $\psi, \bar{\psi}$  and  $V_a^\mu$ , add the term  $-\gamma^\mu \lambda_a$ .
3. For each vertex between  $\psi, \bar{\psi}$  and  $A_a^\mu$ , add the term  $-\gamma^\mu \gamma_5 \lambda_a$ .
4. For each internal  $\psi$ -propagator with momentum  $k$  add the term  $\frac{-i\not{k}+m}{k^2+m^2}$ .
5. Take the trace over the gauge group and the Clifford algebra.
6. The above terms appear in the same order in the trace as they appear in the loop. However, if the current flows clockwise, then we add the terms counterclockwise and vice versa.

According to these rules, the scattering amplitude becomes

$$M_{abS}^{\mu\nu\rho} = \int \frac{d^4 l}{(2\pi)^4} \text{tr} \left[ \gamma_5 \gamma^\rho S \frac{-i(\not{l} - \not{q}) + m}{(l - q)^2 + m^2} \gamma_\mu \lambda_a \frac{-i\not{l} + m}{l^2 + m^2} \gamma_\nu \lambda_b \frac{-i(\not{l} + \not{p}) + m}{(l + p)^2 + m^2} \right] \quad (2.22)$$

Comparing Equation 2.22 with Equation 2.20, we see that the term  $\gamma^\rho \gamma_5 S$  is related to the chiral current. To mimic Equation 2.21, we want to calculate  $\partial_\rho M_{abs}^{\mu\nu\rho}$ . Recall however that Feynman diagrams apply a Fourier transformation on the fields. Hence, we consider contraction  $(p + q)_\rho M_{abs}^{\mu\nu\rho}$ .

By power counting we see that  $M_{abS}^{\mu\nu\rho}$  is proportional to  $\int \frac{d^4 l}{l^3}$  and so is linearly divergent. To overcome this problem, we have to regularize the amplitude. That is, we modify the theory such that the amplitude is finite. The physical situation is when the alteration is negligible small. We work out two different regularization methods.

### 2.2.1 Dimensional regularization

One method is called dimensional regularization. The idea is to generalize equation 2.22 to  $n$  dimensions. Also the Clifford algebra is generalized. However, the grading operator  $\gamma_5$  is still

$$\gamma_5 = -i\gamma^0\gamma^1\gamma^2\gamma^3$$

The result is that  $\gamma_5$  anti-commutes with  $\gamma^0, \gamma^1, \gamma^2$  and  $\gamma^3$ , but commutes with the other  $\gamma_\mu$ . Furthermore, we still assume that the external fields are four dimensional. This method was introduced by 't Hooft and Veltman [1972] and is also called 't Hooft-Veltman regularization.

For dimensional regularization we need to decompose  $l$  into  $k + k_\perp$ , with  $k \in \mathbb{R}^4$  and  $k_\perp \in \mathbb{R}^{n-4}$ . The regularized  $(p + q)_\rho M_{abS}^{\mu\nu\rho}$  becomes

$$(p + q)_\rho M_{abS}^{\mu\nu\rho} = \int \frac{d^n l}{(2\pi)^n} \text{tr} \left[ \gamma_5 (\not{p} + \not{q}) \frac{-i(\not{k} + \not{k}_\perp - \not{q}) + m}{(k + k_\perp - q)^2 + m^2} \gamma_\mu \frac{-i(\not{k} + \not{k}_\perp) + m}{(k + k_\perp)^2 + m^2} \gamma_\nu \times \right. \\ \left. \frac{-i(\not{k} + \not{k}_\perp + \not{p}) + m}{(k + k_\perp + p)^2 + m^2} \right] \text{tr} [S \lambda_a \lambda_b]. \quad (2.23)$$

A little calculation shows that we can write

$$\gamma_5 (\not{q} + \not{p}) = \gamma_5 (\not{q} + \not{p}) - \gamma_5 \not{l} - \not{l} \gamma_5 + 2\gamma_5 \not{k}_\perp \quad (2.24)$$

$$= -\gamma_5 (\not{l} - \not{q} - im) - (\not{l} + \not{p} - im) \gamma_5 - 2im \gamma_5 + 2\gamma_5 \not{k}_\perp. \quad (2.25)$$

The terms  $-\gamma_5(\not{\ell}-\not{q}-im)-(\not{\ell}+\not{p}-im)\gamma_5$  in equation 2.25 are related to  $\bar{\psi}\gamma^\mu\gamma_5(iV_\mu^b+iA_\mu^b\gamma_5)[S,\lambda_b]\psi$  from equation 2.21. Indeed, the term  $-\gamma_5(\not{\ell}-\not{q}-im)$  simplifies the trace in equation 2.23 into

$$\text{tr} \left[ \gamma_5 \gamma_\mu \frac{-i(\not{k} + \not{k}_\perp) + m}{(k + k_\perp)^2 + m^2} \gamma_\nu \frac{-i(\not{k} + \not{k}_\perp + \not{p}) + m}{(k + k_\perp + p)^2 + m^2} \right] \text{tr} [S\lambda_a\lambda_b]. \quad (2.26)$$

This can be interpreted as a one-loop diagram with two external fields where one vertex is connected to an external (axial-) vector field with gauge index  $S\lambda_a$ . This is up to the factor  $-i$  in one to one correspondence with

$$\bar{\psi}\gamma_5\gamma^\mu(iV_\mu^a+iA_\mu^a\gamma_5)S\lambda_a\psi. \quad (2.27)$$

Even more, this part of the scattering amplitude vanish in dimensional regularization. The numerator in equation 2.26 is equal to  $-4i\epsilon^{\mu\nu\sigma\tau}k_\sigma p_\tau \cdot \text{tr} [S\lambda_a\lambda_b]$ . After a shift  $k \rightarrow k - p/2$ , the amplitude of this terms becomes

$$-4i\epsilon^{\mu\nu\sigma\tau}p_\tau \int \frac{d^n l}{(2\pi)^n} \frac{k_\sigma}{((l - p/2)^2 + m^2)((l + p/2)^2 + m^2)}. \quad (2.28)$$

This term vanishes, because it is anti-symmetric in  $k$ .

The only part in equation 2.25 that is not related to the classical conservation law, is the term  $2\gamma_5\not{k}_\perp$ . In the rest of this section we show that this term does not vanish by explicitly calculating

$$M_{abS,an}^{\mu\nu} := 2 \int \frac{d^n l}{(2\pi)^n} \text{tr} \left[ \gamma_5 \not{k}_\perp \frac{-i(\not{k} + \not{k}_\perp - \not{q}) + m}{(k + k_\perp - q)^2 + m^2} \gamma_\mu \frac{-i(\not{k} + \not{k}_\perp) + m}{(k + k_\perp)^2 + m^2} \gamma_\nu \times \right. \\ \left. \frac{-i(\not{k} + \not{k}_\perp + \not{p}) + m}{(k + k_\perp + p)^2 + m^2} \right] \text{tr} [S\lambda_a\lambda_b]. \quad (2.29)$$

The denominator is symmetric in  $k_\perp$ . Therefore all odd orders of  $k_\perp$  in the numerator vanish by antisymmetry. Also the quartic order of  $k_\perp$  disappear, because these terms are proportional to

$$\text{tr}(\gamma_5 k_\perp^4) = \|k_\perp\|^4 \cdot \text{tr}(\gamma_5) = 0. \quad (2.30)$$

The 2<sup>nd</sup> order terms of  $k_\perp^2$  are

$$\begin{aligned} & \text{Tr} [\gamma_5 \not{k}_\perp \not{k}_\perp \gamma_\mu (-i\not{k} + m) \gamma_\nu (-i(\not{k} + \not{p}) + m)] \\ & + \text{Tr} [\gamma_5 \not{k}_\perp (-i(\not{k} - \not{q}) + m) \gamma_\mu \not{k}_\perp \gamma_\nu (-i(\not{k} + \not{p}) + m)] \\ & + \text{Tr} [\gamma_5 \not{k}_\perp (-i(\not{k} - \not{q}) + m) \gamma_\mu (-i\not{k} + m) \gamma_\nu \not{k}_\perp]. \end{aligned} \quad (2.31)$$

This trace can be calculated using the commutation rules and the trace identities<sup>2</sup>. It is equal to

$$4i\|k_{\perp}\|^2\epsilon^{\mu\nu\rho\sigma}q_{\rho}p_{\sigma} \quad (2.32)$$

where  $\epsilon^{\mu\nu\rho\sigma}$  is the Levi-Civita symbol<sup>3</sup>. Combining equation 2.32 and 2.29 we get that  $M_{abS,an}^{\mu\nu}$  equals

$$8\epsilon^{\mu\nu\rho\sigma}q_{\rho}p_{\sigma} \cdot \int \frac{d^n l}{(2\pi)^n} \frac{k_{\perp}^2}{((k-q)^2 + k_{\perp}^2 + m^2)(k^2 + k_{\perp}^2 + m^2)((k+p)^2 + k_{\perp}^2 + m^2)} \quad (2.33)$$

This integral cannot vanish, because the integrand is positive. It can happen that it is linearly dependent on the dimensions of  $k_{\perp}$ . Then it vanishes when  $n \rightarrow 4$ . This is not the case and we will spend the rest of this paragraph to show this.

First we simplify equation 2.33 by using the Feynman trick

$$\frac{1}{ABC} = 2 \int_0^1 dx \int_0^{1-x} dy \frac{1}{(xA + yB + (1-x-y)C)^3} \quad (2.34)$$

where we choose the following values for  $A$ ,  $B$  and  $C$ :

$$A = (k+p)^2 + k_{\perp}^2 + m^2 \quad B = (k-q)^2 + k_{\perp}^2 + m^2 \quad C = k^2 + k_{\perp}^2 + m^2 \quad (2.35)$$

The denominator in equation 2.34 becomes

$$xA + yB + (1-x-y)C = (k + xp - yq)^2 + k_{\perp}^2 + m^2 + xp^2 + yq^2 - (xp - yq)^2. \quad (2.36)$$

If we shift  $k$  to  $k - xp + yq$  and define

$$a = m^2 + xp^2 + yq^2 - (xp - yq)^2, \quad (2.37)$$

then the denominator equals  $(l^2 + a^2)^3$ . The anomalous term  $M_{abS,an}^{\mu\nu}$  simplifies into

$$16\epsilon^{\mu\nu\rho\sigma}q_{\rho}p_{\sigma} \cdot \int_0^1 dx \int_0^{1-x} dy \int \frac{d^n l}{(2\pi)^n} \frac{k_{\perp}^2}{(l^2 + a^2)^3} \text{tr}[S\lambda_a\lambda_b] \quad (2.38)$$

<sup>2</sup>These are  $\text{tr}(\gamma_5) = 0$ ,  $\text{tr}(\gamma_5\gamma_{\mu_1}\dots\gamma_{\mu_{2k+1}}) = 0$ ,  $\gamma(\gamma_5\gamma_{\mu}\gamma_n u) = 0$  and  $\text{tr}(\gamma_5\gamma_{\mu}\gamma_{\nu}\gamma_{\rho}\gamma_{\sigma}) = 4i\epsilon^{\mu\nu\rho\sigma}$ .

<sup>3</sup> $\epsilon^{\mu\nu\rho\sigma} = \pm 1$  depending on the permutation of the indices. If two indices are equal, the Levi-Civita symbol is vanishes.

From symmetry arguments we can replace  $\frac{k_{\perp}^2}{(l^2+a^2)^3}$  with  $\frac{n-4}{n} \cdot \frac{l^2}{(l^2+a^2)^3}$ . Indeed, notice that

$$\frac{d^n l}{(2\pi)^n} \frac{k_{\perp}^2}{(l^2+a^2)^3} = (n-4) \frac{d^n l}{(2\pi)^n} \frac{k_n^2}{(l^2+a^2)^3} \quad (2.39)$$

$$= \frac{n-4}{n} \frac{d^n l}{(2\pi)^n} \frac{k_1^2 + \dots + k_n^2}{(l^2+a^2)^3} \quad (2.40)$$

$$= \frac{n-4}{n} \frac{d^n l}{(2\pi)^n} \frac{l^2}{(l^2+a^2)^3} \quad (2.41)$$

Hence, the integrand from equation 2.38 equals

$$\frac{n-4}{n} \frac{l^2}{(l^2+a^2)^3} \text{tr}[S\lambda_a\lambda_b] = \frac{n-4}{n} \left( \frac{1}{(l^2+a^2)^2} - \frac{a^2}{(l^2+a^2)^3} \right) \text{tr}[S\lambda_a\lambda_b] \quad (2.42)$$

The integral  $\int \frac{d^n l}{(l^2+a^2)^\alpha}$  is called an Feynman integral Bollini and Giambiagi [1972] and its result is set to

$$i\pi^{n/2} \frac{\Gamma(\alpha - \frac{1}{2}n)}{\Gamma(\alpha)} (a^2)^{\frac{1}{2}n-\alpha} \quad (2.43)$$

The term  $\frac{n-4}{n} \frac{a^2}{(l^2+a^2)^3}$  will be a multiple of  $n-4$  after integration. Therefore, this term disappears in the limit  $n \rightarrow 4$ . The other term,  $\frac{1}{(l^2+a^2)^2}$ , induces a factor  $\Gamma(2 - \frac{n}{2})$  which approximates  $-\frac{2}{n-4}$  in the limit  $n \rightarrow 4$ . Therefore, there is a non-zero constant  $c$ , independent of  $x$  and  $y$ , such that

$$\lim_{n \rightarrow 4} \int \frac{d^n l}{2\pi^n} \frac{k_{\perp}^2}{(l^2+a^2)^3} = c. \quad (2.44)$$

The anomalous part  $M_{abS,an}^{\mu\nu}$  is equal to

$$M_{ab,an}^{\mu\nu} = 16c\epsilon^{\mu\nu\rho\sigma} q_{\rho} p_{\sigma} \cdot \int_0^1 dx \int_0^{1-x} dy \text{tr}[S\lambda_a\lambda_b] \quad (2.45)$$

$$= 8c \cdot \epsilon^{\mu\nu\rho\sigma} q_{\rho} p_{\sigma} \cdot \text{tr}[S\lambda_a\lambda_b]. \quad (2.46)$$

In the Abelian case, this does not vanish. For the non-Abelian case,  $M_{ab,an}^{\mu\nu}$  vanishes if and only if  $\text{tr}[S\lambda_a, \lambda_b] = 0$ . We summarize this result in a theorem:

**Theorem 2.2.** *The classical conservation laws for the chiral current does not hold in quantum field theory. Therefore, the chiral symmetry is called anomalous.*

### 2.2.2 Regularization using Pauli-Villars method

Another regularization method is the Pauli-Villars method. The idea is to subtract the same amplitude, but we replace the mass  $m$  of the Dirac fermion with some mass  $M$  (van Nieuwenhuizen [1989]). This is depicted in figure 2. The physical limit is when  $M$  tends to infinity, because in this limit the regulating particle vanishes.




Figure 2: *Pauli-Villars regularization. We subtract the same diagram from the original, but we replace the mass  $m$  of the Dirac fermion with some mass  $M$ . The physical limit is when  $M$  tends to infinity.*

In the example of the AVV-diagram, we show that this method indeed regulates the divergences. The denominator in equation 2.22 is of order  $l^6$ . Thus the current  $(p+q)_\rho M_{abS}^{\mu\nu\rho}$  diverges if the nominator is of degree  $l^2$  or larger. We compare the orders of  $l$  w.r.t. the orders of  $m$  and the number of gamma matrices. It is summarized in table 1. From this table we conclude that the only the mass independent part of the numerator need regularization. In this case the numerator of  $(p+q)_\rho M_{abS}^{\mu\nu\rho}$  equals

$$\text{tr} [\gamma_5 (\not{p} + \not{q}) (\not{l} - \not{q}) \gamma^\mu \not{l} \gamma^\nu (\not{l} + \not{p})] \cdot \text{tr}(S \lambda_a \lambda_b) \quad (2.47)$$

A simple calculation shows that

$Cl$	$m$	$l$	Vanishes?	Convergent?
3	$m^3$	$l^0$	Yes, by trace identities	-
4	$m^2$	$l^0 - l^1$	No	Yes, by power counting
5	$m^1$	$l^0 - l^2$	Yes, by trace identities	-
6	$m^0$	$l^1 - l^3$	Only after regularization	Only after regularization

Table 1: *Order analysis of the numerator in equation 2.22. For a given number of gamma matrices ( $Cl$ ), we give the possible orders of the mass  $m$  and the loop momenta  $l$ . We also state if the given Clifford order vanishes and if it is convergent.*

$$\begin{aligned}
& (p+q)_\rho M_{abS}^{\mu\nu\rho}(m)|_{m \text{ indep. num.}} - (p+q)_\rho M_{abS}^{\mu\nu\rho}(m)|_{M \text{ indep. num.}} = \\
& = i \int \frac{d^4 l}{(2\pi)^4} \text{tr} [\gamma_5 (\not{p} + \not{q}) (\not{l} - \not{q}) \gamma_\mu \not{l} \gamma_\nu (\not{l} + \not{p})] \cdot \text{Tr}[S \lambda_a \lambda_b] \times \\
& \quad \times \frac{1}{((l-q)^2 + m^2)(l^2 + m^2)((l+p)^2 + m^2)} \\
& \quad \times \frac{1}{((l-q)^2 + M^2)(l^2 + M^2)((l+p)^2 + M^2)} \\
& \quad \times (3l^4 * (M^2 - 3m^2) + l^4 \cdot \mathcal{O}(l^{-1}))
\end{aligned} \tag{2.48}$$

By power counting, we see that the integral in equation 2.48 converge. Hence, the Pauli-Villars method is a valid regularization method.

If one sets  $k_\perp = 0$  equation 2.25 is also valid for the Pauli-Villars regularization. Using the same argument as in dimensional regularization, we conclude that the first two terms in equation 2.25 vanishes in the integral. Therefore the physical current is proportional to  $\lim_{M \rightarrow \infty} \langle -2im\gamma_5 + 2iM\gamma_5 \rangle$  and the chiral current is anomalous when  $\lim_{M \rightarrow \infty} \langle 2iM\gamma_5 \rangle$  doesn't vanish. The amplitude  $\langle 2iM\gamma_5 \rangle$  is given by

$$\langle 2iM\gamma_5 \rangle = \int \frac{d^n l}{(2\pi)^n} \text{tr} \left[ 2iMS\gamma_5 \frac{-i(\not{l} - \not{q}) + M}{(l-q)^2 + M^2} \gamma_\mu \lambda_a \frac{-i\not{l} + M}{l^2 + M^2} \gamma_\nu \lambda_b \frac{-i(\not{l} + \not{p}) + M}{(l+p)^2 + M^2} \right] \tag{2.49}$$

From the trace identities<sup>4</sup> and the commutation rules we calculate the trace. Equation

<sup>4</sup>These are  $\text{tr}(\gamma_5) = 0$ ,  $\text{tr}(\gamma_5 \gamma_{\mu_1} \dots \gamma_{\mu_{2k+1}}) = 0$ ,  $\gamma(\gamma_5 \gamma_\mu \gamma_n u) = 0$  and  $\text{tr}(\gamma_5 \gamma_\mu \gamma_\nu \gamma_\rho \gamma_\sigma) = 4i\epsilon^{\mu\nu\rho\sigma}$ .

2.49 equals

$$\langle 2iM\gamma_5 \rangle = \int \frac{d^n l}{(2\pi)^n} \frac{8M^2 \epsilon^{\mu\nu\rho\sigma} p_\rho q_\sigma \cdot \text{tr}[S\lambda_a \lambda_b]}{((l-q)^2 + M^2)(l^2 + M^2)((l+p)^2 + M^2)}. \quad (2.50)$$

We use the same Feynman trick and Feynman integral we used in dimensional regularization. The integral over the denominator in equation 2.50 equals

$$\int d^4 k \frac{1}{(k^2 + \mu^2)^3} = I(4, 3) = i\pi^2 \frac{\Gamma(3-2)}{\Gamma(3)} (\mu^2)^{2-3} \quad (2.51)$$

and this is proportional to  $\frac{1}{M^2+\dots}$ . So in the limit  $M \rightarrow \infty$ , the term  $\langle 2iM\gamma_5 \rangle$  is not zero, but is a multiple of  $\epsilon^{\mu\nu\rho\sigma} q_\rho p_\sigma \cdot \text{Tr}[S\lambda_a \lambda_b]$ . This matches the result found in dimensional regularization.

## 2.3 Other triangle diagrams

Not only the AVV-diagram contains an anomaly. There is another one-loop diagram that shows that the chiral current is anomalous. It is the AAA diagram and it uses the same diagram shown in figure 1, but all external fields are axial-vector fields. We calculate the anomalous part of the part of the scattering using dimensional regularization. That is, we need to find

$$M_{abS,an}^{\mu\nu,AAA} := 2 \int \frac{d^n l}{(2\pi)^n} \text{tr} \left[ \gamma_5 \not{k}_\perp \frac{-i(\not{k} + \not{k}_\perp - \not{q}) + m}{(k + k_\perp - q)^2 + m^2} \gamma_\mu \gamma^5 \frac{-i(\not{k} + \not{k}_\perp) + m}{(k + k_\perp)^2 + m^2} \gamma_\nu \gamma^5 \times \right. \\ \left. \frac{-i(\not{k} + \not{k}_\perp + \not{p}) + m}{(k + k_\perp + p)^2 + m^2} \right] \text{tr}[S\lambda_a \lambda_b]. \quad (2.52)$$

The difference between equation 2.29 and 2.52 is that we have replaced  $\gamma^\mu$  and  $\gamma^\nu$  with  $\gamma^\mu \gamma_5$  resp.  $\gamma^\nu \gamma_5$ . This correspond with the replacement of the external vector fields with axial-vector fields. Using the trace identities we calculate the trace of the numerator and it equals

$$4ik_\perp^2 \epsilon^{\mu\nu\rho\sigma} (2k_\rho(p+q)_\sigma - q_\rho p_\sigma) \cdot \text{Tr}(S\lambda_a \lambda_b) \quad (2.53)$$

Using the same Feynman trick as in equation 2.34 and 2.35, we conclude that  $M_{Sab,an}^{\nu\nu,AAA}$  equals

$$16\epsilon^{\mu\nu\rho\sigma} \int_0^1 dx \int_0^{1-x} dy \int \frac{d^n l}{(2\pi)^n} \frac{k_\perp^2 \cdot (2(k-xp+yq)_\rho(p+q)_\sigma - q_\rho q_\sigma)}{(l^2 + a^2)^3} \text{tr}[S\lambda_a \lambda_b] \quad (2.54)$$

(Note that in the Feynman trick we shifted  $k$  and therefore we have the term  $k - xp + yq$  in equation 2.54.) This can be simplified if we use the antisymmetry of  $\epsilon$  and the fact that the linear order of  $k$  disappears. Hence equation 2.54 can be written as

$$16\epsilon^{\mu\nu\rho\sigma}q_\rho q_\sigma \int \frac{d^n l}{(2\pi)^n} \frac{k_\perp^2}{(l^2 + a^2)^3} \text{tr}[S\lambda_a\lambda_b] \cdot \int_0^1 dx \int_0^{1-x} dy (2(x+y) - 1) \quad (2.55)$$

Comparing equation 2.55 with 2.38, we see that the anomalous amplitude for the AVV- and the AAA-diagram differs by the constant

$$\frac{\int_0^1 dx \int_0^{1-x} dy (2(x+y) - 1)}{\int_0^1 dx \int_0^{1-x} dy} \quad (2.56)$$

and by integration over  $x$  and  $y$  we conclude that this constant equals  $1/3$ .

## 2.4 Anomalies in box- and pentagon-diagrams

Not only triangle diagrams, but also box and pentagon diagrams have anomalous behavior. In this paragraph we calculate the box and pentagon anomaly using dimensional regularization. The calculation is similar to the triangle diagram. However, we first perform order analysis such that the equations doesn't become page filling. We follow the following steps:

1. Get an expression for the scattering amplitude using Feynman rules and contract it with the momenta of the  $A$ -field. This expression is proportional to the chiral current.
2. Generalize  $\gamma_5(\not{p} + \not{q}) = -\gamma_5(\not{l} - \not{q} - im) - (\not{l} + \not{p} - im)\gamma_5 - 2im\gamma_5 + 2\gamma_5\not{k}_\perp$  for the box- and pentagon diagram so that we determine the anomalous part.
3. Apply the Feynman trick and make the denominator in the integrand symmetric.
4. Determine which orders of  $k_\perp$  and  $k$  do vanish and which do not.
5. Calculate the non-vanishing terms.

Consider the box- and pentagon diagrams shown in figure 3. We use the shorthand notation  $\hat{p}_i = p_1 + \dots + p_i$  and  $\hat{p}_0 = 0$ . From the Feynman rules we deduce an




Figure 3: Example of an anomalous scattering processes with four resp. five external fields. In these examples we assume that the top external field is an axial-vector field and the other external fields are vector fields.

expression for the scattering amplitude. These are

$$\begin{aligned}
 M_{abcS}^{\mu\nu\rho\lambda,4} &= \int \frac{d l^n}{(2\pi)^n} \text{Tr} \left[ \frac{\gamma_5 \gamma^\lambda S (\not{l} + im) \gamma^\mu \lambda_a (\not{l} + \hat{p}_1' + im) \gamma^\nu \lambda_b}{(l^2 + m^2)((l + \hat{p}_1)^2 + m^2)} \times \right. \\
 &\quad \left. \times \frac{(\not{l} + \hat{p}_2' + im) \gamma^\rho \lambda_c (\not{l} + \hat{p}_3' + im)}{((l + \hat{p}_2)^2 + m^2)((l + \hat{p}_3)^2 + m^2)} \right] \\
 M_{abcdS}^{\mu\nu\rho\sigma\lambda,5} &= \int \frac{d l^n}{(2\pi)^n} \text{Tr} \left[ \frac{\gamma_5 \gamma^\lambda S (\not{l} + im) \gamma^\mu \lambda_a (\not{l} + \hat{p}_1' + im) \gamma^\nu \lambda_b}{(l^2 + m^2)((l + \hat{p}_1)^2 + m^2)} \times \right. \\
 &\quad \left. \times \frac{(\not{l} + \hat{p}_2' + im) \gamma^\rho \lambda_c (\not{l} + \hat{p}_3' + im) \gamma^\sigma \lambda_d (\not{l} + \hat{p}_4' + im)}{((l + \hat{p}_2)^2 + m^2)((l + \hat{p}_3)^2 + m^2)((l + \hat{p}_4)^2 + m^2)} \right]
 \end{aligned} \tag{2.57}$$

The vertex  $\gamma_5 \gamma^\lambda$  is proportional to the chiral current. To show that these diagrams are anomalous, we need to contract these expressions with  $\hat{p}_{3,\lambda}$  resp.  $\hat{p}_{4,\lambda}$ . Then we can compare the results with classical mechanics.

We continue with the second step. Recall that in dimensional regularization,  $\gamma_5$  anti-commutes with  $\gamma_0, \dots, \gamma_3$ , but commutes with the other gamma matrices. If expand  $l$

into  $k + k_\perp$  with  $k \in \mathbb{R}^4$  and  $k_\perp \in \mathbb{R}^{n-4}$ , we get the identity

$$\begin{aligned}\gamma_5 \hat{p}'_3 &= -\gamma_5(\not{l} - im) - (\not{l} + \hat{p}'_3 - im)\gamma_5 - 2im\gamma_5 + 2\gamma_5 \not{k}_\perp \\ \gamma_5 \hat{p}'_4 &= -\gamma_5(\not{l} - im) - (\not{l} + \hat{p}'_4 - im)\gamma_5 - 2im\gamma_5 + 2\gamma_5 \not{k}_\perp\end{aligned}\quad (2.58)$$

The first three terms in equation 2.58 are in one-to-one correspondence with the classical current conservation law. Therefore the box and pentagon diagrams are anomalous if

$$\begin{aligned}M_{abcS,an}^{\mu\nu\rho,4} &= \int \frac{d l^n}{(2\pi)^n} \text{Tr} \left[ \frac{2\gamma_5 \not{k}_\perp S(\not{l} + im) \gamma^\mu \lambda_a (\not{l} + \hat{p}'_1 + im) \gamma^\nu \lambda_b}{(l^2 + m^2)((l + \hat{p}_1)^2 + m^2)} \times \right. \\ &\quad \left. \times \frac{(\not{l} + \hat{p}'_2 + im) \gamma^\rho \lambda_c (\not{l} + \hat{p}'_3 + im)}{((l + \hat{p}_2)^2 + m^2)((l + \hat{p}_3)^2 + m^2)} \right] \\ M_{abcdS,an}^{\mu\nu\rho\sigma,5} &= \int \frac{d l^n}{(2\pi)^n} \text{Tr} \left[ \frac{2\gamma_5 \not{k}_\perp S(\not{l} + im) \gamma^\mu \lambda_a (\not{l} + \hat{p}'_1 + im) \gamma^\nu \lambda_b}{(l^2 + m^2)((l + \hat{p}_1)^2 + m^2)} \times \right. \\ &\quad \left. \times \frac{(\not{l} + \hat{p}'_2 + im) \gamma^\rho \lambda_c (\not{l} + \hat{p}'_3 + im) \gamma^\sigma \lambda_d (\not{l} + \hat{p}'_4 + im)}{((l + \hat{p}_2)^2 + m^2)((l + \hat{p}_3)^2 + m^2)((l + \hat{p}_4)^2 + m^2)} \right]\end{aligned}\quad (2.59)$$

does not vanish.

In the next step we symmetrize the denominator by applying the Feynman trick. In the general case, the Feynman trick is

$$\frac{1}{A_1 \cdots A_m} = (m-1)! \int_0^1 d z_1 \cdots d z_m \frac{\delta(1 - \sum_i z_i)}{(\sum_i z_i A_i)^m}. \quad (2.60)$$

Comparing equation 2.59 and 2.60 we choose  $A_i = (k + \hat{p}_i)^2 + k_\perp^2 + m^2$ . The denominators in equation 2.59 can be written as

$$\begin{aligned}\left(\sum_i A_i z_i\right)^m &= \left(\sum_i z_i (k^2 + k_\perp^2 + m^2 + 2k \cdot \hat{p}_i + \hat{p}_i^2)\right)^m \\ &= \left(\left(k + \sum_i z_i \hat{p}_i\right)^2 + k_\perp^2 + m^2 + \sum_i z_i \hat{p}_i^2 - \left(\sum_i z_i \hat{p}_i\right)^2\right)^m\end{aligned}\quad (2.61)$$

Let  $\mu^2 = m^2 + \sum_i z_i \hat{p}_i^2 - (\sum_i z_i \hat{p}_i)^2$  and shift  $k$  to  $k - \sum_i z_i \hat{p}_i$ . If we set  $B_i =$

$\hat{p}_i - \sum_j z_j \hat{p}_j$ , equation 2.59 simplifies to

$$\begin{aligned} M_{abcS,an}^{\mu\nu\rho,4} &= \int \frac{d^n l}{(2\pi)^n} \text{Tr} \left[ \frac{2\gamma_5 k_\perp S(\not{l} + \not{B}_0 + im)\gamma^\mu \lambda_a(\not{l} + \not{B}_1 + im)\gamma^\nu \lambda_b}{(l^2 + \mu^2)^4} \times \right. \\ &\quad \left. \times (\not{l} + \not{B}_2 + im)\gamma^\rho \lambda_c(\not{l} + \not{B}_3 + im) \right] \\ M_{abcdS,an}^{\mu\nu\rho\sigma,5} &= \int \frac{d^n l}{(2\pi)^n} \text{Tr} \left[ \frac{2\gamma_5 k_\perp S(\not{l} + \not{B}_0 + im)\gamma^\mu \lambda_a(\not{l} + \not{B}_1 + im)\gamma^\nu \lambda_b}{(l^2 + \mu^2)^5} \times \right. \\ &\quad \left. \times (\not{l} + \not{B}_2 + im)\gamma^\rho \lambda_c(\not{l} + \not{B}_3 + im)\gamma^\sigma \lambda_d(\not{l} + \not{B}_4 + im) \right]. \end{aligned} \quad (2.62)$$

Note that the denominator is symmetric in  $k$  and  $k_\perp$ . The numerator is symmetric if the terms are a multiple of  $k_\perp^2 (l^2)^m$  for some  $m \in \mathbb{N}$ . For these terms we can use the following symmetry argument.

$$\int \frac{d^n l}{(2\pi)^n} \frac{k_\perp^2 l^{2m}}{(l^2 + \mu^2)^d} = \frac{n-4}{n} \int \frac{d^n l}{(2\pi)^n} \frac{l^{2m+2}}{(l^2 + \mu^2)^d} \quad (2.63)$$

We expand the numerator of equation 2.62 in a polynomial in  $l$ . In table 2 and 3 we determine which orders will vanish. There we notice that all terms are a multiple of  $\frac{(\mu^2)^\alpha}{(l^2 + \mu^2)^\beta}$ . In dimensional regularization this integrates to  $i\pi^{n/2} \frac{\Gamma(\beta - n/2)}{\Gamma(\beta)} (\mu^2)^{n/2 + \alpha - \beta}$ . By the following proposition, we show that for some values of  $\beta$  it vanishes.

**Proposition 2.3.** *Let  $\beta \in \mathbb{N}$  and  $f$  a smooth map. The limit*

$$\lim_{n \rightarrow 4} \frac{n-4}{n} \cdot i\pi^{\frac{n}{2}} \frac{\Gamma(\beta - \frac{n}{2})}{\Gamma(\beta)} \int_0^1 dz_1 \dots dz_m \delta \left( 1 - \sum_i z_i \right) f(z_i) \cdot (\mu^2)^{\frac{n}{2} + \alpha - \beta}$$

*is zero if  $\beta > 2$  and converges if  $\beta = 0$ .*

We postpone the proof of this proposition to the end of this chapter. From table 2 and 3 we conclude that the only non-vanishing terms of equation 2.62 are

$$\begin{aligned} M_{abcS,an}^{\mu\nu\rho,4} &= \int \frac{d^n l}{(2\pi)^n} \frac{\text{Tr}[S\lambda_a\lambda_b\lambda_c]}{(l^2 + \mu^2)^4} \times \\ &\quad \times (\text{Tr}[2\gamma_5 k_\perp (\not{B}_0 + im)\gamma^\mu \not{l} \gamma^\nu \not{l} \gamma^\rho] + \text{Tr}[2\gamma_5 k_\perp \not{l} \gamma^\mu (\not{B}_1 + im)\gamma^\nu \not{l} \gamma^\rho] + \\ &\quad \text{Tr}[2\gamma_5 k_\perp \not{l} \gamma^\mu \not{l} \gamma^\nu (\not{B}_2 + im)\gamma^\rho] + \text{Tr}[2\gamma_5 k_\perp \not{l} \gamma^\mu \not{l} \gamma^\nu \not{l} \gamma^\rho (\not{B}_3 + im)]) \\ M_{abcdS,an}^{\mu\nu\rho\sigma,5} &= \int \frac{d^n l}{(2\pi)^n} \frac{\text{Tr}[S\lambda_a\lambda_b\lambda_c\lambda_d]}{(l^2 + \mu^2)^5} \text{Tr}[2\gamma_5 k_\perp \not{l} \gamma^\mu \not{l} \gamma^\nu \not{l} \gamma^\rho \not{l} \gamma^\sigma]. \end{aligned} \quad (2.64)$$

Order(s) of $l$	Vanishes?
0,2,4	Yes, Then the numerator of equation 2.62 is antisymmetric in $k$ or $k_{\perp}$ .
1	<p>Yes, the first order of equation 2.62 is proportional to</p> $\frac{k_{\perp}^2}{(l^2 + \mu^2)^4} = \frac{n-4}{n} \frac{l^2}{(l^2 + \mu^2)^4} \quad (2.65)$ $= \frac{n-4}{n} \left( \frac{1}{(l^2 + \mu^2)^3} - \frac{\mu^2}{(l^2 + \mu^2)^4} \right) \quad (2.66)$ <p>By Proposition 2.3 this vanishes.</p>
3	No

Table 2: Order analysis of equation 2.62 for the box diagram. We expand this equation in  $l$  and we determine which orders vanishes.

Using the Clifford algebra, we work out the spin traces. We conclude that for the box diagram this trace is a multiple of  $\epsilon^{\mu\nu\rho\tau} p_{(i),\tau}$ . For the pentagon diagram equation 2.64 is a multiple of  $\epsilon^{\mu\nu\rho\sigma}$ .

We finish this paragraph with the proof of Proposition 2.3. We simplify our notation with  $\int_0^1 dz_1 \dots dz_m \delta(1 - \sum_i z_i) = \int_{\Delta} dz_i$ . Observe that

$$\lim_{n \rightarrow 4} \frac{n-4}{n} \cdot i\pi^{\frac{n}{2}} \frac{\Gamma(\beta - \frac{n}{2})}{\Gamma(\beta)} \int_{\Delta} dz_i f(z_i) \cdot (\mu^2)^{\frac{n}{2} - \beta} \quad (2.71)$$

$$= \left( \lim_{n \rightarrow 4} \frac{-2i\pi^{\frac{n}{2}}}{n\Gamma(\beta)} \right) \left( \lim_{n \rightarrow 4} (2 - n/2) \Gamma(\beta - n/2) \right) \left( \lim_{n \rightarrow 4} \int_{\Delta} dz_i f(z_i) \cdot (\mu^2)^{\frac{n}{2} - \beta} \right) \quad (2.72)$$

The first part of equation 2.72 can be easily calculated and is equal to  $\frac{-i\pi^2}{2\Gamma(\beta)}$ . The second term vanishes if  $\beta > 2$ . If  $\beta = 2$ , the second term does not vanish, but is equal to

$$\lim_{n \rightarrow 4} (2 - n/2) \Gamma(2 - n/2) = \lim_{n \rightarrow 4} \Gamma(3 - n/2) = 1. \quad (2.73)$$

We finally focus on the last term of equation 2.72. We need to show that this integral

Order(s) of $l$	Vanishes?
0,2,4	Yes, Then the numerator of equation 2.62 is antisymmetric in $k$ or $k_{\perp}$ .
1	<p>Yes, the first order of equation 2.62 is proportional to</p> $\frac{k_{\perp}^2}{(l^2 + \mu^2)^5} = \frac{n-4}{n} \frac{l^2}{(l^2 + \mu^2)^5} \quad (2.67)$ $= \frac{n-4}{n} \left( \frac{1}{(l^2 + \mu^2)^4} - \frac{\mu^2}{(l^2 + \mu^2)^5} \right) \quad (2.68)$ <p>By Proposition 2.3 this vanishes.</p>
3	<p>Yes, the third order of equation 2.62 is proportional to</p> $\frac{k_{\perp}^2 l^2}{(l^2 + \mu^2)^5} = \frac{n-4}{n} \frac{l^4}{(l^2 + \mu^2)^5} \quad (2.69)$ $= \frac{n-4}{n} \left( \frac{1}{(l^2 + \mu^2)^3} - \frac{2\mu^2}{(l^2 + \mu^2)^4} + \frac{\mu^4}{(l^2 + \mu^2)^5} \right) \quad (2.70)$ <p>By Proposition 2.3 this vanishes.</p>
5	No

Table 3: *Order analysis of equation 2.62 for the pentagon diagram. We expand this equation in  $l$  and we determine which orders vanishes.*

converges. Recall that

$$\mu^2 - m^2 = \sum_i z_i \hat{p}_i^2 - \left( \sum_i z_i \hat{p}_i \right)^2 \geq 0. \quad (2.74)$$

Hence the map  $f \cdot (\mu^2)^{n/2-\beta}$  does not have  $\frac{1}{0}$  behavior on  $\Delta$ . This concludes that  $f \cdot (\mu^2)^{n/2-\beta}$  is integrable for each value of  $n \in \mathbb{R}$ . This proves the proposition.  $\square$

## 2.5 Non-Abelian Anomalies in Feynman diagrams

Till now, we ignored the traces like  $\text{Tr}(S\lambda_a \dots \lambda_c)$ . These terms are determined by the properties of the gauge group. It turns out that in some cases these terms vanishes.

We give two examples of this phenomena. In both cases we assume Bose symmetry. Recall that this is the symmetry when one interchanges external lines.

**Example 2.4.** If  $S = 1$ , then the pentagon anomaly vanishes due to Bose symmetry.

Indeed, The pentagon anomaly is proportional to

$$\epsilon^{\mu\nu\rho\sigma} \cdot \text{tr}(S\lambda_a\lambda_b\lambda_c\lambda_d) \quad (2.75)$$

Bose symmetry requires that the above expression is symmetric under the permutation of the pairs  $(a, \mu)$ ,  $(b, \nu)$ ,  $(c, \rho)$  and  $(d, \sigma)$ . By adding all the permuted versions of equation 2.75 we get an expression with 24 terms, which can be written as

$$\epsilon^{\mu\nu\rho\sigma} \cdot \text{tr}(S\{[\lambda_a, \lambda_b], [\lambda_c, \lambda_d]\} + S\{[\lambda_a, \lambda_c], [\lambda_d, \lambda_b]\} + S\{[\lambda_a, \lambda_d], [\lambda_b, \lambda_c]\}) \quad (2.76)$$

We now assume that  $S = 1$ . Note that due to the cyclic property of the trace, the anti-commutator in equation 2.76 simplifies to

$$2\epsilon^{\mu\nu\rho\sigma} \cdot \text{tr}([\lambda_a, \lambda_b][\lambda_c, \lambda_d] + [\lambda_a, \lambda_c][\lambda_d, \lambda_b] + [\lambda_a, \lambda_d][\lambda_b, \lambda_c]) \quad (2.77)$$

and this can be written as

$$2\epsilon^{\mu\nu\rho\sigma} \cdot \text{tr}(\lambda_a([\lambda_b, [\lambda_c, \lambda_d]] + [\lambda_c, [\lambda_d, \lambda_b]] + [\lambda_d, [\lambda_b, \lambda_c]])) \quad (2.78)$$

From the Jacobi identity we conclude that the Abelian pentagon anomaly vanishes.

**Example 2.5.** If all triangle anomalies vanishes, then the box and pentagon anomalies vanishes due to Bose symmetry.

Under Bose symmetry the triangle anomaly is proportional to

$$\epsilon^{\mu\nu\rho\sigma} q_\rho p_\sigma \cdot \text{tr}(S\lambda_a\lambda_b) + \epsilon^{\nu\mu\rho\sigma} p_\rho q_\sigma \cdot \text{tr}(S\lambda_b\lambda_a) \quad (2.79)$$

which equals  $\epsilon^{\mu\mu\rho\sigma} q_\rho p_\sigma \cdot \text{tr}(S\{\lambda_a, \lambda_b\})$ . Hence all triangle anomalies vanishes if and only if  $\text{tr}(\{\lambda_a, \lambda_b\}\lambda_c) = 0$  for all generators of the lie algebra  $\lambda_a, \lambda_b, \lambda_c$ . We call  $\text{tr}(\{\lambda_a, \lambda_b\}\lambda_c)$  to be the *d-symbol* and we denote it as  $d_{abc}$ . We show that the box and pentagon anomalies can be written as linear combinations of *d-symbols*.

Assume that  $S = \lambda_e$ . We can write equation 2.76 in terms of  $d$ -symbols using the structure constants<sup>5</sup>  $f_{ab}^c$ :

$$\epsilon^{\mu\nu\rho\sigma} \cdot (f_{ab}^f f_{cd}^g + f_{ac}^f f_{db}^g + f_{ad}^f f_{bc}^g) d_{efg} \quad (2.80)$$

Clearly this vanishes if the  $d$ -symbol vanishes. We repeat the calculation for the box diagram. Under Bose symmetry, the anomaly is proportional to

$$\begin{aligned} & \epsilon^{\mu\nu\rho\tau} p_{(i),\tau} \cdot \text{tr}(\lambda_e \lambda_a \lambda_b \lambda_c) + \epsilon^{\mu\nu\rho\tau} p_{(i),\tau} \cdot \text{tr}(\lambda_e \lambda_a \lambda_b \lambda_c) \\ & + \epsilon^{\nu\rho\mu\tau} p_{(i),\tau} \cdot \text{tr}(\lambda_e \lambda_b \lambda_c \lambda_a) + \epsilon^{\nu\mu\rho\tau} p_{(i),\tau} \cdot \text{tr}(\lambda_e \lambda_b \lambda_c \lambda_a) \\ & + \epsilon^{\rho\mu\nu\tau} p_{(i),\tau} \cdot \text{tr}(\lambda_e \lambda_c \lambda_a \lambda_b) + \epsilon^{\rho\nu\mu\tau} p_{(i),\tau} \cdot \text{tr}(\lambda_e \lambda_c \lambda_a \lambda_b). \end{aligned} \quad (2.81)$$

We write this in terms of (anti)-commutators

$$\frac{1}{2} \epsilon^{\mu\nu\rho\sigma} p_{(i),\sigma} \text{tr}(\lambda_d(\{\lambda_a, [\lambda_b, \lambda_c]\}) + \{\lambda_b, [\lambda_c, \lambda_a]\} + \{\lambda_c, [\lambda_a, \lambda_b]\}) \quad (2.82)$$

This is a linear combination of  $d$ -symbols, because it equals

$$\frac{1}{2} \epsilon^{\mu\nu\rho\sigma} p_{(i),\sigma} (f_{bc}^e d_{aed} + f_{ca}^e d_{bed} + f_{ab}^e d_{ced}) \quad (2.83)$$

So when the  $d$ -symbol vanishes, the box and pentagon anomaly also vanishes. Georgi and Glashow calculated the  $d$ -symbol for many matrix groups and showed that in most cases the  $d$ -symbol vanishes. For more information, see Georgi and Glashow [1972].

---

<sup>5</sup>Recall:  $f_{ab}^c$  are uniquely defined by  $[\lambda_a, \lambda_b] = f_{ab}^c \lambda_c$

### 3 Calculating anomalies using path integrals

In the previous chapter, we have seen that anomalies are classical conservation laws that are broken when one considers its results in one-loop diagrams. In this chapter we examine another point of view, namely we show that anomalies arise due to the fact that the path integral measure is not invariant under the symmetry.

To show this, we first consider the simple case of massless Dirac fermions in the path integral formalism. Using the method introduced by Fujikawa, we again show that the chiral symmetry is anomalous. After this, we look at other examples and we relate them to the results found when using perturbation theory.

Finally we analyze Fujikawa's method from a more mathematical perspective. We show that the anomaly only depends on the topology of the gauge bundle and is a specific application of the Atiyah-Singer index theorem. In the next chapters we prove the index theorem using Fujikawa's method.

#### 3.1 The Fujikawa method

In classical mechanics we investigated how the action  $D$  changed under symmetry transformations of the fields  $\psi$ . From this we deduced conserved quantities we called Noethers currents. In quantum field theory the dynamics of a field are not determined by the action, but by the generating functional

$$\mathcal{Z} = \int \mathcal{D}\bar{\psi} \mathcal{D}\psi \exp(iS[\psi]). \quad (3.1)$$

Here  $\int \mathcal{D}\psi$  is the path integral over all fields  $\psi$ . With Fujikawa's method (Fujikawa [1980], Fujikawa and Suzuki [2004]) we mimic Noethers theorem for generating functionals. We get a result that differs from the classical theory and this difference is the anomaly. As an example we consider a Dirac fermion  $\psi$  of mass  $m$  in quantum electrodynamics. Let  $A_\mu$  be the electromagnetic gauge potential and let  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$  be the field strength. The action for this fermion is given by

$$S = \int_{\mathbb{R}^4} d^4x \bar{\psi} i \gamma^\mu (\partial_\mu - i A_\mu) \psi - m \bar{\psi} \psi - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \quad (3.2)$$

As usual we work with a flat four dimensional spacetime, but instead of using the Minkowski metric, we use the Euclidean metric. For this we need to replace the  $i$  with  $-1$  in equation 3.2. We use the following properties of path integrals:

1. The path integral is different for fermionic than for bosonic particles. For fermionic particles physicists use Berezin integration for which the Jacobian is given by the inverse of regular Jacobian.
2. The Dirac field  $\psi$  is actually a bispinor. That is,  $\mathbb{C}^4$  has 2 irreducible spin-representations and thus describes 2 spin particles. Therefore we treat  $\bar{\psi}$  and  $\psi$  as separate particles and so we integrate them separately.

We assume that the Dirac fermion is invariant under the transformation

$$\psi \mapsto e^{i\alpha(x)\gamma_5} \psi \quad (3.3)$$

$$\bar{\psi} \mapsto \bar{\psi} e^{i\alpha(x)\gamma_5} \quad (3.4)$$

where  $\alpha$  is an arbitrary but infinitely small real-valued map on  $\mathbb{R}^4$ . Under this symmetry the action  $S$  changes into

$$\tilde{S} = S + \int_{\mathbb{R}^4} d^4 x \alpha(x) [\partial_\mu (\bar{\psi} \gamma^\mu \gamma_5 \psi) + 2im \bar{\psi} \gamma_5 \psi] \quad (3.5)$$

If we denote the Jacobian of the path integral as  $J_\psi$ , then the generating functional  $\mathcal{Z}$  transforms into

$$\tilde{\mathcal{Z}} = \int \mathcal{D}\bar{\psi} \mathcal{D}\psi J_{\bar{\psi}} J_\psi \exp \left( -S[\psi] - \int_{\mathbb{R}^4} d^4 x \alpha(x) [\partial_\mu (\bar{\psi} \gamma^\mu \gamma_5 \psi) + 2im \bar{\psi} \gamma_5 \psi] \right). \quad (3.6)$$

Clearly, we get the classical chiral current conservation if and only if the Jacobians are equal to one. To show that the chiral symmetry is anomalous we need to calculate the Jacobians. For this we must assume that  $\alpha$  is small enough such that it behaves as a constant. Formally, the Jacobian is given by

$$J_\psi \equiv \det^{-1} \exp(i\alpha(x)\gamma_5) = \exp(-i \text{Tr}(\alpha(x)\gamma_5)) + \mathcal{O}(\dot{\alpha}, \alpha^2) \quad (3.7)$$

where  $\text{Tr}$  is the trace over spacetime and all spin indices. The expression is exactly the same for  $J_{\bar{\psi}}$  and therefore is the combined Jacobian

$$J = \exp(-2i \text{Tr}(\alpha(x)\gamma_5)) + \mathcal{O}(\dot{\alpha}, \alpha^2) \quad (3.8)$$

To evaluate this explicitly, we consider an orthonormal basis of eigenvectors  $\phi_n$  for the operator  $D = \gamma^\mu (\partial_\mu - iA_\mu)$ . We denote the corresponding eigenvalues with  $\lambda_n$ . Formally the trace equals  $\text{Tr}(\cdot) = \sum_n \int d^4 x \langle \phi_n(x) | \cdot | \phi_n(x) \rangle$  and so

$$J = \exp \left[ -2i \sum_n \int d^4 x \langle \phi_n(x) | \alpha(x) \gamma_5 | \phi_n(x) \rangle \right] + \mathcal{O}(\dot{\alpha}, \alpha^2) \quad (3.9)$$

This is a formal calculation and may not converge. Indeed, there may be infinitely many eigenspaces and each eigenspace may be infinite dimensional. It is also possible that the eigenvectors are not normalizable. That is, they diverge when we integrate over them. We tackle this problem by modifying the Jacobian slightly. For this observe the following: Recall that the sum over an infinite sequence  $\{c_n\}$  is defined as the limit of the partial sums  $\sum_{n=0}^M c_n$ . Consider the step function  $\theta: \mathbb{R} \rightarrow \{0, 1\}$  with  $\theta(x) = 1$  if  $x \leq 1$  and  $\theta(x) = 0$  if  $x > 1$ . The partial sum equals

$$\sum_{n=0}^M c_n = \sum_{n=0}^M c_n \theta\left(\frac{n}{M}\right) = \sum_{n=0}^{\infty} c_n \theta\left(\frac{n}{M}\right) \quad (3.10)$$

We modify the Jacobian by replacing the step function with another smooth map  $f(x)$  that rapidly approaches zero when  $x$  is large and  $f(0) = 1$ . See figure 4. For a well-chosen map  $f$  the Jacobian converges absolutely. An example is  $e^{-x}$ . We study this regulator more thoroughly in chapter 4. In this chapter we investigate the second method and we assume that the combined Jacobian equals

$$J = \exp \left[ -2i \lim_{t \rightarrow 0} \sum_n \int d^4 x \alpha(x) \langle \phi_n(x) | \gamma_5 f(t \cdot \lambda_n^2) | \phi_n(x) \rangle \right] + \mathcal{O}(\dot{a}, a^2) \quad (3.11)$$

Using functional calculus we rewrite equation 3.11 as




Figure 4: The regulator  $f(x)$ , used in the Fujikawa method, mimics the behavior of the step function. It must value 1 at  $x = 0$  and it must rapidly decrease at infinity.

$$\log J = -2i \lim_{t \rightarrow 0} \sum_n \int d^4 x \alpha(x) \langle \phi_n(x) | \gamma_5 f(t \cdot D^2) | \phi_n(x) \rangle + \mathcal{O}(\dot{a}, a^2) \quad (3.12)$$

$$= -2i \lim_{t \rightarrow 0} \text{Tr} [\gamma_5 \alpha \cdot f(t \cdot D^2)] + \mathcal{O}(\dot{a}, a^2). \quad (3.13)$$

Next we perform a change of basis. Namely, we expand the Jacobian in terms of the plane-waves  $e^{ikx}$ . Because plane-waves form only a basis over  $\mathbb{R}^4$ , we still have to take the trace over the spin indices. We denote the trace over the spin indices as  $\text{tr}$ . In the plane-wave basis equation 3.13 becomes

$$\log J = -2i \lim_{t \rightarrow 0} \int \frac{d^4 k}{(2\pi)^4} \langle k | \text{tr} [\gamma_5 \alpha \cdot f(t \cdot D^2)] | k \rangle \quad (3.14)$$

$$= -2i \lim_{t \rightarrow 0} \int d^4 x \alpha(x) \int \frac{d^4 k}{(2\pi)^4} e^{-ik \cdot x} \text{tr} [\gamma_5 f(t \cdot D^2)] e^{ik \cdot x}. \quad (3.15)$$

Note that the operator  $D$  is the Dirac operator with respect to the covariant derivative  $\nabla_\mu = \partial_\mu - iA_\mu$ . Theorem 1.22 states that  $D^2 = -\nabla^\mu \nabla_\mu - \frac{i}{4}[\gamma^\mu, \gamma^\nu] F_{\mu\nu}$  and so

$$\log J = -2i \lim_{t \rightarrow 0} \int d^4 x \alpha(x) \int \frac{d^4 k}{(2\pi)^4} e^{-ik \cdot x} \text{tr} \left[ \gamma_5 f \left( -t \nabla^\mu \nabla_\mu - \frac{it}{4} [\gamma^\mu, \gamma^\nu] F_{\mu\nu} \right) \right] e^{ik \cdot x}. \quad (3.16)$$

We simplify the equation 3.16 by pulling  $e^{ik \cdot x}$  to the left. For this we use the Leibniz rule  $[\nabla_\mu, e^{ik \cdot x}] = ik_\mu$  and hence

$$\log J = -2i \lim_{t \rightarrow 0} \int d^4 x \alpha(x) \int \frac{d^4 k}{(2\pi)^4} \text{tr} \left[ \gamma_5 f \left( -t(\nabla^\mu + ik_\mu)^2 - \frac{it}{4} [\gamma^\mu, \gamma^\nu] F_{\mu\nu} \right) \right]. \quad (3.17)$$

We calculate the non vanishing part of this integral. First we rescale  $k_\mu$  by  $t^{-1/2}k_\mu$  and the integrand of  $\int d^4 x$  in equation 3.17 equals

$$\int \frac{d^4 k}{(2\pi)^4} t^{-2} \text{tr} \left[ \gamma_5 f \left( -(t^{1/2} \nabla_\mu + ik_\mu)^2 - \frac{it}{4} [\gamma^\mu, \gamma^\nu] F_{\mu\nu} \right) \right] \quad (3.18)$$

We consider the Taylor series of  $f$  in  $t^{\frac{1}{2}}$  at  $t = 0$ . We only have to consider the first four orders, because all higher orders are linear to  $t^{1/2}$  and vanishes when we perform the limit  $t \rightarrow 0$ . In table 4 we analyze these orders and we see that only the  $t$ -constant term doesn't vanish. Hence, when  $t$  tends to zero, then equation 3.18 equals

$$-\frac{1}{32} \int \frac{d^4 k}{(2\pi)^4} f''(k^2) \text{tr} [\gamma_5 [\gamma^\mu, \gamma^\nu] [\gamma^\rho, \gamma^\sigma]] F_{\mu\nu} F_{\rho\sigma} \quad (3.19)$$

$t$	Clifford order	Vanishes?
$t^{-2}$	$\text{tr}(\gamma_5)$	Yes
$t^{-3/2}$	$\text{tr}(\gamma_5)$	Yes
$t^{-1}$	$\text{tr}(\gamma_5)$	Yes
	$\text{tr}(\gamma_5 \gamma^\mu \gamma^\nu)$	Yes
$t^{-1/2}$	$\text{tr}(\gamma_5)$	Yes
	$\text{tr}(\gamma_5 \gamma^\mu \gamma^\nu)$	Yes
$t^0$	$\text{tr}(\gamma_5)$	Yes
	$\text{tr}(\gamma_5 \gamma^\mu \gamma^\nu)$	Yes
	$\text{tr}(\gamma_5 \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma)$	No

Table 4: *Order analysis of the Taylor approximation of equation 3.18 in  $t^{1/2}$  at  $t = 0$ . For a given order of  $t^{1/2}$ , we give the possible traces over the spin indices. Using the trace identities we conclude if a given order vanishes*

From the trace identity  $\gamma(\gamma_5 \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma) = 4\epsilon^{\mu\nu\rho\sigma}$  where  $\epsilon$  is the Levi-Civita symbol, the regularized Jacobian becomes

$$\log J = \frac{i}{4} e^{\mu\nu\rho\sigma} \int d^4 x \alpha(x) \int \frac{d^4 k}{(2\pi)^4} f''(k^2) F_{\mu\nu} F_{\rho\sigma} \quad (3.20)$$

$$= \frac{i}{16} e^{\mu\nu\rho\sigma} \int d^4 x \alpha(x) F_{\mu\nu} F_{\rho\sigma}. \quad (3.21)$$

In the last step we used the that  $f(0) = 1$  and  $f(\infty) = 0$ . We use equation 3.21 in the transformed generating functional and so equation 3.6 equals

$$\tilde{\mathcal{Z}} = \int \mathcal{D}\bar{\psi} \mathcal{D}\psi \exp \left( -S[\psi] - \int_{\mathbb{R}^4} d^4 x \alpha(x) \left[ \partial_\mu (\bar{\psi} \gamma^\mu \gamma_5 \psi) + 2im \bar{\psi} \gamma_5 \psi - \frac{i}{16} e^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma} \right] \right). \quad (3.22)$$

Because  $\mathcal{Z}$  is conserved under the chiral transformation we get the conservation law

$$\partial_\mu (\bar{\psi} \gamma^\mu \gamma_5 \psi) = -2im \bar{\psi} \gamma_5 \psi + \frac{i}{16} e^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma} \quad (3.23)$$

This shows that the chiral current is anomalous.

### 3.2 The covariant anomaly

Previously, we have only considered Fujikawas method for abelian anomalies. It is natural to ask whether this method also works for non-abelian gauge fields. Most steps we can copy directly, but there are some subtleties we have to be aware of. For reference see Bertlmann [1996], which give a detailed overview of different non-Abelian anomalies.

In the rest of this paragraph we consider the chiral symmetry for a fermion  $\psi$  with mass  $m$  we studied in chapter 2.1. The action  $S[\psi]$  is given by

$$S[\psi] = \int_{\mathbb{R}^4} -\bar{\psi}(\not{\partial} + m)\psi + i\bar{\psi}(V + \not{A}\gamma_5)\psi \, dx \quad (3.24)$$

where  $V_\mu$  and  $A_\mu$  are gauge fields. In the case of the chiral symmetry we assume that the physical system is invariant under the transformation

$$\psi \mapsto e^{i\alpha(x)S\gamma_5}\psi \quad \bar{\psi} \mapsto \bar{\psi}e^{i\alpha(x)S\gamma_5}. \quad (3.25)$$

Here  $\alpha$  is a small real valued function. When we consider the Abelian chiral symmetry we assume that  $S$  equals the identity. Otherwise we assume that  $S$  is a generator of the Lie algebra. We rewrite the action and the symmetry in terms of the operators  $P_\pm = \frac{1}{2}(1 \pm \gamma_5)$ . These operators project the fields into the  $\pm 1$  eigenspace of  $\gamma_5$ . Because they are projection operators they obey the properties

$$P_\pm^2 = P_\pm, \quad P_+P_- = P_-P_+ = 0, \quad P_+ + P_- = 1, \quad \text{and} \quad \gamma_5 P_\pm = \pm P_\pm. \quad (3.26)$$

We write  $\psi_\pm$  for the projection of  $\psi$  to the  $\pm 1$  eigenspace of  $\gamma_5$ . In terms of these new fields the action can be written as

$$S[\psi] = \int_{\mathbb{R}^4} -\bar{\psi}_+(\not{\partial} + m - iV - i\not{A})\psi_+ + -\bar{\psi}_-(\not{\partial} + m - iV + i\not{A})\psi_- \, dx \quad (3.27)$$

If we denote  $A^\pm = V \pm A$  and define  $D_\pm = \not{\partial} - i\not{A}^\pm$  the action splits into the action of two non-interacting Dirac particles. That is, the action becomes

$$S[\psi_+, \psi_-] = \int_{\mathbb{R}^4} -\bar{\psi}_+(D_+ + m)\psi_+ + -\bar{\psi}_-(D_- + m)\psi_- \, dx. \quad (3.28)$$

Using Fujikawas method we calculate how the generating functional changes under the chiral symmetry. The generating functional  $\mathcal{Z} = \int \mathcal{D}\bar{\psi}_+ \mathcal{D}\psi_+ \mathcal{D}\bar{\psi}_- \mathcal{D}\psi_- \exp[-S]$  transforms under equation 3.25 into

$$\tilde{\mathcal{Z}} = \int \mathcal{D}\bar{\psi}_+ \mathcal{D}\psi_+ \mathcal{D}\bar{\psi}_- \mathcal{D}\psi_- J_{\bar{\psi}_+} J_{\psi_+} J_{\bar{\psi}_-} J_{\psi_-} \exp[-\tilde{S}] \quad (3.29)$$

where  $\tilde{S}$  equals

$$\tilde{S} = S + \int_{\mathbb{R}^4} d^4 x \alpha(x) \left[ \partial_\mu (\bar{\psi} \gamma^\mu \gamma_5 S \psi) + \bar{\psi} \gamma^\mu \gamma_5 (iV_\mu^b + iA_\mu^b \gamma_5) [S, \lambda_b] \psi - 2im \bar{\psi} \gamma_5 S \psi \right] \quad (3.30)$$

The transformed action  $\tilde{S}$  was found using Noethers theorem and was calculated in equation 2.21. We note that  $P_\pm \gamma_5 = \gamma_5 P_\pm$  and hence the chiral symmetry equals

$$\psi_\pm \mapsto e^{i\alpha(x)S\gamma_5} \psi_\pm \quad \bar{\psi}_\pm \mapsto \bar{\psi}_\pm e^{i\alpha(x)S\gamma_5}. \quad (3.31)$$

By definition of the fermionic path integral the Jacobian equals

$$J_{\bar{\psi}_\pm} = J_{\psi_\pm} = \det^{-1} \exp(i\alpha S\gamma_5) \quad (3.32)$$

If we assume that  $\alpha$  and  $\dot{\alpha}$  are infinitely small, we get by the Baker–Campbell–Hausdorff formula

$$J_{\bar{\psi}_\pm} = J_{\psi_\pm} = \exp \text{Tr}(-i\alpha S\gamma_5) + \mathcal{O}(\alpha^2, \dot{\alpha}). \quad (3.33)$$

Again, this might diverge and we need to regulate this. We regulate the same way as we regulated the Jacobian in the previous paragraph. For this we need to pick a real valued maps  $f_+, f_-: \mathbb{R} \rightarrow \mathbb{R}$  that vanishes at infinity and  $f_\pm(0) = 1$ . We regulate  $J_{\bar{\psi}_\pm}$  and  $J_{\psi_\pm}$  with  $D_\pm$ . The combined regularized Jacobian equals

$$J = \lim_{t_\pm \rightarrow 0} \exp \text{Tr} \left[ -2i\alpha S\gamma_5 \cdot (f_+(t_+ D_+^2) + f_-(t_- D_-^2)) \right] \quad (3.34)$$

Here  $\text{Tr}$  denotes the trace over the fields, spin-indices and the Lie algebra of the gauge group. We split this trace into the trace  $\text{tr}_g$  over the Lie algebra and the trace  $\text{Tr}$  over the fields and spin indices. Equation 3.34 then equals

$$\log J = \text{tr}_g \left[ S \cdot \left( \lim_{t_+ \rightarrow 0} \text{Tr} [-2i\alpha \cdot \gamma_5 f_+(t_+ D_+^2)] + \lim_{t_- \rightarrow 0} \text{Tr} [-2i\alpha \cdot \gamma_5 f_-(t_- D_-^2)] \right) \right] \quad (3.35)$$

We already calculated  $\lim_{t_\pm \rightarrow 0} \text{Tr} [-2i\alpha \cdot \gamma_5 f_\pm(t_\pm D_\pm^2)]$  in the previous paragraph. By equation 3.21 we conclude

$$\log J = \frac{i}{16} e^{\mu\nu\rho\sigma} \int d^4 x \alpha(x) \text{tr}_g [S \cdot (F_{\mu\nu}^+ F_{\rho\sigma}^+ + F_{\mu\nu}^- F_{\rho\sigma}^-)] \quad (3.36)$$

where  $F_{\mu\nu}^{\pm}$  is the field strength with respect to  $A_{\mu}^{\pm}$ . Hence equation 3.29 equals

$$\begin{aligned} \tilde{\mathcal{Z}} = & \int \mathcal{D}\bar{\psi}_+ \mathcal{D}\psi_+ \mathcal{D}\bar{\psi}_- \mathcal{D}\psi_- \exp(-S[\psi_+, \psi_-]) \times \\ & \times \left( - \int_{\mathbb{R}^4} d^4x \alpha(x) [\partial_{\mu}(\bar{\psi}\gamma^{\mu}\gamma_5\psi) - \bar{\psi}\gamma^{\mu}\gamma_5(iV_{\mu}^b + iA_{\mu}^b\gamma_5)[S, \lambda_b]\psi - 2im\bar{\psi}\gamma_5\psi] \right) \times \\ & \times \left( + \int_{\mathbb{R}^4} d^4x \alpha(x) \text{tr}_{\mathfrak{g}} [S \cdot (F_{\mu\nu}^+ F_{\rho\sigma}^+ + F_{\mu\nu}^- F_{\rho\sigma}^-)] \right) \end{aligned} \quad (3.37)$$

Because  $\mathcal{Z}$  is conserved under the chiral transformation we get a conservation law. However it does not coincide with the classical conservation equation. This shows that the Abelian chiral current is anomalous. We call this anomaly the **covariant anomaly**.

### 3.3 The Bardeen anomaly

In Fujikawas method, we have to regulate in order to get a well-defined Jacobian. However, the result depends on the method of regularization. In this paragraph we consider another regulator for the chiral current. Again the result is anomalous, but this time it coincides with the one-loop calculations.

We consider the chiral symmetry given in equation 3.25:

$$\psi \mapsto e^{i\alpha(x)S\gamma_5}\psi \quad \bar{\psi} \mapsto \bar{\psi}e^{i\alpha(x)S\gamma_5} \quad (3.25)$$

The generating functional  $\mathcal{Z}$  transforms under this transformation and the Jacobian of the path integral equals

$$J_{\psi} = J_{\bar{\psi}} = \det^{-1} \exp(i\alpha\gamma_5) = \exp(-i \text{Tr}[\alpha \cdot \gamma_5]) \quad (3.38)$$

We need to regulate the Jacobian in order to get a well-defined result. For the covariant anomaly we used  $D_{\pm}$  as the regulator. We now try to regulate the Jacobian with the differential operator  $\not{D} - iV - iA\gamma_5$ . This operator is not Hermitian. Even more, it doesn't commute with its adjoint. Hence it is not suitable to use in functional calculus. To solve this, we take the analytic continuation of  $A_{\mu}$  and we transform  $A_{\mu}$  to  $iA_{\mu}$ . We regulate the Jacobian with the differential operator  $D = \not{D} - iV + A\gamma_5$ . At the end of our calculation we undo the transformation. Note that  $\text{Tr}[\alpha\gamma_5 \exp(-tD^2)]$  diverges in the limit  $t \rightarrow 0$ . This is due to the factor  $\gamma_5$  in the differential operator  $D$ . There are two approaches to solve this problem:

1. We can renormalize the theory such that the divergent terms are canceled.<sup>6</sup>
2. According to Hu et al. [1984], the  $t$ -divergent terms cancels if we choose a different regulator for  $d\bar{\psi}$  and  $d\psi$ : They regulate the Jacobian for  $d\psi$  with  $e^{-t D^2}$  but they use  $e^{-t \bar{D}^2}$  (with  $\bar{D}_\mu := \partial_\mu - iV_\mu - A_\mu$ ) for  $d\bar{\psi}$ .

We use the second method, but to prevent extra terms from the Baker-Campbell-Hausdorff formula we regulate both  $d\psi$  and  $d\bar{\psi}$  with  $\frac{1}{2}(e^{-t D^2} + e^{-t \bar{D}^2})$ . The regulated Jacobian equals

$$J = \lim_{t \rightarrow 0} \exp \left[ -i \text{Tr}(\gamma_5 \alpha (e^{-t D^2} + e^{-t \bar{D}^2})) \right] \quad (3.39)$$

where  $\text{Tr}$  is the trace over spinor fields and over all spin indices. We expand this trace using the plane wave basis  $e^{ik \cdot x}$ . This yields an expression for the trace over space-time, but we still need a trace over the spin-indices. If we denote the trace over the spin indices as  $\text{tr}$ , we get

$$\log J = -i \lim_{t \rightarrow 0} \int d^4 x \alpha(x) \cdot \int \frac{d^4 k}{(2\pi)^4} \text{tr} \left( \gamma_5 e^{-ik \cdot x} (e^{-t D^2} + e^{-t \bar{D}^2}) e^{ik \cdot x} \right) \quad (3.40)$$

Using the Leibniz rule  $[D, e^{ik \cdot x}] = i \not{k} e^{ik \cdot x}$ , we pull  $e^{ik \cdot x}$  to the left

$$\log J = -i \lim_{t \rightarrow 0} \int d^4 x \alpha(x) \cdot \int \frac{d^4 k}{(2\pi)^4} \text{tr} \left( \gamma_5 \left( e^{-t (D+i\not{k})^2} + e^{-t (\bar{D}+i\not{k})^2} \right) \right) \quad (3.41)$$

and as before we rescale  $k_\mu$  with  $t^{-1/2} k_\mu$ :

$$\log J = -i \lim_{t \rightarrow 0} \int d^4 x \alpha(x) \cdot \int \frac{d^4 k}{(2\pi)^4} t^{-2} \text{tr} \left( \gamma_5 \left( e^{-(t^{1/2} D + i\not{k})^2} + e^{-(t^{1/2} \bar{D} + i\not{k})^2} \right) \right) \quad (3.42)$$

We consider the Taylor series in  $t^{1/2}$  at  $t = 0$ . In this approximation the integrand of equation 3.42 equals

$$\begin{aligned} e^{-k^2} \text{tr} [ & \gamma_5 (2t^{-2} + \\ & - it^{-3/2} (\{D, \not{k}\} + \{\bar{D}, \not{k}\}) \\ & - t^{-1} \left( D^2 + \bar{D}^2 + \frac{1}{2} (\{D, \not{k}\})^2 + \frac{1}{2} (\{\bar{D}, \not{k}\})^2 \right) \\ & + \frac{i}{2} t^{-1/2} (D^2 \{D, \not{k}\} + \bar{D}^2 \{\bar{D}, \not{k}\}) ) ] + \dots \end{aligned} \quad (3.43)$$

---

<sup>6</sup>For the mathematicians: Renormalization is the process of modifying the action with  $t$  dependent terms such that the final result is finite.

The order  $t^{-2}$  vanishes because it is proportional to  $\text{tr}(\gamma_5) = 0$ . We notice that the orders  $t^{-3/2}$  and  $t^{-1/2}$  are odd in  $k$ . Hence these orders also vanishes when we integrate over  $k$ . To calculate the  $t^{-1}$  order let  $D_0 = \not{\partial} + iV$  and notice that

$$D^2 + \bar{D}^2 + \frac{1}{2} (\{D, k\})^2 + \frac{1}{2} (\{\bar{D}, k\})^2 = 2D_0^2 + 2A^2 + (\{D_0, k\})^2 + ([k, A])^2 \quad (3.44)$$

All notions of  $\gamma_5$  disappears. We apply the trace identities by counting the Clifford degree. The terms  $D_0^2$  and  $A^2$  vanishes in the trace. The other terms in equation 3.44 are proportional to  $\epsilon^{\mu\nu\rho\sigma} k_\mu k_\nu$ . However, the integral  $\int d^4 k e^{-k^2} k_\mu k_\nu$  equals  $\frac{\pi^2}{2} \eta_{\mu\nu}$  and so the  $t^{-1}$  order vanishes completely. We conclude that the Jacobian does not diverge when  $t \rightarrow 0$ .

The calculation of the  $t$ -independent part of equation 3.42 is tedious and hence we use computer algebra to solve this problem. We calculate the Jacobian using FORM(Vermaseren [2000]) and the source code is given in appendix 11.1. The basic structure of the calculation is as follows:

1. We expand the exponent in equation 3.42 using the series  $\exp(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!}$ . We only consider the taylor series with five orders. All higher order terms vanishes because they are a multiple of  $t^{1/2}$ .
2. We work out the trace using the gamma matrix identities

$$\text{tr}(\gamma_5) = 0 \quad \text{tr}(\gamma_5 \gamma^\mu \gamma^\nu) = 0 \quad \text{tr}(\gamma_5 \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma) = 4\epsilon^{\mu\nu\rho\sigma} \quad (3.45)$$

3. We integrate over  $k$  using the identities:

$$\begin{aligned} \int d^4 k e^{-k^2} &= \pi^2 & \int d^4 k e^{-k^2} k_\mu k_\nu &= \frac{\pi^2}{2} \eta_{\mu\nu} \\ \int d^4 k e^{-k^2} k_\mu k_\nu k_\rho k_\sigma &= \frac{\pi^2}{4} (\eta_{\mu\nu} \eta_{\rho\sigma} + \eta_{\mu\rho} \eta_{\nu\sigma} + \eta_{\mu\sigma} \eta_{\nu\rho}) \end{aligned} \quad (3.46)$$

4. We simplify the result and we recall that we transformed  $iA_\mu$  to  $A_\mu$ .

After the calculation we conclude that the Jacobian equals

$$\begin{aligned} J = \exp \left[ \int \frac{d^4 k}{16\pi^2} \alpha(x) \epsilon^{\mu\nu\rho\sigma} \text{tr} \left( F_{\mu\nu} F_{\rho\sigma} + \frac{1}{3} G_{\mu\nu} G_{\rho\sigma} + \frac{32}{3} A_\mu A_\nu A_\rho A_\sigma - \right. \right. \\ \left. \left. - \frac{8}{3} (A_\mu A_\nu F_{\rho\sigma} + A_\mu F_{\nu\rho} A_\sigma + F_{\mu\nu} A_\rho A_\sigma) \right) \right] \end{aligned} \quad (3.47)$$

where

$$F_{\mu\nu} = \partial_\mu V_\nu - \partial_\nu V_\mu + [V_\mu, V_\nu] + [A_\mu, A_\nu] \quad (3.48)$$

$$G_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [V_\mu, A_\nu] + [A_\mu, V_\nu]. \quad (3.49)$$

This result coincide with the results in Bertlmann [1996] and van Nieuwenhuizen [1989] and was first found by Bardeen [1969]. Therefore,  $F_{\mu\nu}$  and  $G_{\mu\nu}$  are called Bardeen curvatures.

We compare the difference between the Bardeen anomaly and the covariant anomaly. We notice that the Bardeen anomaly is not covariant under gauge transformation. From this we immediately see that the Covariant anomaly and the Bardeen anomaly must differ. There are two possible factors where the difference can come from:

1. We used different regulators for the different anomalies.
2. We changed  $A_\mu$  to  $iA_\mu$  when we calculated the Bardeen anomaly.

When we set  $A_\mu$  to zero, the Dirac operators  $D$ ,  $\bar{D}$  and  $D^\pm$  become equal and hence the regulators are equal. The act of making making  $A_\mu$  imaginary is irrelevant, when  $A_\mu = 0$ . Hence the results must be equal when  $A_\mu = 0$ . This is indeed true, because the Bardeen anomaly simplifies to

$$\epsilon^{\mu\nu\rho\sigma} (\partial_\mu V_\nu - \partial_\nu V_\mu + [V_\mu, V_\nu]) (\partial_\rho V_\sigma - \partial_\sigma V_\rho + [V_\rho, V_\sigma]). \quad (3.50)$$

This is equal to the Covariant anomaly.

Earlier we calculated the chiral anomaly using Feynman diagrams. Recall that the rules for Feynman diagrams comes from the perturbation theory of the path integral formalism. In table 5 we compare our earlier calculated results with the results found by Bardeen. For example, the term  $\epsilon^{\mu\nu\rho\sigma} A_\mu A_\nu A_\rho A_\sigma$  corresponds to an interaction with four external  $A$ -fields. Up to first loop approximation, this is calculated in a pentagon diagram<sup>7</sup>. In this diagram we denote

$$F_{\mu\nu}^{lin} = \partial_\mu V_\nu - \partial_\nu V_\mu \quad (3.51)$$

$$G_{\mu\nu}^{lin} = \partial_\mu A_\nu - \partial_\nu A_\mu \quad (3.52)$$

See that the factor  $\frac{1}{3}$  difference in the AVV and AAA diagram corresponds to the factor  $\frac{1}{3}$  between  $F_{\mu\nu}^{lin} F_{\rho\sigma}^{lin}$  and  $G_{\mu\nu}^{lin} G_{\rho\sigma}^{lin}$ . We see that the one loop approximation fully

---

<sup>7</sup>Although a pentagon diagram has 5 external fields, one is contracted to calculate the anomaly. Hence, it correspond to  $\epsilon^{\mu\nu\rho\sigma} A_\mu A_\nu A_\rho A_\sigma$

captures the anomaly. This is first found by Adler and Bardeen [1969] and is called the Adler-Bardeen theorem:

**Theorem 3.1** (Adler-Bardeen). *The chiral anomaly can be fully determined in the one loop approximation.*

Bardeen anomaly	Factor	Feynman diagram	Amplitude
$F_{\mu\nu}^{lin} F_{\rho\sigma}^{lin}$	1	AVV	$\epsilon^{\mu\nu\rho\sigma} q_\rho p_\sigma$
$G_{\mu\nu}^{lin} G_{\rho\sigma}^{lin}$	$\frac{1}{3}$	AAA	$\frac{1}{3} \epsilon^{\mu\nu\rho\sigma} q_\rho p_\sigma$
$F_{\mu\nu}^{lin} V_\rho V_\sigma + V_\mu V_\nu F_{\rho\sigma}^{lin}$	2	AVVV	$\epsilon^{\mu\nu\rho\sigma} p_\sigma^{(i)}$
$G_{\mu\nu}^{lin} V_\rho A_\sigma + G_{\mu\nu}^{lin} A_\rho V_\sigma +$ $V_\mu A_\nu G_{\rho\sigma}^{lin} + A_\mu V_\nu G_{\rho\sigma}^{lin} -$ $-F_{\mu\nu}^{lin} A_\rho A_\sigma - 4A_\mu F_{\nu\rho}^{lin} A_\sigma - A_\mu A_\nu F_{\rho\sigma}^{lin}$	$\frac{2}{3}$	AAAV	-
$V_\mu V_\nu V_\rho V_\sigma$	4	AVVVV	$\epsilon^{\mu\nu\rho\sigma}$
$V_\mu A_\nu V_\rho A_\sigma - V_\mu V_\nu A_\rho A_\sigma +$ $A_\mu V_\nu A_\rho V_\sigma - A_\mu A_\nu V_\rho V_\sigma +$ $V_\mu A_\nu A_\rho V_\sigma - 4A_\mu V_\nu V_\rho A_\sigma$	$\frac{4}{3}$	AAAVV	-
$A_\mu A_\nu A_\rho A_\sigma$	$-\frac{4}{3}$	AAAAA	-

Table 5: *The Bardeen anomaly has an one loop approximation using Feynman diagrams. In this table every term in the Bardeen anomaly is related to a triangle, box or pentagon diagram. The prefactor of each term in the Bardeen anomaly are given relative to  $F_{\mu\nu}^{lin} F_{\rho\sigma}^{lin}$ . Also, the results from chapter 2 are again stated.*

### 3.4 The consistent anomaly

When we calculated the covariant anomaly, we simplified the calculation by considering the  $\gamma_5$  eigenbasis. We concluded that the action given in equation 3.28 describes two non-interacting particles. We now ask what happens to the chiral anomaly if we only

consider massless right-handed particles  $\psi_+$ . That is, what is the chiral anomaly if we consider the action

$$S[\psi_+] = \int_{\mathbb{R}^4} -\bar{\psi}_+(\not{\partial} - iA^+)\psi_+ \, dx \quad (3.53)$$

However, this does not define a well-defined quantum theory. Using  $\psi_+ = P_+\psi = \frac{1}{2}(1 + \gamma_5)$  and  $D_+ = \not{\partial} - iA^+$  we rewrite this action as

$$S[\psi] = \int_{\mathbb{R}^4} -\bar{\psi}(D_+)P_+\psi \, dx \quad (3.54)$$

The operator  $D_+P_+$  maps right-handed chirality spinors into left-handed chirality spinors. Therefore, this operator has no eigenvalues and the generating functional, which is the formal determinant of  $iD_+P_+$ , cannot be defined. Alvarez-Gaumé and Ginsparg [1984] [1985] noticed that if we consider the action

$$S[\psi] = \int_{\mathbb{R}^4} -\bar{\psi}(D_+P_+ + \not{\partial}P_-)\psi \, dx \quad (3.55)$$

we get a well-defined quantum theory. Also the gauge couples only to the positive chirality spinors and the non-zero eigenmodes are all right-handed. Hence the gauge theories coincide. We assume that  $\psi_+$  is invariant under the symmetry  $\psi_+ \mapsto e^{i\alpha(x)\gamma_5}\psi_+$ . Here we assume that  $\alpha$  is an infinitely small Lie algebra valued smooth map. The action transforms into

$$\tilde{S}[\psi, A_\mu^+] = S[\psi, A_\mu^+] + \int_{\mathbb{R}^4} -i\bar{\psi}\gamma^\mu(\tilde{D}_\mu^+\alpha(x))\psi_+ \, dx + \mathcal{O}(\alpha^2) \quad (3.56)$$

where  $\tilde{D}_\mu^+ = \delta_\mu - i[A_\mu^+, \cdot]$ . The generating functional  $\mathcal{Z}[A_\mu^+]$  transforms into

$$\tilde{\mathcal{Z}}[A_\mu^+] = \int \mathcal{D}\bar{\psi}\mathcal{D}\psi J \cdot \exp\left(-\tilde{S}[\psi, A_\mu^+]\right) \quad (3.57)$$

where  $J$  is the Jacobian of the path integral. We can use Fujikawas method to calculate the Jacobian. However, this is a special case (Andrianov and Bonora [1984]) of Bardeens anomaly if we substitute

$$V_\mu \rightarrow \frac{1}{2}A_\mu^+ \quad A_\mu \rightarrow \frac{1}{2}A_\mu^+ \quad (3.58)$$

The Jacobian  $J$  equals  $\int d^4x \, \text{tr}(\alpha(x) \cdot G[A^+])$  where  $G[A^+]$  is

$$G[A^+] = \frac{1}{32\pi^2}\epsilon^{\mu\nu\rho\sigma} \left( \frac{1}{3}A_\mu^+A_\nu^+ (\partial_\rho A_\sigma^+) - \frac{1}{3}A_\mu^+(\partial_\nu A_\rho^+)A_\sigma^+ \right. \\ \left. + \frac{1}{3}(\partial_\mu A_\nu^+)A_\rho^+A_\sigma^+ + \frac{2}{3}(\partial_\mu A_\nu^+)(\partial_\rho A_\sigma^+) \right). \quad (3.59)$$

We simplify  $G[A^+]$  into

$$G[A^+] = \frac{1}{24\pi^2} \epsilon^{\mu\nu\rho\sigma} \partial_\mu \left( A_\nu^+ (\partial_\rho A_\sigma^+) + \frac{1}{2} A_\nu^+ A_\rho^+ A_\sigma^+ \right) \quad (3.60)$$

This result is called the **consistent anomaly**. It is called this way, because  $G[A^+]$  satisfies the Wess-Zumino consistency condition [1971].

We now derive this condition. By assumption the generating functional is invariant under the chiral symmetry. So from Equations 3.56 and 3.57 it follows that

$$\mathcal{Z}[A_\mu^+ + \tilde{D}_\mu^+ \alpha(x)] = \tilde{\mathcal{Z}}[A_\mu^+ + \tilde{D}_\mu^+ \alpha(x)] \quad (3.61)$$

$$= \int \mathcal{D}\bar{\psi} \mathcal{D}\psi \exp \left( -S[\psi, A_\mu^+] + \int \alpha(x) G[A^+] \, d^4 x \right) \quad (3.62)$$

and we conclude

$$\lim_{t \rightarrow 0} \frac{1}{t} \left( \log \mathcal{Z}[A_\mu^+ + t \tilde{D}_\mu^+ \alpha(x)] - \log \mathcal{Z}[A_\mu^+] \right) = \int \alpha(x) G[A^+](x) \, d^4 x. \quad (3.63)$$

In terms of the variational derivative we yield the **anomalous Ward identity**  $\frac{\delta}{\delta \alpha} \log \mathcal{Z} = G[A^+]$ . The Ward identity can also be written in terms of differential forms<sup>8</sup>. For this we view  $\mathcal{Z}$  as a smooth real-valued map on the space of all fields  $A_\mu^+$ . That is,  $\mathcal{Z}$  is a 0-form on an infinite dimensional manifold. The left hand side of equation 3.63 can be interpreted as the Lie derivative of  $\log(\mathcal{Z})$  in the direction  $\tilde{D}_\mu^+ \alpha$ . The right hand side is the  $L^2$  inner product on the space of Lie algebra valued smooth maps. Hence, the anomalous Ward identity equals

$$\mathcal{L}_{\tilde{D}^+ \alpha} \log \mathcal{Z} = \langle \alpha, G[A^+] \rangle_{L^2}. \quad (3.64)$$

Let  $\tilde{G}$  be the exterior derivative  $d \log(\mathcal{Z})$ . It is the unique 1-form such that  $\tilde{G}(\delta A_\mu) = \mathcal{L}_{\delta A_\mu} \log(\mathcal{Z})$ . By equation 3.64 we conclude  $\tilde{G}(\tilde{D}^+ \alpha) = \langle \alpha, G[A^+] \rangle_{L^2}$ . Using Cartan's magic formula we calculate the exterior derivative of  $\tilde{G}$ .

$$d \tilde{G}(\tilde{D}^+ \alpha, \tilde{D}^+ \beta) = \iota_{\tilde{D}^+ \beta} \circ \iota_{\tilde{D}^+ \alpha} \circ d \tilde{G} \quad (3.65)$$

$$= -\iota_{\tilde{D}^+ \beta} \circ d \circ \iota_{\tilde{D}^+ \alpha} \tilde{G} + \iota_{\tilde{D}^+ \beta} \circ \mathcal{L}_{\tilde{D}^+ \alpha} \tilde{G} \quad (3.66)$$

$$= d \circ \iota_{\tilde{D}^+ \beta} \circ \iota_{\tilde{D}^+ \alpha} \tilde{G} - \mathcal{L}_{\tilde{D}^+ \beta} \circ \iota_{\tilde{D}^+ \alpha} \tilde{G} + \iota_{\tilde{D}^+ \beta} \circ \mathcal{L}_{\tilde{D}^+ \alpha} \tilde{G} \quad (3.67)$$

<sup>8</sup>See the appendix for a short introduction.

We notice that  $\iota_{\tilde{D}+\beta} \circ \iota_{\tilde{D}+\alpha} \tilde{G}$  is a  $(-1)$ -form. These forms does not exists and hence  $d \circ \iota_{\tilde{D}+\beta} \circ \iota_{\tilde{D}+\alpha} \tilde{G}$  is zero. We also recall that the commutator between a Lie derivative and the interior product is the interior product of a Lie bracket. That is,  $[L_x, \iota_Y] = \iota_{[X,Y]}$  and hence

$$d \tilde{G}(\tilde{D}^+\alpha, \tilde{D}^+\beta) = - [\mathcal{L}_{\tilde{D}+\alpha}, \iota_{\tilde{D}+\beta}] \tilde{G} + \mathcal{L}_{\tilde{D}+\alpha} \circ \iota_{\tilde{D}+\beta} \tilde{G} - \mathcal{L}_{\tilde{D}+\beta} \circ \iota_{\tilde{D}+\alpha} \tilde{G} \quad (3.68)$$

$$= - \iota_{[\tilde{D}+\alpha, \tilde{D}+\beta]} \tilde{G} + \mathcal{L}_{\tilde{D}+\alpha} \circ \iota_{\tilde{D}+\beta} \tilde{G} - \mathcal{L}_{\tilde{D}+\beta} \circ \iota_{\tilde{D}+\alpha} \tilde{G}. \quad (3.69)$$

By the definition of the exterior derivative  $d^2 = 0$ , and so equation 3.69 equals zero. Using the Ward identity we simplify equation 3.69 into

$$\mathcal{L}_{\tilde{D}+\alpha} \langle \tilde{D}^+\beta, G[A^+] \rangle_{L_2} - \mathcal{L}_{\tilde{D}+\beta} \langle \tilde{D}^+\alpha, G[A^+] \rangle_{L_2} - \langle [\tilde{D}^+\alpha, \tilde{D}^+\beta], G[A^+] \rangle_{L_2} = 0 \quad (3.70)$$

This is the Wess-Zumino consistency condition.

### 3.5 Mathematical interpretation of the Fujikawa method

At last we give a mathematical interpretation to the Fujikawa method. In section 3.1 we calculated the chiral anomaly for a Dirac particle in quantum electrodynamics. The anomaly was due to the Jacobian of the path integral. By equation 3.21 the Jacobian equals

$$\log J = \frac{i}{16} e^{\mu\nu\rho\sigma} \int_{\mathbb{R}^4} dx \alpha(x) F_{\mu\nu} F_{\rho\sigma}. \quad (3.21)$$

For simplicity we assume that  $\alpha(x) = 1$ . We rewrite  $J$  in terms of differential forms<sup>9</sup>. Recall that  $F^{\mu\nu}$  are the components of the curvature tensor  $F$ . For quantum electrodynamics  $F$  is a complex valued 2-form. By definition the wedge product between  $F$  and itself equals  $(F \wedge F)^{0123} = \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma}$ . This shows that

$$\log J = \frac{i}{8} \int_{\mathbb{R}^4} F \wedge F. \quad (3.71)$$

By stokes theorem we conclude that the Jacobian is determined by the cohomology class of  $F \wedge F$ . In section ?? we will show that this cohomology class only depends on *topology* of the gauge bundle. At the same time, the regulated Jacobian equals

$$\log J = -2i \lim_{t \rightarrow 0} \text{Tr} [\gamma_5 f(t \cdot D^2)] \quad (3.72)$$

---

<sup>9</sup>See the appendix for a basic introduction.

In the orthonormal eigenbasis  $\phi_n$  of  $D$  where  $\lambda_n$  is the corresponding eigenvalue this trace is given by

$$\log J = -2i \lim_{t \rightarrow 0} \sum_n \langle \phi_n | \gamma_5 f(t \cdot D^2) | \phi_n \rangle \quad (3.73)$$

$$= -2i \lim_{t \rightarrow 0} \sum_n f(t \cdot \lambda_n^2) \langle \phi_n | \gamma_5 | \phi_n \rangle \quad (3.74)$$

Recall that  $\gamma_5$  anti-commutes with the Dirac operator  $D$  and hence  $\gamma_5 \phi_n$  is an eigenvector of  $D$  with eigenvalue  $-\lambda_n$ . By the orthonormality of the eigenspaces we conclude that  $\langle \phi_n | \gamma_5 | \phi_n \rangle = 0$  if  $\lambda_n \neq 0$ . If  $n^\pm$  denotes the number of independent left- resp. right-handed zero modes of  $D$  then,

$$\log J = -2i \sum_{\text{left-handed zero modes}} \langle \phi_n | \gamma_5 | \phi_n \rangle - 2i \sum_{\text{right-handed zero modes}} \langle \phi_n | \gamma_5 | \phi_n \rangle \quad (3.75)$$

$$= 2i \sum_{\text{left-handed zero modes}} \langle \phi_n | \phi_n \rangle - 2i \sum_{\text{right-handed zero modes}} \langle \phi_n | \phi_n \rangle \quad (3.76)$$

$$= -2i(n^+ - n^-) \quad (3.77)$$

This result is found by McKean and Singer [1967]. In mathematics the quantity  $n^+ - n^-$  is called the *index of  $D$* . Using the Fujikawa method we see that the index is determined by the topology of a vector bundle. This relation is first found by Atiyah and Singer [1968] and is called the Atiyah-Singer index theorem.

In the next chapters we prove this theorem for Dirac operators on compact spaces<sup>10</sup> by using Fujikawa's method. In chapter 4 we define the operator  $\exp(-tD^2)$ . We will not define it using the eigenvalues of  $D$ , but as the unique operator that satisfies  $(\frac{\partial}{\partial t} + D^2)e^{-tD^2} = 0$ . In chapter 5 we show that the trace over  $\gamma_5 e^{-tD^2}$  is finite and does not depend on the choice of basis. Hence, we show that  $e^{-tD^2}$  is a well-chosen regulator for  $J$ .

In the next step Fujikawa considered the trace in the plane wave basis  $e^{ik \cdot x}$  and showed that the index of  $D$  equals the integral over a trace. In chapter 4 we show that  $e^{-tD^2}$  has a kernel. That is, we show that there exists an operator  $k_t$  such that  $e^{-tD^2}$  is the integral over this operator. Later in chapter 5 we show that the trace over  $\gamma_5 e^{-tD^2}$

<sup>10</sup>That is, we assume that the space-time is of finite size and we assume that it does not contain any singularities.

reduces to the integral  $\int \text{tr}(\gamma_5 k_t)$ .

In the last step Fujikawa calculated a Taylor series. In chapter 7 we formalize this in the theory of graded and filtered algebras. We show that the number of gamma matrices induce a grading. Also, Taylor series in  $k_\mu$  forms a grading. However the grading Fujikawa used is a combination of both and is due to Getzler [1983]. Till now we explicitly calculated each term of the Taylor series. This is tedious and is not useful when we consider the general case. Hence we investigate how the Getzler grading behaves under the differential equation  $\frac{\partial}{\partial t} + D^2$ . This yields another differential equation that we can explicitly solve.

## 4 Smoothing operators and Heat kernels

In chapter 3 we saw how the Abelian anomaly was related to the index of a Dirac operator. In the following chapters we analyze this method and work out the technical details. In this chapter we question whether a Dirac operator can be exponentiated. This is indeed possible and we also show that we can write it as an integral.

We follow the approach given by Berline et al. [2004]. We construct the exponential by solving a differential equation. This is still a formal solution, because we need to show that this formal solution converges. We prove this in two steps. First we modify the formal solution into an approximate solution that does converge. Secondly we increase the accuracy of the approximate solution. So the basic steps in this chapter are

$$\text{Formal solution} \implies \text{Approximate solution} \implies \text{Existence.}$$

### 4.1 Definitions

We want to study  $e^{-tD^2}$ . From the Lichnerowicz formula, we know that

$$D^2 = \nabla^* \nabla + F^S + \frac{1}{4} \kappa \quad (4.1)$$

where  $F^S + \frac{1}{4} \kappa$  is a section of  $\text{End}(S)$ . In this chapter we don't need the Clifford structure and hence we study generalized Laplacians:

**Definition 4.1.** *Let  $(M, g)$  be a Riemannian manifold and let  $E$  be a vector bundle over  $M$  with a positive definite inner product. A self-adjoint map  $H: \Gamma(E) \rightarrow \Gamma(E)$  is a **generalized Laplacian** if there exists a compatible connection  $\nabla$  such that  $H - \nabla^* \nabla$  is a section of  $\text{End}(E)$ .*

We want to write  $e^{-tD^2}$  as an integral. Although vector-valued integration is well-defined, vector bundle valued integration is not. To solve this, we construct an operator which maps all fibers into a single one. Then vector bundle integration reduces to vector space integration which is well defined. Such an operator is called a kernel. Informally, for two vector bundles  $E^1$  and  $E^2$  and  $x, y \in M$  a kernel is a "smooth" linear map  $p(x, y)$  from  $E_y^1$  to  $E_x^2$ . To define smoothness we recall that  $\text{Hom}(E_y^1, E_x^2)$

is isomorphic to  $(E_y^1)^* \otimes E_x^2$ . Using the canonical projections  $\text{pr}_1$  and  $\text{pr}_2$  from  $M \times M$  to  $M$  we can write this as

$$\text{pr}_1^* E^2 \otimes \text{pr}_2^* E^1_{(x,y)} \simeq (E^2)_x \otimes (E_y^1)^* \simeq \text{Hom}(E_y^1, E_x^2). \quad (4.2)$$

To simplify notation we write the vector bundle  $\text{pr}_1^* E^2 \otimes \text{pr}_2^* E^1 \rightarrow M \times M$  as  $E^1 \boxtimes E^2 \rightarrow M \times M$ . We see  $p$  is a section of  $E^1 \boxtimes E^2$ . Smoothness of a kernel follows from the fact that sections are smooth.

**Definition 4.2.** Let  $(M, g)$  be a compact Riemannian manifold and let  $E^1$  and  $E^2$  be vector bundles over  $M$  with positive definite inner products. A **kernel** is a section  $p \in \Gamma(E^1 \boxtimes E^2)$  (which is a linear map  $p(x, y): E_y^1 \rightarrow E_x^2$ ). A linear operator  $P: \Gamma(M, E_1) \rightarrow \Gamma(M, E_2)$  is a **smoothing operator** if there exists a kernel  $p \in \Gamma(E^1 \boxtimes E^2)$  such that

$$(Ps)(x) = \int_{y \in M} p(x, y) s(y) \text{Vol}(g)$$

There are two methods to define  $e^{-tD^2}$ . One method is to consider the eigenvalues  $\lambda_i$  of  $D^2$  and construct an eigenbasis  $\{v_i\}$  of  $D^2$ . Then we define  $e^{-tD^2}$  by  $e^{-tD^2} v_i = e^{-t\lambda_i} v_i$ . For this method we need to estimate the eigenvalues of  $D^2$  before we can show that the exponent converges. Also, if we want to show that this is a smoothing operator, then we need to prove this separately. We follow a more direct approach: We consider all smoothing operators that satisfy  $\frac{\partial}{\partial t} e^{-tD^2} = -D^2 e^{-tD^2}$ . We then show that there is a unique operator satisfying this equation which has therefore the properties we would formally expect from  $e^{-tD^2}$ .

**Definition 4.3.** Let  $(M, g)$  be a compact Riemannian manifold,  $E$  be a vector bundle over  $M$  with a positive definite inner product and  $H$  be a generalized Laplacian. A **heat kernel** for  $H$  is a section  $k^H$  of the vector bundle  $E \boxtimes E \rightarrow \mathbb{R}_+ \times M \times M$  which has the following properties:

1. The kernel  $k^H$  is at least once continuous differentiable in the first component
2. The kernel  $k^H$  is at least twice continuous differentiable in the second component
3. The kernel  $k$  satisfies the heat equation:

$$\left( \frac{\partial}{\partial t} + H_x \right) k^H(t, x, y) = 0 \quad \forall x, y \in M, t \in \mathbb{R}_+ \quad (4.3)$$

4. The kernel  $k^H$  satisfies the boundary condition. That is, in the supremum norm

$$\lim_{t \rightarrow 0} \int_{y \in M} k^H(t, x, y) s(y) \text{Vol}(g) = s(x) \quad (4.4)$$

for all  $x \in M$  and  $s \in \Gamma(E)$ .

We denote  $k^H(t, \cdot, \cdot)$  as  $k_t^H(\cdot, \cdot)$ .

## 4.2 Examples of heat kernels

In this paragraph we give some useful examples of heat kernels for Laplacians on the real line. Although  $\mathbb{R}$  is not compact we can define a heat kernel if we restrict the boundary condition to only square integrable sections of  $\mathbb{R}$ . Note that on the real line  $p_t$  reduces to a smooth map  $\mathbb{R} \times \mathbb{R} \times \mathbb{R}_+ \rightarrow \mathbb{R}$ .

The simplest generalized Laplacian is the standard Laplacian  $H = -\frac{\partial}{\partial x^2}$ . The heat equation  $\frac{\partial}{\partial t} - \frac{\partial}{\partial x^2} = 0$  suggests that the heat kernel is a Gaussian function. By trial and error we can find that the map  $k_t^H(x, y) = \frac{1}{\sqrt{4\pi t}} e^{-(x-y)^2/4t}$  satisfies

$$\left( \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right) k_t^H(x, y) = 0. \quad (4.5)$$

This suggests that  $k_t^H$  is a heat kernel. This is indeed true.

**Lemma 4.4** (Berline et al. [2004], Lemma 2.12). *The map  $k_t^H(x, y) = \frac{1}{\sqrt{4\pi t}} e^{-(x-y)^2/4t}$  is a heat kernel for the generalized Laplacian  $H = -\frac{\partial^2}{\partial x^2}$  on  $\mathbb{R}$ .*

*Proof.* We are left to show that  $\int_{\mathbb{R}} k_t^H(x, y) s(y) dy = s(x)$  for all  $s \in L^2(\mathbb{R})$ . Recall that compactly supported smooth maps are dense in  $L^2(\mathbb{R})$  and on this compact support we can approximate these maps with polynomials. Hence it is sufficient to show that this is true if  $s(x) = x^k$ . Consider the map

$$A(s, t) = \frac{1}{\sqrt{4\pi t}} \int_{\mathbb{R}} e^{-(x-y)^2/4t + sy} dy \quad (4.6)$$

This map converges, because if we substitute  $y$  with  $y = \sqrt{4t}\tilde{y} + 2st + x$  we get

$$A(s, t) = \frac{1}{\sqrt{\pi}} \int_{\mathbb{R}} e^{-(x - (\sqrt{4t}\tilde{y} + 2st + x))^2 / 4t + s(\sqrt{4t}\tilde{y} + 2st + x)} d\tilde{y} \quad (4.7)$$

$$= \frac{1}{\sqrt{\pi}} \int_{\mathbb{R}} e^{-\tilde{y}^2 + s^2 t + sx} d\tilde{y} \quad (4.8)$$

$$= \exp(s^2 t + sx) \quad (4.9)$$

We calculate the Taylor series of  $A(s, t)$  in  $s$ . Comparing equation 4.6 and 4.9 we see that

$$\sum_{k=0}^{\infty} \frac{s^k}{k!} \left( \frac{1}{\sqrt{4\pi t}} \int_{\mathbb{R}} e^{-(x-y)^2 / 4t} y^k d y \right) = \sum_{k=0}^{\infty} \frac{s^k}{k!} \cdot (st + x)^k \quad (4.10)$$

In the limit  $t \rightarrow 0$  this simplifies to

$$\sum_{k=0}^{\infty} \frac{s^k}{k!} \left( \lim_{t \rightarrow 0} \frac{1}{\sqrt{4\pi t}} \int_{\mathbb{R}} e^{-(x-y)^2 / 4t} y^k d y \right) = \sum_{k=0}^{\infty} \frac{s^k}{k!} \cdot x^k \quad (4.11)$$

This proves  $\int_{\mathbb{R}} k_t^H(x, y) s(y) d y = s(x)$  for all  $s \in L^2(\mathbb{R})$ .  $\square$

Another example is the quantum mechanical harmonic oscillator in one dimension . Up to constants the Schrödinger equation for this system is

$$\frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} + \omega^2 x^2 = 0 \quad (4.12)$$

where  $\omega \in \mathbb{R}$  is the angular frequency. Notice that this is the heat equation for the generalized Laplacian

$$H = -\frac{\partial^2}{\partial x^2} + \omega^2 x^2 \quad (4.13)$$

The heat kernel is found by Mehler [1866] and in the next lemma we give the formula.

**Lemma 4.5** (Mehler [1866]). *Let  $H$  be the generalized Laplacian on  $\mathbb{R}$  that is given by  $H = -\frac{\partial^2}{\partial x^2} + \omega^2 x^2$ . Then the heat kernel w.r.t.  $H$  exists and equals*

$$k_t^H(x, y) = \sqrt{\frac{\omega}{2\pi \sinh(2\omega t)}} \exp \left[ \frac{-\omega(x^2 + y^2) \coth(2\omega t) + 2\omega \operatorname{cosech}(2\omega t)xy}{2} \right]$$

*Proof.* To solve the heat equation, we use the ansatz:

$$k_t^H(x, y) = \exp \left( \frac{a(t)}{2}(x^2 + y^2) + b(t)xy + c(t) \right) \quad (4.14)$$

The heat equation we need to solve is then

$$0 = \frac{\dot{\alpha}}{2}x^2 + \dot{\beta}xy + \frac{\dot{\alpha}}{2}y^2 + \dot{\gamma} - (\alpha x + \beta y)^2 - \alpha + \omega^2 x^2 \quad (4.15)$$

$$= x^2 \left( \frac{\dot{\alpha}}{2} - \alpha^2 + \omega^2 \right) + y^2 \left( \frac{\dot{\alpha}}{2} - \beta^2 \right) + xy \left( \dot{\beta} - 2\alpha\beta \right) + (\dot{\gamma} - \alpha) \quad (4.16)$$

This equation must be valid for any  $x, y \in \mathbb{R}$ . Hence, we have the following system of differential equations

$$\frac{\dot{\alpha}(t)}{2} = \alpha^2(t) - \omega^2 = \beta^2(t) \quad \dot{\beta}(t) = 2\alpha(t)\beta(t) \quad \dot{\gamma}(t) = \alpha(t) \quad (4.17)$$

The differential equation for  $\alpha$  can be written as  $\int_{\mathbb{R}} \frac{1}{\alpha^2 - \omega^2} da = 2t + c_1$  where  $c_1 \in \mathbb{R}$  is an integration constant. Using change of variables this simplifies to  $\int_{\mathbb{R}} \frac{\dot{\alpha}(t)}{\alpha^2(t) - \omega^2} dt = 2t + c_1$ . The primitive is a cotangent hyperbolic<sup>11</sup> and  $\alpha$  equals

$$\alpha(t) = -\omega \coth(2\omega t + c_1) \quad (4.18)$$

By integration and differentiation we find expressions for  $\beta$  and  $\gamma$  and hence we have

$$\alpha(t) = -\omega \coth(2\omega t + c_1) \quad (4.19)$$

$$\beta(t) = +\omega \operatorname{cosech}(2\omega t + c_1) \quad (4.20)$$

$$\gamma(t) = -\frac{1}{2} \log(\sinh(2\omega t + c_1) * c_2) \quad (4.21)$$

Here  $c_1$  and  $c_2$  are integration constants and we find them by using the boundary condition. Already we notice that  $k_t^H$  is a bounded map and so it is a well defined operator on  $L^2(\mathbb{R})$ .

When we apply the smoothing operator on 1 and  $x^2$ , we conclude that  $c_1$  must vanish. We approximate the resulting kernel using a taylor series in  $t$  and we get that

$$k_t^H(x, y) = \frac{1}{\sqrt{4\pi t}} e^{-(x-y)^2/4t} \cdot \left( \sqrt{\frac{2\pi}{\omega c_2}} + \mathcal{O}(t) \right) \quad (4.22)$$

Using Lemma 4.4, we conclude that  $k_t^H$  satisfies the boundary condition if  $c_2 = \frac{\omega}{2\pi}$ .  $\square$

<sup>11</sup> Recall that  $\coth = \frac{\cosh}{\sinh}$  and  $\operatorname{cosech} = \frac{1}{\sinh}$

In our last example we generalize Mehlers kernel for  $\mathbb{R}^n$ . This extension will be important in the proof of Atiyahs index theorem.

**Lemma 4.6** (Roe [1998], Proposition 12.25). *Let  $R \in M_{n \times n}(\mathbb{R})$  be a skew-symmetric matrix and let  $F \in \mathbb{R}$ . Let  $H$  be the generalized Laplacian on  $\mathbb{R}^n$  given by  $H = -\sum_i \left( \frac{\partial}{\partial x_i} + \frac{1}{4} \sum_j R_{ij} x_j \right)^2 + F$ . Then the heat kernel  $k_t^H$  w.r.t.  $H$  is given by*

$$k_t^H(x, 0) = \frac{1}{(4\pi t)^{n/2}} \det^{1/2} \left( \frac{tR/2}{\sinh(tR/2)} \right) \exp \left[ -\frac{1}{4t} \left\langle \frac{tR}{2} \coth \left( \frac{tR}{2} \right) x, x \right\rangle - tF \right]$$

*Proof.* Assume that the heat kernel  $k_t^H$  can be written as

$$k_t^H(x, y) = w_t(x, y) e^{-tF} \quad (4.23)$$

The heat equation  $(\frac{\partial}{\partial t} + H_x)K_t^H(x, y) = 0$  simplifies to

$$e^{-tF} \left[ \frac{\partial w_t}{\partial t} - \sum_i \left( \frac{\partial}{\partial x_i} + \frac{1}{4} \sum_j R_{ij} x_j \right)^2 w_t \right] = 0 \quad (4.24)$$

and this gives us a differential equation for  $w_t$ . Next we consider the eigenvalue decomposition of  $R$ . Extend  $R$  to  $R_{\mathbb{C}}: \mathbb{C}^n \rightarrow \mathbb{C}^n$  and let  $\{z_i\}$  be the eigenbasis of  $R_{\mathbb{C}}$ . The eigenvalues  $\lambda_i$  w.r.t.  $z_i$  must be imaginary, because  $R$  is skew symmetric. So there exists  $\theta_i \in \mathbb{R}$  such that  $\lambda_i = i\theta_i$ . We split  $z_i$  in its real and complex components  $z_i = x_i + iy_i$ . We decompose the eigenvalue equation  $R_{\mathbb{C}}z_i = \lambda_i z_i$  into its real and complex components. It follows that

$$Rx_i = -\theta_i y_i \quad Ry_i = \theta_i x_i \quad (4.25)$$

Because  $\bar{z}_i$  is an eigenvector of  $R_{\mathbb{C}}$  and  $x_i$  and  $y_i$  forms a linear combination of  $z_i$  and  $\bar{z}_i$ , we get that the set  $\{x_i, y_i\}$  forms a basis of  $\mathbb{R}^n$ . Even more, this basis can be chosen orthonormal. This follows from the relations

$$\langle x_i, Rx_i \rangle = -\langle Rx_i, x_i \rangle = \theta_i \langle y_i, x_i \rangle = 0 \quad (4.26)$$

$$\langle x_i, Ry_i \rangle = -\langle Rx_i, y_i \rangle = \theta_i \|x_i\|^2 = \theta_i \|y_i\|^2 \quad (4.27)$$

In the orthonormal basis  $\{x_i, y_i\}$ , equation 4.24 becomes

$$\frac{\partial w_t}{\partial t} - \sum_i \left( \frac{\partial}{\partial x_i} + \frac{1}{4} \theta_i y_i \right)^2 w_t - \sum_i \left( \frac{\partial}{\partial y_i} - \frac{1}{4} \theta_i x_i \right)^2 w_t = 0 \quad (4.28)$$

We also assume that  $w_t = \prod_i u_i \cdot v_i$ , where  $u_i$  is a smooth map in  $x_i$  and  $t$  and  $v_i$  is a map in  $y_i$  and  $t$ . From this ansatz the heat equation reduces a differential equation for all  $i \leq n$ :

$$v_i \left( \frac{\partial u_i}{\partial t} - \frac{\partial^2 u_i}{\partial x_i^2} - \frac{\theta_i}{16} x_i^2 u_i \right) + u_i \left( \frac{\partial v_i}{\partial t} - \frac{\partial^2 v_i}{\partial y_i^2} - \frac{\theta_i}{16} y_i^2 v_i \right) = -\frac{\theta_i}{4} \left( x_i \frac{\partial}{\partial y_i} - y_i \frac{\partial}{\partial x_i} \right) u_i \cdot v_i \quad (4.29)$$

We further postulate that the left hand side and right hand side of equation 4.29 are independent differential equations and are both equal to zero. From the left hand side we notice that  $u_i$  and  $v_i$  are the heat kernels for the harmonic oscillator. That is

$$u_i(x_i, 0, t) = \frac{1}{\sqrt{4\pi t}} \sqrt{\frac{it\theta_i/2}{\sinh(it\theta_i/2)}} \exp \left[ -\frac{1}{8} i\theta_i x_i^2 \coth(it\theta_i/2) \right] \quad (4.30)$$

$$v_i(x_i, 0, t) = \frac{1}{\sqrt{4\pi t}} \sqrt{\frac{it\theta_i/2}{\sinh(it\theta_i/2)}} \exp \left[ -\frac{1}{8} i\theta_i y_i^2 \coth(it\theta_i/2) \right] \quad (4.31)$$

Notice that the product  $u_i \cdot v_i$  is rotation invariant. Therefore is  $\left( x_i \frac{\partial}{\partial y_i} - y_i \frac{\partial}{\partial x_i} \right) u_i \cdot v_i = 0$ . Thus the right hand side of equation 4.29 also vanishes. We finally have a solution of the heat equation which is given by

$$k_t^H(x, 0) = e^{-tF} \prod_i \frac{1}{4\pi t} \frac{it\theta/2}{\sinh(it\theta/2)} \exp \left[ -\frac{1}{8} i\theta (x_i^2 + y_i^2) \coth(it\theta/2) \right] \quad (4.32)$$

This equals the heat kernel proposed in the lemma. At last we need to check the boundary condition. Note that

$$\frac{tR/2}{\sinh(tR/2)} = \text{Id} + \mathcal{O}(t^2) \quad \frac{tR}{2} \coth \frac{rR}{2} = \text{Id} + \mathcal{O}(t^2), \quad (4.33)$$

Hence in the limit  $t \rightarrow 0$ ,  $\int_{\mathbb{R}^n} k_t^H(x, 0) s(x) = s(0)$  for all  $s \in L^2(\mathbb{R}^n)$ .  $\square$

### 4.3 Uniqueness of the heat kernel

**Proposition 4.7** (Berline et al. [2004], Proposition 2.17). *Let  $(M, g)$  be a compact Riemannian manifold and let  $E$  be a vector bundle on  $M$  with a positive definite*

inner product. Let  $H$  be a generalized Laplacian and suppose that  $H$  admits a heat kernel  $k_t^H \in \Gamma(E \boxtimes E)$ . Then  $k_t$  is unique and is self adjoint in the sense that

$$k_t^H(x, y)^* = k_t^H(y, x) \quad (4.34)$$

for all  $x, y \in M$  and  $t \in \mathbb{R}_+$ .

*Proof.* Let  $\tilde{k}_t^H \in \Gamma(E \boxtimes E)$  be another heat kernel of  $H$  and let  $K_t^H$  and  $\tilde{K}_t^H$  be the corresponding smoothing operators. Pick two sections  $u, v \in \Gamma(E)$  and denote  $\langle \cdot, \cdot \rangle$  as the inner product on  $E$ . Consider the smooth map

$$f(\theta) = \langle K_\theta^H u, \tilde{K}_{t-\theta}^H v \rangle \quad 0 < \theta < t \quad (4.35)$$

From the heat equation it follows that

$$\frac{\partial f}{\partial \theta} = \langle -H K_\theta^H u, \tilde{K}_{t-\theta}^H v \rangle + \langle K_\theta^H u, H \tilde{K}_{t-\theta}^H v \rangle. \quad (4.36)$$

This equals zero for all  $0 < \theta < t$ , because  $H$  is self-adjoint and hence  $f$  is constant. Taking the limits  $\theta \rightarrow 0$  and  $\theta \rightarrow t$  we conclude that

$$\langle K_t^H u, v \rangle = \langle u, \tilde{K}_t^H v \rangle \quad (4.37)$$

If we multiply  $u$  and  $v$  with bump functions, then for all  $x, y \in M$  and  $t \in \mathbb{R}_+$  we get  $(k_t^H)^*(x, y) = \tilde{k}_t^H(y, x)$ . If we choose  $\tilde{k}_t^H = k_t^H$ , we conclude that  $k_t^H$  is self adjoint. In the general case we get for all  $x, y \in M$  and  $t \in \mathbb{R}_+$

$$k_t^H(x, y) = (\tilde{k}_t^H)^*(y, x) = \tilde{k}_t^H(x, y) \quad (4.38)$$

Therefore the heat kernel is unique.  $\square$

With a slight modification of this proof we show that smoothing operators of heat kernels form a semi group. That is,  $K_{s+t}^H = K_s^H \circ K_t^H$ . This property will be useful later, when we show that the trace of  $e^{-tD^2}$  is well-defined.

**Lemma 4.8** (Berline et al. [2004], Proposition 2.17(3)). *Let  $(M, g)$  be a compact Riemannian manifold, let  $E$  be a vector bundle on  $M$  with a positive definite inner product and let  $H$  be a generalized Laplacian. If  $H$  admits a heat kernel  $k_t^H \in \Gamma(E \boxtimes E)$ , then the corresponding smoothing operator  $K_t^H$  satisfies*

$$K_{s+t}^H = K_s^H \circ K_t^H \quad (4.39)$$

for all  $s, t \in \mathbb{R}_+$ .

*Proof.* Let  $u, v \in \Gamma(E)$  be two sections on  $E$ . Consider the differentiable map

$$f(\theta) = \langle K_{\theta+t}^H u, K_{s-\theta}^H v \rangle \quad -t < \theta < s \quad (4.40)$$

From the heat equation it follows that

$$\frac{\partial f}{\partial \theta} = \langle -H K_{\theta+t}^H u, K_{s-\theta}^H v \rangle + \langle K_{\theta+t}^H u, H K_{s-\theta}^H v \rangle \quad (4.41)$$

Again, this equals zero for all  $-t < \theta < s$ , because  $H$  is self-adjoint. Hence,  $f$  is constant. Comparing  $f$  at different values of  $\theta$  we conclude

$$\langle K_t^H u, K_s^H v \rangle = \langle K_{s+t}^H u, v \rangle \quad (4.42)$$

and so  $K_s^H \circ K_t^H = K_{s+t}^H$ .  $\square$

#### 4.4 The formal solution of the heat kernel

In section 4.2 we found that  $\frac{1}{\sqrt{4\pi t}} e^{-x^2/4t}$  was the heat kernel for the Laplacian on  $\mathbb{R}$ . For an  $n$  dimensional Riemannian manifold this kernel can be generalized into  $\frac{1}{(4\pi t)^{n/2}} \exp(-r^2/4t)$  where  $r$  is the geodesic distance. In order to construct the heat kernel  $k_t^H$  we consider the map  $s_t$  which is defined by  $s_t = (4\pi t)^{n/2} \exp(r^2/4t) \cdot k_t^H$ . This map measures the difference between the 'Euclidian' heat kernel and the heat kernel we are interested in. We show that if  $s_t$  is a formal power series  $\sum_i t^i \Phi_i$ , then  $s_t$  has a unique solution.

In this section we work with the following setup: Let  $(M, g)$  be an  $n$  dimensional Riemannian manifold and let  $y \in M$ . Consider a neighborhood  $U_y$  such that the map  $\exp_y^{-1}: U_y \rightarrow T_y M$  is a chart of  $M$ . Let  $E \rightarrow U_y$  be a vector bundle with a positive definite inner product and let  $H$  be a generalized Laplacian on  $E$ . The "Euclidian" heat kernel we denote by

$$q_t(x, y) = \frac{1}{(4\pi t)^{n/2}} \exp(-d(x, y)^2/4t) \quad \forall x \in U_y \quad (4.43)$$

where  $d(x, y)$  is the geodesic distance between  $x$  and  $y$ .

**Definition 4.9.** A formal power series  $k_t^H \in \Gamma(E \boxtimes E)$  of the form

$$k_t^H(x, y) = q_t(x, y) \sum_{i=0}^{\infty} t^i \Phi_i(x, y) \quad (4.44)$$

is a formal solution of the heat equation if

$$\left(\frac{\partial}{\partial t} + H_x\right) k_t^H = 0 \quad (4.45)$$

If we let  $k_t^H = q_t \cdot s_t$  be a candidate for the heat kernel, then  $s_t$  must satisfy a differential equation. In the next lemmas we make this explicit.

**Lemma 4.10** (Roe [1998], Lemma 7.12 and 7.13). *Let  $s_t \in \Gamma(E \rightarrow U \times \mathbb{R}_+) \otimes E_y^*$  be a time-dependent section on  $E$  and let  $q_t(\cdot, y): U \times \mathbb{R}_+ \rightarrow \mathbb{R}$  be the map defined in equation 4.43. Then*

$$H(q_t \cdot s_t) - q_t H s_t = (\Delta_x q_t) s_t - 2 \nabla_{\nabla q_t} s_t \quad (4.46)$$

*Proof.* By definition of the generalized Laplacian, the only non-commuting part of  $H$  is the Laplacian. In a Riemannian normal coordinate system  $\{x_\mu\}$  it is given by  $-\sum_\mu \nabla_\mu \nabla_\mu$ . In this coordinate system the commutator between  $H$  and  $q_t$  is given by

$$[H, q_t] s_t = - \sum_\mu [\nabla_\mu \nabla_\mu, q_t] s_t \quad (4.47)$$

Using the properties of the commutator this can be written as

$$[H, q_t] s_t = - \sum_\mu \nabla_\mu [\nabla_\mu, q_t] s_t + [\nabla_\mu, q_t] \nabla_\mu s_t \quad (4.48)$$

$$= - \sum_\mu [\nabla_\mu, [\nabla_\mu, q_t]] s_t + 2 [\nabla_\mu, q_t] \nabla_\mu s_t \quad (4.49)$$

According to the Leibniz rule, the commutator between a connection and a smooth function is the Lie derivative and hence

$$[H, q_t] s_t = - \sum_\mu (\mathcal{L}_\mu \mathcal{L}_\mu q_t) s_t + 2 (\mathcal{L}_\mu q_t) \nabla_\mu s_t \quad (4.50)$$

Recall that  $-\sum_\mu (\mathcal{L}_\mu \mathcal{L}_\mu q_t)$  is the Laplacian of  $q_t$ . We only need to show that  $\sum_\mu \mathcal{L}_\mu q_t \frac{\partial}{\partial x^\mu}$  is the gradient of  $q_t$ . Indeed, the gradient is the dual of the exterior derivative. The dual of  $\sum_\mu \mathcal{L}_\mu q_t \frac{\partial}{\partial x^\mu}$  equals  $\sum_\mu \mathcal{L}_\mu q_t \, dx^\mu$ . This is the exterior derivative of  $q_t$  in local coordinates.  $\square$

We need to calculate the Laplacian and the gradient of  $q_t$  in a suitable coordinate system. Namely, take the Riemannian normal coordinate frame and pick the polar coordinates  $\{r, \phi_1, \dots, \phi_{n-1}\}$  on  $T_y M$ . In this coordinate frame the map  $q_t$  reduces to

$$q_t = \frac{1}{(4\pi t)^{n/2}} \exp(-r^2/4t) \quad (4.51)$$

For this calculation we also need the Hodge dual<sup>12</sup>, because the Laplacian can be written as  $- * d * d$ . The next lemma calculates  $* d r$  and after this we calculate  $\Delta q_t$  and  $\nabla q_t$ .

**Lemma 4.11.** *Let  $\{r, \phi_1, \dots, \phi_{n-1}\}$  be the polar coordinates on  $T_y M$  and use the exponential map as a coordinate frame around  $y$ . Then,*

$$* d r = r^{n-1} \sqrt{g} d \phi_1 \wedge \dots \wedge d \phi_{n-1} \quad (4.52)$$

where  $g$  is the determinant of the metric.

*Proof.* First we show that  $* d r$  is a multiple of  $d \phi_1 \wedge \dots \wedge d \phi_{n-1}$ . For simplicity we write  $d \phi_1 \wedge \dots \wedge d \phi_{n-1} = d \Omega$ . Indeed,  $d r$  can be expanded into

$$* d r = c_r d \Omega + \sum_i c_i d r \wedge d \phi_1 \wedge \dots \wedge d \phi_{i-1} \wedge d \phi_{i+1} \wedge \dots \wedge d \phi_{n-1} \quad (4.53)$$

where  $c_r, c_i \in \mathbb{R}$  are the components of the vector field. We calculate  $d \phi_k \wedge * d r$  in this local basis and using the Hodge dual. Comparing them gives

$$d \phi_k \wedge * d r = (-1)^{k+1} c_k d r \wedge d \Omega = \langle d \phi_k, d r \rangle \text{Vol}(g). \quad (4.54)$$

By Gauss lemma it follows that  $\langle d \phi_k, d r \rangle = 0$  in a local neighborhood around the origin and so  $c_k = 0$  for all  $k$ . This shows that  $* d r = c_r d \Omega$ .

We explicitly calculate  $c_r$  by computing  $d r \wedge * d r$ . Because  $\left\| \frac{\partial}{\partial r} \right\|_g = 1$  it follows that

$$d r \wedge * d r = c_r d r \wedge d \Omega = \langle d r, d r \rangle \text{Vol}(g) = \text{Vol}(g). \quad (4.55)$$

For polar coordinates the volume form is given by  $r^{n-1} \sqrt{g} d r \wedge d \Omega$ . Hence we conclude that  $* d r = r^{n-1} \sqrt{g} d \Omega$ .  $\square$

---

<sup>12</sup>Recall that the Hodge dual  $*$  is the unique operator which is defined by the property  $\alpha \wedge * \beta = \langle \alpha, \beta \rangle \text{Vol}(g)$  for all  $\alpha, \beta \in \Omega^\bullet(M)$ .

**Lemma 4.12.** *In the coordinates defined in Lemma 4.11 the gradient of  $q_t$  is*

$$\nabla q_t = -q_t \frac{r}{2t} \frac{\partial}{\partial r} \quad (4.56)$$

*and the Laplacian of  $q_t$  is*

$$\Delta q_t = q_t \left( -\frac{r^2}{4t^2} + \frac{n}{2t} + \frac{r}{4gt} \frac{\partial g}{\partial r} \right) \quad (4.57)$$

*where  $g$  is the determinant of the metric.*

*Proof.* The gradient is the dual of the exterior derivative and so

$$\nabla q_t = (d q_t)^\flat = \left( -\frac{q_t}{2t} r d r \right)^\flat = -q_t \frac{r}{2t} \frac{\partial}{\partial r}. \quad (4.58)$$

For the Laplacian we need to calculate  $- * d * d q_t$ . From Lemma 4.11, we know that

$$* d q_t = -\frac{q_t}{2t} r * d r = -q_t \frac{r^n}{2t} \sqrt{g} d \phi_1 \wedge \dots \wedge d \phi_{n-1}. \quad (4.59)$$

The Laplacian can now be easily calculated

$$\Delta q_t = - * q_t \left( \frac{r^2}{4t^2} - \frac{n}{2t} - \frac{r}{4gt} \frac{\partial g}{\partial r} \right) r^{n-1} \sqrt{g} d r \wedge d \phi_1 \wedge \dots \wedge d \phi_{n-1} \quad (4.60)$$

and this proves the result.  $\square$

**Theorem 4.13** (Roe [1998], Theorem 7.15). *For any generalized Laplacian  $H$ , there exists a unique formal solution of the heat equation of the form*

$$k_t(x, y) = \frac{1}{(4\pi t)^{n/2}} e^{-d(x,y)^2/4t} \sum_{i=0}^{\infty} t^i \Phi_i(x, y) \quad (4.61)$$

*such that  $\Phi_0(y, y)$  is the identity map on  $E_y$ .*

*Proof.* Denote  $s_t = \sum_i t^i \Phi_i$ . The kernel  $k_t^H = q_t s_t$  must satisfy the heat equation. By Lemmas 4.10 - 4.12 the heat equation applied on  $k_t^H$  equals

$$\left( \frac{\partial}{\partial t} + H \right) k_t^H = \frac{\partial}{\partial t} (q_t \cdot s_t) + q_t H k_t + (\Delta q_t) s_t - 2 \nabla_{\nabla q_t} s_t = 0 \quad (4.62)$$

$$= \frac{\partial q_t}{\partial t} s_t + q_t \left( \frac{\partial}{\partial t} + H - \frac{r^2}{4t^2} + \frac{n}{2t} + \frac{r}{4gt} \frac{\partial g}{\partial r} - 2 \frac{1}{t} \nabla_{r\partial/\partial r} \right) s_t \quad (4.63)$$

The  $t$ -derivative of  $q_t$  can be easily calculated and it equals

$$\frac{\partial}{\partial t} q_t = q_t \left( -\frac{n}{2t} + \frac{r^2}{4t^2} \right) \quad (4.64)$$

and so the heat equation simplifies to

$$\left( \frac{\partial}{\partial t} + H \right) k_t^H = q_t \left( \frac{\partial}{\partial t} + H + \frac{r}{4gt} \frac{\partial g}{\partial r} - 2 \frac{1}{t} \nabla_{r\partial/\partial r} \right) s_t = 0 \quad (4.65)$$

This induces a differential equation for  $s_t$ . By expanding  $s_t$  into  $\sum_i \Phi_i$  we get a differential equation for each factor of  $t$ . We get the system of equations

$$\left( \frac{r}{4g} \frac{\partial g}{\partial r} + \nabla_{r\partial/\partial r} \right) \Phi_0 = 0 \quad (4.66)$$

$$\left( i + \frac{r}{4g} \frac{\partial g}{\partial r} + \nabla_{r\partial/\partial r} \right) \Phi_i = -H \Phi_{i-1} \quad \forall i > 0 \quad (4.67)$$

This can be simplified into

$$\nabla_{\partial/\partial r} (g^{1/4} \Phi_0) = 0 \quad (4.68)$$

$$\nabla_{\partial/\partial r} (r^i g^{1/4} \Phi_i) = -r^{i-1} g^{1/4} H \Phi_{i-1} \quad \forall i > 0, \quad (4.69)$$

Equation 4.68 and 4.69 are first order differential equations. Both are uniquely determined by its initial value at the origin. We set the integration constant for  $\Phi_0$  such that  $\Phi_0(y, y)$  is the identity map on  $E_y$ . Note that  $r^i g^{1/4} \Phi_i$  will be of order  $r^i$  if and only if the integration constant is set to zero. So the requirement that  $\Phi_i$  is smooth determines  $\Phi_i$  uniquely.  $\square$

## 4.5 The existence of the heat kernel

We have seen that for any generalized Laplacian has a unique formal solution of the heat equation. However we do not know if this formal solution is globally defined

and we do not know if the power series converges. In this section we construct the heat kernel by considering globally defined approximations to the formal solution. These approximations are not necessary smooth, but we require them to be  $l$  times continuous differentiable for some  $l \in \mathbb{N}$ . In this paragraph we denote the space of  $l$ -times differentiable sections on a vector bundle  $E \rightarrow M$  as  $\Gamma^l(E)$ . We often use the norm

$$\|s\|_l(x) = \sup_{k \leq l} \sup_{v \in T_x M} \|(\mathcal{L}_v)^k s\|. \quad (4.70)$$

The first approximation to  $k_t^H = q_t \sum_i t^i \Phi_i$  we consider is the partial sum multiplied by a bump function. The next proposition states some properties this approximation has.

In this section we work with the following setup: Let  $E \rightarrow (M, g)$  be a vector bundle with a positive definite inner product over an  $n$  dimensional *compact* Riemannian manifold, let  $H$  be a generalized Laplacian and let  $k_t^H = \sum_i t^i \Phi_i$  be the formal solution of the heat equation. Let  $y \in M$  and consider a neighborhood  $U_y$  such that the map  $\exp_y^{-1}: U_y \rightarrow T_y M$  is a chart of  $M$ .

**Proposition 4.14** (Berline et al. [2004], Theorem 2.20). *For a small enough  $\epsilon > 0$  pick a smooth map  $\psi: \mathbb{R} \rightarrow \mathbb{R}$  such that*

$$\psi(x) = \begin{cases} 1 & \text{if } x < \epsilon^2/4 \\ 0 & \text{if } x > \epsilon^2 \end{cases} \quad (4.71)$$

*Then  $k_t^{H,N}(x, y) = \frac{1}{(4\pi t)^{n/2}} e^{-d(x,y)^2/4t} \cdot \psi(d^2(x, y)) \sum_{i=0}^N t^i \Phi_i(x, y)$  is a smooth family of sections of  $E \boxtimes E \rightarrow M \times M$  for which the following holds:*

1. *The smoothing operators  $K_t^{H,N}$  which has  $k_t^{H,N}$  as their kernel, form a uniform bounded family of operators on  $\Gamma^l(E)$  for all  $0 \leq t \leq T$  for all  $T$ .*
2. *For all  $l \in \mathbb{N}$  and  $s \in \Gamma^l(E)$  the norm  $\|K_t^{H,N} s - s\|_l$  tends to zero when  $t$  tends to zero.*
3. *For all  $l \in \mathbb{N}$ , there exists a constant  $C \in \mathbb{R}$  such that the kernel satisfies the estimate*

$$\left\| \left( \frac{\partial}{\partial t} + H \right) k_t^{H,N}(x, y) \right\| \leq C t^{N - \frac{n+l}{2}} \quad (4.72)$$

*Proof.* Pick  $\epsilon > 0$  such that the ball of radius  $\epsilon$  centered at  $y$  lies inside  $U_y$ . For this  $\epsilon$  the map  $k_t^{H,N}$  can be globally extended on  $M \times M$  by zero. Hence  $k_t^{H,N}$  is a smooth family of sections of  $E \boxtimes E$ . The corresponding smoothing operator equals

$$\left(K_t^{H,N} s\right)(x) = \frac{1}{(4\pi t)^{n/2}} \int_{y \in M} e^{-\frac{d(x,y)^2}{4t}} \psi(d(x,y)^2) \sum_{i=0}^N t^i \Phi_i(x,y) s(y) \text{Vol}(g) \quad (4.73)$$

The integrand is almost everywhere zero, except in the neighborhood of  $x \in M$ . For this neighborhood we pick the local coordinates  $y = \exp_x \mathbf{y}$  and we have

$$\left(K_t^{H,N} s\right)(x) = \frac{1}{(4\pi t)^{n/2}} \int_{\mathbf{y} \in T_x M} e^{-\frac{\|\mathbf{y}\|^2}{4t}} \sum_{i=0}^N t^i \Psi_i(x, \vec{y}) s(\exp_x(\mathbf{y})) d^n \mathbf{y} \quad (4.74)$$

for some compactly supported  $\Psi_i \in \Gamma(E \boxtimes E)$ . It follows from applying the vector space transformation  $\mathbf{y} = t^{1/2} v$  that

$$\left(K_t^{H,N} s\right)(x) = (4\pi)^{-n/2} \int_{\mathbb{R}^n} e^{-\|v\|^2/4} \sum_{i=0}^N t^i \Psi_i(x, t^{1/2} v) s(\exp_x(t^{1/2} v)) d^n v. \quad (4.75)$$

This is bounded for all  $t$ . Because  $[0, T]$  is compact for all  $T > 0$ , we conclude that  $K_t^N$  is uniformly bounded on  $[0, T]$ . This proves the first part.

In the limit  $t \rightarrow 0$  equation 4.75 equals

$$\left(K_0^{H,N} s\right)(x) = (4\pi)^{-n/2} \int_{\mathbb{R}^n} e^{-\|v\|^2/4} \Psi_0(x, 0) s(\exp_x(0)) d^n v \quad (4.76)$$

$$= (4\pi)^{-n/2} \int_{\mathbb{R}^n} e^{-\|v\|^2/4} \psi(d(x,x)^2) \Psi_0(x, x) s(x) d^n v. \quad (4.77)$$

Clearly, the distance between  $x$  and itself is zero and hence  $\psi(d(x,x)^2) = 1$ . Recall that  $\Psi_0(x, x)$  is the identity operator on  $E_x$ . So equation 4.77 simplifies to

$$\left(K_t^{H,N} s\right)(x) = s(x) \cdot (4\pi)^{-n/2} \int_{\mathbb{R}^n} e^{-\|v\|^2/4} d^n v. \quad (4.78)$$

By comparing this result to the Euclidean heat kernel we conclude that  $\lim_{t \rightarrow 0} K_t^{H,N} s = s$ . This shows the second part.

Finally we estimate  $\left\| \left( \frac{\partial}{\partial t} + H \right) k_t^{H,N}(x, y) \right\|$ . For simplicity we write  $r_t^N = \left( \frac{\partial}{\partial t} + H \right) k_t^{H,N}(x, y)$ . From lemmas Lemmas 4.10 - 4.12, we know that

$$r_t^N(x, y) = q_t \left( \frac{\partial}{\partial t} + H + \frac{1}{t} \nabla_{r\partial/\partial r} + \frac{r}{4gt} \frac{\partial g}{\partial r} \right) \psi(d(x, y)^2) \sum_{i=0}^N t^i \Phi_i(x, y) \quad (4.79)$$

$$\begin{aligned} &= q_t \psi(d(x, y)^2) \left( \frac{\partial}{\partial t} + H + \frac{1}{t} \nabla_{r\partial/\partial r} + \frac{r}{4gt} \frac{\partial g}{\partial r} \right) \sum_{i=0}^N t^i \Phi_i(x, y) + \\ &\quad + q_t \left( \Delta \psi(r^2) - 2 \nabla_{\nabla \psi(r^2)} + \frac{r}{t} \frac{\partial \psi(r^2)}{\partial r} \right) \sum_{i=0}^N t^i \Phi_i(x, y) \end{aligned} \quad (4.80)$$

The kernels  $\Phi_i$  will cancel most terms. However, we are left with

$$\begin{aligned} r_t^N(x, y) &= q_t \psi(d(x, y)^2) t^N H \Phi_N(x, y) + \\ &\quad + q_t \left( \Delta \psi(r^2) - 2 \nabla_{\nabla \psi(r^2)} + \frac{r}{t} \frac{\partial \psi(r^2)}{\partial r} \right) \sum_{i=0}^N t^i \Phi_i(x, y) \end{aligned} \quad (4.81)$$

The first terms is of order  $t^{N-n/2}$ . We show that this is true for

$$q_t \left( \Delta \psi(r^2) - 2 \nabla_{\nabla \psi(r^2)} + \frac{r}{t} \frac{\partial \psi(r^2)}{\partial r} \right) \sum_{i=0}^N t^i \Phi_i(x, y) \quad (4.82)$$

This vanishes if  $d(x, y) < \epsilon$ , because  $\psi$  is constant in this area. If this term is zero, then it is of order  $t^{N-n/2}$ . So we only need to show that equation 4.82 is of the right degree when  $d(x, y) > \epsilon$ .

If  $d(x, y) > \epsilon$ , then  $\|t^{-N+\frac{n}{2}} q_t(x)\|_0$  is bounded by  $\|t^{-N+\frac{n}{2}} q_t(\epsilon)\|_0$ . This follows from the fact that  $q_t$  is a decreasing map. For  $t > 1$  the norm  $\|t^{-N+\frac{n}{2}} q_t(x)\|_0$  is bounded by one. Because  $t^{-N+\frac{n}{2}} q_t(\epsilon)$  is continuous in  $t$  and  $[0, 1]$  is compact follows that  $t^{-N+\frac{n}{2}} q_t(\epsilon)$  has an upper bound. So  $r_t^N$  is bounded in the supremum norm by  $t^{N-\frac{n}{2}}$ .

To consider higher order derivatives of  $r_t^N(x, y)$  in  $x$ , we note that we can only lower the degree of  $t$  by differentiating over  $q_t$ . Because  $\frac{\partial}{\partial x} q_t = \mathcal{O}(t^{-1/2})$ , we conclude that  $\|r_t^N\|_l < C t^{N-\frac{n+l}{2}}$ .  $\square$

New kernels can be constructed using old kernels. Indeed let  $p, q \in \Gamma(E \boxtimes E)$  be two kernels. The composition  $\int_{z \in M} p(x, z) q(z, y) \text{Vol}(g)$  is a map from  $E_y$  to  $E_x$ .

Therefore it is a section on  $E \boxtimes E$ . For smooth families of kernels  $p_t, q_t \in \Gamma(E \boxtimes E \rightarrow M \times M \times \mathbb{R}_+)$  we consider the composition

$$(x, y, t) \mapsto \int_0^t ds \int_{z \in M} p_{t-s}(x, z) q_s(z, y) \text{Vol}(g) \quad (4.83)$$

This is also a section of  $\Gamma(E \boxtimes E \rightarrow M \times M \times \mathbb{R}_+)$ . As our second approximation attempt we consider compositions of  $k_t^{H,N}$  and  $(\frac{\partial}{\partial t} + H_x) k_t^{H,N}$ . To simplify our notation we inductively define

$$r_t^{H,N,1}(x, y) = \left( \frac{\partial}{\partial t} + H_x \right) k_t^{H,N} \quad (4.84)$$

$$r_t^{H,N,m+1}(x, y) = \int_0^t ds \int_{z \in M} r_{t-s}^{H,N,1}(x, z) r_s^{H,N,m}(z, y) \text{Vol}(g) \quad (4.85)$$

$$k_t^{H,N,0}(x, y) = k_t^{H,N}(x, y) \quad (4.86)$$

$$k_t^{H,N,m}(x, y) = \int_0^t ds \int_{z \in M} k_{t-s}^{H,N}(x, z) r_s^{H,N,m}(z, y) \text{Vol}(g) \quad (4.87)$$

In the following lemma we give an estimation of  $k_t^{H,N,m}$  and  $r_t^{H,N,m}$ .

**Lemma 4.15** (Berline et al. [2004], Lemma 2.21 and 2.22(1)). *Let  $k_t^{H,N}$  be the family of kernels defined in Proposition 4.14. Then for all  $l, m \in \mathbb{N}$  and  $N > \frac{n+l}{2}$  the kernel  $r_t^{H,N,m+1}(x, y)$ , which is defined in equation 4.84 and 4.85, is  $l$ -times continuous differentiable with respect to  $x$  and  $y$  and satisfies the estimate*

$$\|r_t^{H,N,m+1}\|_l \leq C^{m+1} t^{(m+1)(N - \frac{n+l}{2})} \text{Vol}(M)^m \frac{t^m}{m!} \quad (4.88)$$

*for some  $C \in \mathbb{R}$ . The kernel  $k_t^{H,N,m}$ , which is defined in equation 4.86 and 4.87, is also  $l$ -times continuous differentiable with respect to  $x$  and  $y$  and satisfies the estimate*

$$\|k_t^{H,N,m+1}\|_l \leq \tilde{C} C^{m+1} t^{(m+1)(N - \frac{n+l}{2})} \text{Vol}(M)^m \frac{t^m}{m!}. \quad (4.89)$$

*for some  $\tilde{C} \in \mathbb{R}$ .*

*Proof.* From Theorem 4.14 follows that  $r_t^{H,N,1}$  is bounded by  $t$ . Hence we extend  $r_t^{H,N,1}$  (and its derivatives) continuously to  $t = 0$  and so  $r_t^{H,N,1}$  has a well defined

$l$ -norm. This is equivalent to the fact that it is  $l$ -times continuously differentiable.

We show the first estimate using induction. From Definition 4.84 and Theorem 4.14 it follows that

$$\|r_t^{H,N,0+1}\|_l \leq C t^{N-\frac{n+l}{2}}. \quad (4.90)$$

This coincides with equation 4.88 for  $m = 0$ . Now assume that equation 4.88 holds for some  $m \in \mathbb{N}$ . Then  $r_t^{H,N,m+1}$  is estimated by

$$\|r_t^{H,N,m+1}(x, y)\|_l \leq \int_0^t ds \int_{z \in M} \|r_{t-s}^{H,N,1}(x, z)\|_l \cdot \|r_s^{H,N,m}(x, z)\|_l \quad (4.91)$$

From the induction hypothesis it follows that

$$\|r_t^{H,N,m+1}(x, y)\|_l \leq \int_0^t ds \int_{z \in M} C(t-s)^{N-\frac{n+l}{2}} \cdot C^m s^{m(N-\frac{n+l}{2})} \text{Vol}(M)^{m-1} \frac{s^{m-1}}{m-1!} \quad (4.92)$$

and this simplifies to

$$\|r_t^{H,N,m+1}(x, y)\|_l \leq C^{m+1} \text{Vol}(M)^m \int_0^t ds (t-s)^{N-\frac{n+l}{2}} s^{m(N-\frac{n+l}{2})} \frac{s^{m-1}}{m-1!} \quad (4.93)$$

$$\leq C^{m+1} t^{(m+1)(N-\frac{n+l}{2})} \text{Vol}(M)^m \int_0^t ds \frac{s^{m-1}}{m-1!}. \quad (4.94)$$

Integrating  $\int_0^t ds \frac{s^{m-1}}{m-1!}$  yields a factor  $\frac{t^m}{m!}$  and this proves that equation 4.88 is satisfied for  $m + 1$ . By induction we conclude that it is satisfied for all  $m \in \mathbb{N}$ .

From Theorem 4.14 it follows that the smoothing operators w.r.t.  $k_t^N$  are uniform bounded on  $0 \leq t \leq T$  for all  $T$ . Hence for all  $s \in [0, t]$  and  $x, y \in M$

$$\left\| \int_{z \in M} k_{t-s}^{H,N}(x, z) r_s^{H,N,m}(z, y) \text{Vol}(g) \right\|_l \leq \tilde{C} \|r_s^{H,N,m}(x, y)\|_l \quad (4.95)$$

for some  $\tilde{C} \in \mathbb{R}$ . Notice that the left hand side is a norm estimate of  $k_t^{H,N,m}$ . From this and equation 4.88 we conclude the result.  $\square$

The kernels  $r_t^{H,N,m}$  and  $k_t^{H,N,m}$  are bounded by  $\frac{(C \text{Vol}(M) t^{N-\frac{n+l}{2}})^m}{m!}$ . Hence we can use them for constructing convergent power series, because we can bound these series by  $\exp(C \text{Vol}(M) t^{N-\frac{n+l}{2}})$ . To find which series is the heat kernel, we need to investigate how the heat equation behaves for  $k_t^{H,N,m}$ :

**Lemma 4.16** (Berline et al. [2004], Lemma 2.22(2)). *For all  $l, m \in \mathbb{N}$  and  $N > \frac{n+l}{2}$  the kernel  $k_t^{H,N,m}(x, y)$ , which is defined in equation 4.86 and 4.87, satisfies*

$$\left(\frac{\partial}{\partial t} + H_x\right) k_t^{H,N,m} = r_t^{H,N,m+1} + r_t^{H,N,m} \quad (4.96)$$

*Proof.* Consider the map  $b(t, s, x, y) = \int_{z \in M} k_{t-s}^{H,N}(x, z) r_s^{H,N,m}(z, y) \text{Vol}(g)$ . The integral  $\int_0^t b(t, s, x, y) \, ds$  equals the map  $k_t^{H,N,m}$ . The map  $b(t, s, x, y)$  is continuous in  $s \in [0, t]$ , because the smoothing operator  $K_t^N$  is uniform bounded. So the heat equation applied on  $k_t^{H,N,m}$  equals

$$\left(\frac{\partial}{\partial t} + H_x\right) k_t^{H,N,m} = \left(\frac{\partial}{\partial t} + H_x\right) \int_0^t b(t, s, x, y) \, ds \quad (4.97)$$

$$= b(t, t, x, y) - b(t, 0, x, y) + \int_0^t \left(\frac{\partial}{\partial t} + H_x\right) b(t, s, x, y) \, ds \quad (4.98)$$

From Theorem 4.14 it follows that  $b(t, 0, x, y) = \int_{z \in M} k_t^{H,N}(x, z) r_0^{H,N,m}(z, y) \text{Vol}(g) = 0$ . Also,  $b(t, t, x, y) = \int_{z \in M} k_0^{H,N}(x, z) r_t^{H,N,m}(z, y) \text{Vol}(g) = r_t^{H,N,m}(x, y)$ . Equation 4.98 therefore becomes

$$\left(\frac{\partial}{\partial t} + H_x\right) k_t^{H,N,m} = r_t^{H,N,m}(x, y) + \int_0^t \left(\frac{\partial}{\partial t} + H_x\right) b(t, s, x, y) \, ds. \quad (4.99)$$

We calculate the heat equation acted on  $b(t, s, x, y)$ . By definition it equals:

$$\left(\frac{\partial}{\partial t} + H_x\right) b(t, s, x, y) = \left(\frac{\partial}{\partial t} + H_x\right) \int_{z \in M} k_{t-s}^{H,N}(x, z) r_s^{H,N,m}(z, y) \text{Vol}(g) \quad (4.100)$$

The heat equation acts only on terms which depend on  $t$  and  $x$  and so

$$\left(\frac{\partial}{\partial t} + H_x\right) b(t, s, x, y) = \int_{z \in M} \left(\left(\frac{\partial}{\partial t} + H_x\right) k_{t-s}^{H,N}(x, z)\right) r_s^{H,N,m}(z, y) \text{Vol}(g) \quad (4.101)$$

$$= \int_{z \in M} r_{t-s}^{H,N,1}(x, z) r_s^{H,N,m}(z, y) \text{Vol}(g). \quad (4.102)$$

By equation 4.85 it follows that

$$\left(\frac{\partial}{\partial t} + H_x\right) k_t^{H,N,m} = r_t^{H,N,m}(x, y) + r_t^{H,N,m+1}(x, y) \quad (4.103)$$

and we finish the proof.  $\square$

Assume that the heat kernel is of the form  $\sum_{m=0}^{\infty} c_m k_t^{H,N,m}$ . We now search for which values of  $c_m \in \mathbb{R}$  the heat equation is satisfied. From the previous lemma it follows that

$$\left(\frac{\partial}{\partial t} + H\right) \sum_{m=0}^{\infty} c_m k_t^{H,N,m} = \sum_m c_m (r_t^{H,N,m} + r_t^{H,N,m+1}) \quad (4.104)$$

$$= \sum_m r_t^{H,N,m} (c_m + c_{m-1}) \quad (4.105)$$

The alternating sum of  $k_t^{H,N,m}$  will satisfy the heat equation. We show that it is indeed the heat kernel.

**Theorem 4.17** (Berline et al. [2004], Theorem 2.23). *Let  $(M, g)$  be a compact Riemannian manifold of dimension  $n$  and let  $E \rightarrow M$  be a vector bundle with a positive definite inner product. Let  $H$  be a generalized Laplacian and let  $k_t$  be the formal solution to the heat kernel. Then the following is true:*

1. *For any  $N \in \mathbb{N}$  and  $l \in \mathbb{N}$  such that  $N > \frac{n+l+1}{2}$ , the series*

$$p_t^{H,N}(x, y) = \sum_{m=0}^{\infty} (-1)^m k_t^{H,N,m}(x, y) \quad (4.106)$$

*converges in the  $\|\cdot\|_{l+1}$ -norm over  $M \times M$  and it is continuous differentiable in  $t$ . It satisfies the heat equation.*

2. *The kernel  $k_t^{H,N} \in \Gamma(E \boxtimes E)$  approximates  $p_t$  in the sense that*

$$\left\| \frac{\partial^m}{\partial t^m} (p_t^{H,N} - k_t^{H,N}) \right\|_l = \mathcal{O}(t^{N - \frac{n+l}{2} - m + 1}) \quad (4.107)$$

*for all  $m \in \mathbb{N}$  and  $N > \frac{n+l+1}{2}$  when  $t$  approaches zero.*

3. *The kernel  $p_t^H = p_t^{H,n+1}$  is a heat kernel for the operator  $H$ .*

*Proof.* In the above discussion we showed that  $p_t$  converges because it can be estimated using the exponential series. Therefore the kernel  $p_t$  converges in the  $l+1$ -norm. From Lemma 4.16 it follows that

$$\frac{\partial}{\partial t} k_t^{H,N,m} = r_t^{H,N,m+1} + r_t^{H,N,m} - H_x k_t^{H,N,m} \quad (4.108)$$

and we estimate

$$\left\| \frac{\partial}{\partial t} p_t \right\|_l = \lim_{M \rightarrow \infty} \left\| \sum_{m=0}^M (-1)^m (r_t^{H,N,m+1} + r_t^{H,N,m} - H_x m_t^{H,N,m}) \right\|_l \quad (4.109)$$

$$= \lim_{M \rightarrow \infty} \left\| (-1)^M r_t^{H,N,M+1} - H_x p_t \right\|_l \quad (4.110)$$

This converges for all  $t$  and so  $p_t$  is continuous differentiable in the  $t$  component. It also satisfies the heat equation. This prove the first part.

To show the second part, recall that  $q_t^{H,N,0} = k_t^{H,N}$ . From Lemma 4.15 we estimate

$$\|p_t - k_t^N\|_l = \left\| \sum_{j=1}^{\infty} (-1)^j q_t^{j,N} \right\|_l = \mathcal{O}(t^{N - \frac{n+l}{2} + 1}) \quad (4.111)$$

The second part of the theorem follows after  $m$ -times differentiation.

Before we show that  $p_t^H$  is a heat kernel, we prove that  $p_t^{H,N}$  is a  $C^l$ -heat kernel if  $N > \frac{n+l+1}{2}$ . We only need to show the boundary condition

$$\lim_{t \rightarrow 0} \int_{y \in M} p_t^{H,N}(x, y) s(y) = s(x) \quad \text{for all } s \in \Gamma^l(E). \quad (4.112)$$

From the second part of this theorem we know that  $\lim_{t \rightarrow 0} \left\| \int_{y \in M} p_t^{H,N}(x, y) s(y) - s(x) \right\|_l = \lim_{t \rightarrow 0} \left\| \int_{y \in M} k_t^{H,N}(x, y) s(y) - s(x) \right\|_l$ . By Theorem 4.14 is equals zero. Hence,  $p_t^{H,N}$  is a  $C^l$ -heat kernel w.r.t.  $H$ .

Finally notice that  $\Gamma^l(E) \subseteq \Gamma^0(E)$ . Hence,  $p_t^{H,N}$  and  $p_t^{H,n+1}$  are both  $C^0$ -heat kernels. By unicity of the heat kernel it follows that they are equal. Hence,  $p_t^{H,n+1}$  is the smooth heat kernel w.r.t.  $H$ .  $\square$

## 5 Traces and the Index of a Dirac operator

In chapter 3 we saw how the Fujikawa method relates the chiral anomaly to the index of a Dirac operator. Although we calculated the trace over  $\gamma_5 e^{-tD^2}$  we did not check whether it converges. We also never checked if the trace was independent of basis. Now we rigorously introduce a class of operators for which the trace is well defined and show that the heat kernel is in this class.

Because the trace is the sum of the eigenvalues, we retrieve some properties of the eigenvalues of the heat kernel. Using the heat equation we get that a generalized Laplacian has countably many eigenvalues. Using this spectrum analysis we can prove McKean-Singer formula. It relates the index of a Dirac operator with the trace of the heat kernel.

Next we investigate how the trace interacts with the Clifford action. We prove the trace identities which we used in chapter 2 and we generalize them to higher dimensions.

At last we study the topological properties of traces. We study the trace over differential forms and we show that they characterize vector bundles. We introduce the characteristic classes needed for the Atiyah-Singer index theorem.

The theory of traceclass operators is standard. For more information see Murphy [1990].

### 5.1 Traceclass operators

**Definition 5.1.** A linear operator  $A$  on a Hilbert space  $\mathcal{H}$  is called **Hilbert-Schmidt** if  $\sum_i \|Ae_i\|^2$  is finite for all orthonormal basis  $\{e_i\}$ .

Hilbert-Schmidt operators are bounded operators. Indeed let  $\mathcal{H}$  be a Hilbert space,  $A$  a Hilbert-Schmidt operator and let  $v \in \mathcal{H}$ . In an orthonormal basis  $\{e^i\}$  of  $H$  we can write  $v$  as  $\sum_i v_i e^i$ . Let  $v_{max}$  be the largest component of  $v$ . That is,  $v_{max} = \max_i |v_i|$ . We can estimate

$$\frac{\|Av\|^2}{\|v\|^2} \leq \sum_i \frac{|v_i|^2}{\|v\|^2} \cdot \|Ae^i\|^2 \leq \frac{v_{max}^2}{\|v\|^2} \sum_i \|Ae^i\|^2 \leq \sum_i \|Ae^i\|^2 \quad (5.1)$$

and so the operator norm is bounded by

$$\|A\|_{op}^2 = \sup_{v \in \mathcal{H}} \frac{\|Av\|^2}{\|v\|^2} \leq \sup_{v \in \mathcal{H}} \sum_i \|Ae^i\|^2 = \sum_i \|Ae^i\|^2. \quad (5.2)$$

By definition  $\sum_i \|Ae^i\|^2$  is finite and hence  $A$  is bounded.

Linear combinations of Hilbert-Schmidt operators are also Hilbert-Schmidt. To see this let  $A$  and  $B$  be Hilbert-Schmidt operators on a Hilbert space  $\mathcal{H}$ , let  $\lambda \in \mathbb{C}$  and let  $\{e^i\}$  be an orthonormal basis of  $\mathcal{H}$ . For all  $N \in \mathbb{N}$  we estimate

$$\sum_{i=0}^N \|(A + \lambda B)e^i\|^2 \leq \sum_{i=0}^N \|Ae^i\|^2 + |\lambda|^2 \cdot \sum_{i=0}^N \|Be^i\|^2 \quad (5.3)$$

and in the limit  $N \rightarrow \infty$  we conclude that  $\sum_{i=0}^\infty \|(A + \lambda B)e^i\|^2$  is finite.

The product between a Bounded operator  $B$  and a Hilbert-Schmidt operator  $A$  is also Hilbert-Schmidt. This follows from the estimate

$$\sum_i \|BAe^i\|^2 \leq \|B\|_{op}^2 \cdot \sum_i \|Ae^i\|^2 \quad (5.4)$$

This shows that the space of Hilbert-Schmidt operators is an ideal in the space of Bounded operators.

**Lemma 5.2.** *Let  $\mathcal{H}$  be a Hilbert space. Then for all orthonormal frames  $\{e^i\}$  of  $\mathcal{H}$  the map  $A \mapsto \sqrt{\sum_i \|Ae^i\|^2}$  is a norm on the space of Hilbert-Schmidt operators.*

*Proof.* Let  $A$  be a Hilbert-Schmidt operator. Clearly,  $\sum_i \|Ae^i\|^2 \geq 0$  and so this norm is positive. It is also definite. In the case that  $\sum_i \|Ae^i\|^2 = 0$ , then by Equation 5.2 the operator norm of  $A$  is zero. Hence the norm is positive definite.

This map is also absolutely scalable: For all  $N \in \mathbb{N}$  and  $\lambda \in \mathbb{C}$  the finite sum  $\sum_{i=0}^N \|\lambda Ae^i\|^2$  equals  $|\lambda|^2 \sum_{i=0}^N \|Ae^i\|^2$ . Absolutely scalability follows if we take the limit  $N \rightarrow \infty$ .

Finally we show that it satisfies the triangle inequality. Let  $A$  and  $B$  be Hilbert-Schmidt operators. Then for all  $N \in \mathbb{N}$  these operators satisfy

$$\sum_{i=0}^N \|(A + B)e^i\|^2 \leq \sum_{i=0}^N \|Ae^i\|^2 + \sum_{i=0}^N \|Be^i\|^2 \leq \sum_{i=0}^\infty \|Ae^i\|^2 + \sum_{i=0}^\infty \|Be^i\|^2 \quad (5.5)$$

The triangle inequality follows if we take  $N \rightarrow \infty$ .  $\square$

**Lemma 5.3.** *If  $A$  is a Hilbert-Schmidt operator on a Hilbert space  $\mathcal{H}$  then the following is true:*

1. *The adjoint  $A^*$  is a Hilbert-Schmidt operator.*
2. *The norm  $\sqrt{\sum_i \|A^* e^i\|^2}$  equals  $\sqrt{\sum_i \|A e^i\|^2}$ .*
3. *The norm  $\sqrt{\sum_i \|A e^i\|^2}$  does not depend on the choice of orthonormal basis  $\{e^i\}$ .*

*Proof.* Let  $\{e^i\}$  and  $\{\tilde{e}^i\}$  be two orthonormal bases of  $\mathcal{H}$ . By Parsevals identity it follows that for all  $N \in \mathbb{N}$

$$\sum_{i=0}^N \|A^* e^i\|^2 = \sum_{i=0}^N \sum_{j=0}^{\infty} |\langle A^* e^i, \tilde{e}^j \rangle|^2 = \sum_{j=0}^{\infty} \sum_{i=0}^N |\langle e^i, A \tilde{e}^j \rangle|^2 \leq \sum_{j=0}^{\infty} \|A \tilde{e}^j\|^2 \quad (5.6)$$

Because  $\sum_{j=0}^{\infty} \|A \tilde{e}^j\|^2$  is finite, we conclude that  $A^*$  is Hilbert-Schmidt. Using a similar method we can show that  $\sum_{i=0}^{\infty} \|A e^i\|^2 \leq \sum_{i=0}^{\infty} \|A^* \tilde{e}^i\|^2$ . Hence  $A$  has the same norm as  $A^*$  and

$$\sqrt{\sum_i \|A e_i\|^2} = \sqrt{\sum_i \|A^* \tilde{e}_i\|^2} = \sqrt{\sum_i \|A e_i\|^2} \quad (5.7)$$

So the norm is independent of the choice of basis.  $\square$

**Definition 5.4.** *Let  $\mathcal{H}$  be a Hilbert space and let  $\{e^i\}$  be an orthonormal basis of  $\mathcal{H}$ . The **Hilbert-Schmidt norm** is the norm on the space of all Hilbert-Schmidt operators on  $\mathcal{H}$  defined by*

$$\|A\|_{HS}^2 = \sum_i \|A e^i\|^2 \quad (5.8)$$

Given a norm, we can ask whether it is induced from an inner product. Recall that any Hermitian inner product  $\langle \cdot, \cdot \rangle$  on Hilbert space  $\mathcal{H}$  satisfies the polarization identity

$$\langle u, v \rangle = \frac{1}{4} \sum_{k=0}^3 i^k \|u + i^k v\|^2 \quad \forall u, v \in \mathcal{H} \quad (5.9)$$

Hence we investigate if the pairing  $(A, B) \mapsto \frac{1}{4} \sum_{k=0}^3 i^k \|A + i^k B\|_{HS}^2$  is an inner product on the space of Hilbert-Schmidt operators.

**Lemma 5.5.** *Let  $\mathcal{H}$  be a Hilbert space. The function that maps two Hilbert-Schmidt operators  $A$  and  $B$  to*

$$\frac{1}{4} \sum_{k=0}^3 i^k \|A + i^k B\|_{HS}^2 \quad (5.10)$$

*is an inner product on the space of Hilbert-Schmidt operators. Given an orthonormal basis  $\{e^i\}$  this inner product is given by*

$$(A, B) \mapsto \sum_i \langle Ae^i, Be^i \rangle \quad (5.11)$$

*Proof.* This pairing is clearly positive definite, because the norm is positive definite. We only need to check for linearity and conjugate-symmetry. Let  $\{e^i\}$  be an orthonormal basis of  $\mathcal{H}$ . Then the pairing of  $A$  and  $B$  equals

$$\frac{1}{4} \sum_{k=0}^3 \sum_j i^k \|Ae^j + i^k Be^j\|_{HS}^2 \quad (5.12)$$

Applying the polarization identity on  $\mathcal{H}$  Equation 5.12 simplifies to

$$\frac{1}{4} \sum_j \sum_{k=0}^3 i^k \|Ae^j + i^k Be^j\|_{HS}^2 = \sum_j \langle Ae^j, Be^j \rangle \quad (5.13)$$

This concludes Equation 5.11. From this Equation we directly conclude that the pairing is linear and skew-symmetric.  $\square$

**Definition 5.6.** *Let  $\mathcal{H}$  be a Hilbert space,  $\{e^i\}$  be an orthonormal basis of  $\mathcal{H}$  and let  $A$  and  $B$  be two Hilbert-Schmidt operators. The **Hilbert-Schmidt inner product** is the inner product that maps  $A$  and  $B$  to*

$$\sum_i \langle Ae^i, Be^i \rangle \quad (5.14)$$

*and is denoted by  $\langle \cdot, \cdot \rangle_{HS}$ .*

For a finite dimensional Hilbert space  $\mathcal{H}$ , all linear operators are Hilbert Schmidt. Note that for any linear operator  $A, B$  on  $\mathcal{H}$ ,

$$\langle A, B \rangle_{HS} = \sum_i \langle Ae_i, Be_i \rangle = \sum_i \langle B^* Ae_i, e_i \rangle = \text{Tr}(B^* A) \quad (5.15)$$

Equation 5.15 suggests that we can define a trace for products of Hilbert Schmidt operators.

**Definition 5.7.** A linear operator  $T$  on a Hilbert space  $\mathcal{H}$  is **traceclass** if it is the product of two Hilbert-Schmidt operators.

Let  $T$  be a traceclass operator. Hence, there exists two Hilbert-Schmidt operators  $A$  and  $B$  such that  $T = AB$ . Now suppose that there is another decomposition of  $T$ . That is, there exists also some Hilbert-Schmidt operators  $\tilde{A}$  and  $\tilde{B}$  such that  $T = AB = \tilde{A}\tilde{B}$ . In both cases we have

$$\begin{aligned} \langle B, A^* \rangle_{HS} &= \sum_i \langle Be^i, A^* e^i \rangle = \sum_i \langle ABe^i, A^* e^i \rangle = \sum_i \langle Te^i, e^i \rangle \\ \langle \tilde{B}, \tilde{A}^* \rangle_{HS} &= \sum_i \langle \tilde{B}e^i, \tilde{A}^* e^i \rangle = \sum_i \langle \tilde{A}\tilde{B}e^i, A^* e^i \rangle = \sum_i \langle Te^i, e^i \rangle \end{aligned} \quad (5.16)$$

and we conclude that  $\langle B, A^* \rangle_{HS}$  does not depend on the decomposition of  $T$ . Using Equation 5.15 we extend the definition of the trace to traceclass operators.

**Definition 5.8.** Let  $T$  be a traceclass operator and let  $A$  and  $B$  Hilbert-Schmidt operators such that  $T = AB$ . The **trace** of  $T$  is the Hilbert-Schmidt inner product  $\text{Tr}(T) = \langle B, A^* \rangle_{HS}$

In the previous chapter we studied generalized Laplacians. We showed that for each generalized Laplacian  $H$  there exists a smoothinh operator  $e^{-tH}$  that satisfies the heat equation. We now show that  $e^{-tH}$  is traceclass. Observe that it is enough to show that  $e^{-tH}$  is Hilbert-Schmidt, because of the semi-group property. Indeed, if we show that  $e^{-tH}$  is Hilbert-Schmidt for all  $t \in \mathbb{R}_+$  then  $e^{-\frac{t}{2}H} \cdot e^{-\frac{t}{2}H}$  is traceclass. By the semi-group property  $e^{-\frac{t}{2}H} \cdot e^{-\frac{t}{2}H}$  simplifies to  $e^{-tH}$ .

**Proposition 5.9.** *Let  $E$  be a vector bundle with a positive definite inner product over a compact Riemannian manifold  $(M, g)$  and let  $H$  be a generalized Laplacian. Let  $k_t^H$  be the heat kernel and let  $e^{-tH}$  be the corresponding smoothing operator of  $k_t^H$ . Then  $e^{-tH}$ , which we view as an operator on the space of sections of  $E$  with  $L^2$  norm, is traceclass.*

*Proof.* First we observe that it is enough to prove that  $e^{-tH}$  is Hilbert-Schmidt for all  $t \in \mathbb{R}_+$ , since in this case  $e^{-tH} = e^{-\frac{t}{2}H} \cdot e^{-\frac{t}{2}H}$  by the semi-group property. Pick an orthonormal basis  $\{e^i\}$  on the space of sections of  $E$ . The Hilbert-Schmidt norm of  $e^{-tH}$  equals

$$\|e^{-tH}\|_{HS}^2 = \sum_{i=0}^{\infty} \|e^{-tH}e^i\|_{L^2}^2 = \sum_{i=0}^{\infty} \langle e^{-tH}e^i, e^{-tH}e^i \rangle_{L^2} \quad (5.17)$$

In Lemma 4.7 we showed that  $e^{-tD^2}$  is self-adjoint. Using this and the fact that  $e^{-tH}$  is a smoothing operator we conclude that the Hilbert-Schmidt norm of  $e^{-tH}$  equals

$$\|e^{-tH}\|_{HS}^2 = \sum_{i=0}^{\infty} \langle e^{-2tH}e^i, e^i \rangle_{L^2} \quad (5.18)$$

$$= \sum_{i=0}^{\infty} \int_{x \in M} \langle k_{2t}(\cdot, x)e^i(x), e^i \rangle_{L^2} \quad (5.19)$$

Let  $\{f_j\}$  be an orthonormal frame of the fiber  $E_x$ . Using the identity  $e^i(x) = \sum_{j=0}^{\dim E} \langle e^i(x), f^j \rangle f^j$  we see

$$\|e^{-tH}\|_{HS}^2 = \sum_{j=0}^{\dim E} \sum_{i=0}^{\infty} \int_{x \in M} \langle \langle k_{2t}(\cdot, x)f^j, e^i \rangle_{L^2} \cdot e^i(x), f^j \rangle \quad (5.20)$$

Because  $\{e^i\}$  is a basis of  $\Gamma(E)$  we conclude that  $\sum_{i=0}^{\infty} \langle k_{2t}(\cdot, x)f^j, e^i \rangle_{L^2} \cdot e^i$  equals  $k_{2t}(\cdot, x)$  and hence

$$\|e^{-tH}\|_{HS}^2 = \sum_{j=0}^{\dim E} \int_{x \in M} \langle k_{2t}(x, x)f^j, f^j \rangle = \int_{x \in M} \text{tr}(k_{2t}(x, x)) \quad (5.21)$$

$$(5.22)$$

Because the manifold is compact, we conclude that  $e^{-tH}$  is Hilbert-Schmidt.  $\square$

**Proposition 5.10** (Berline et al. [2004], Proposition 2.32). *Let  $E$  be a vector bundle with a positive definite inner product over a compact Riemannian manifold  $(M, g)$ , let  $H$  be a generalized Laplacian, let  $k_t^H$  be its heat kernel and let  $e^{-tH}$  be the corresponding smoothing operator. Let  $B$  be a bounded operator on  $\Gamma(E)$ . The operator  $Be^{-tH}$  is trace-class and its trace is given by*

$$\mathrm{Tr}(Be^{-tH}) = \int_{x \in M} \mathrm{tr}(Bk_t(x, x)) \quad (5.23)$$

*Proof.* Because Hilbert-Schmidt operators is an ideal in the space of bounded operators it follows that  $Be^{-t/2H}$  is Hilbert-Schmidt. Hence the product  $Be^{-t/2H} \cdot e^{-t/2H} = Be^{-tH}$  is traceclass.

To show the explicit expression of the trace, let  $\{e^i\}$  be an orthonormal basis of the space of all sections on  $E$ . Equip  $\Gamma(E)$  with the  $L^2$  norm. In this orthonormal basis the trace equals

$$\mathrm{Tr}(Be^{-tH}) = \sum_{i=0}^{\infty} \langle Be^{-tH} e^i, e^i \rangle_{L^2} \quad (5.24)$$

From the definition of a smoothing operator follows that

$$\mathrm{Tr}(Be^{-tH}) = \sum_{i=0}^{\infty} \int_{x \in M} \langle Bk_t(\cdot, x) e^i(x), e^i \rangle_{L^2} \quad (5.25)$$

Let  $f^j$  be an orthonormal basis of  $E_x$ . Using the identity  $e^i(x) = \sum_{j=1}^{\dim E_x} \langle e^i(x), f^j \rangle f^j$  we see

$$\mathrm{Tr}(Be^{-tH}) = \sum_{j=1}^{\dim E_x} \sum_{i=0}^{\infty} \int_{x \in M} \langle e^i(x), f^j \rangle \cdot \langle Bk_t(\cdot, x) f^j, e^i \rangle_{L^2} \quad (5.26)$$

$$= \sum_{j=1}^{\dim E_x} \sum_{i=0}^{\infty} \int_{x \in M} \langle \langle Bk_t(\cdot, x) f^j, e^i \rangle_{L^2} \cdot e^i(x), f^j \rangle \quad (5.27)$$

Because  $\{e^i\}$  is a basis of  $\Gamma(E)$  we conclude that  $\sum_{i=0}^{\infty} \langle k_t(\cdot, x) f^j, e^i \rangle_{L^2} \cdot e^i$  equals

$k_t(\cdot, x)$  and hence

$$\mathrm{Tr}(Be^{-tH}) = \sum_{j=1}^{\dim E_x} \int_{x \in M} \langle Bk_t(x, x)f^j, f^j \rangle \quad (5.28)$$

$$= \int_{x \in M} \mathrm{tr}(Bk_t(x, x)) \quad (5.29)$$

□

The grading operator  $\gamma_5$  is a bounded operator. On a Clifford bundle with Dirac operator  $D$  we have the generalized Laplacian  $D^2$ . We conclude that  $\gamma_5 e^{-tD^2}$  is traceclass and so  $\mathrm{Tr}(\gamma_5 e^{-tD^2})$  is well-defined. In the next section we will study the eigenvalues of the Dirac operator so that we can express  $\mathrm{Tr}(\gamma_5 e^{-tD^2})$  in terms of these eigenvalues.

## 5.2 Spectral theory of generalized Laplacians

In functional analysis there is a generalized notion of eigenvalues and it is called the spectrum. The spectrum of an operator  $A$  is the collection  $\lambda \in \mathbb{C}$  such that  $A - \lambda \mathrm{Id}$  is not invertible. There is a class of operators for which the spectrum behaves well, namely the class of compact<sup>13</sup> operators. We recall the following theorems from functional analysis. For more information see Murphy [1990].

**Theorem 5.11.** *On a Hilbert space, the space of finite rank operators lies dense in the space of compact operators.*

Using this theorem we can create compact operator by considering the limit of a converging sequence of finite rank operators. The next two theorems, we describe the spectrum of compact operators

**Theorem 5.12.** *If  $A$  is a compact operator on a Banach space  $\mathcal{B}$ , then the spectrum of  $A$  is at most countable, each non-zero element of the spectrum is an eigenvalue and all eigenvalues are isolated.*

<sup>13</sup>Do not confuse it with the notion of compact spaces!

Not all elements in the spectrum are eigenvalues, but all eigenvalues are part of the spectrum. The previous theorem states that for compact operators, the spectrum and the set of eigenvalues coincide (except for zero). Isolated means that eigenvalues does not lie infinitely close to each other.

**Theorem 5.13.** *If  $A$  is a compact self-adjoint operator on a Hilbert space  $\mathcal{H}$ , then  $\mathcal{H}$  can be decomposed in a countable family of finite eigenspaces.*

We show that Hilbert-Schmidt operators are compact. Especially we get that smoothing operators of heat kernels are compact. Using the eigenspace decomposition of we then calculate the spectrum of a generalized Laplacian.

**Lemma 5.14.** *Hilbert-Schmidt operators are compact.*

*Proof.* Let  $K: \mathcal{H} \rightarrow \mathcal{H}$  be a Hilbert-Schmidt operator on a Hilbert space  $\mathcal{H}$ . Let  $\{e^i\}$  be an orthonormal basis of  $\mathcal{H}$ . Denote  $K^N$  as the projection of  $K$  to the span of  $\{e^1, \dots, e^N\}$ . That is  $K^N = \sum_{i=1}^N K(e^i)e_i^b$  where  $b$  denotes the dual. We estimate the difference of  $K$  and  $K^N$  in the operator norm:

$$\|K - K^N\|_{op} \leq \sum_{i=N+1}^{\infty} \|(Ke^i)e_i^b\|_{op} \leq \sum_{i=N+1}^{\infty} \|Ke^i\|$$

This is arbitrary small when  $N$  tends to infinity, because  $K$  is Hilbert-Schmidt. Hence,  $K$  is a limit of finite rank operators and thus compact.  $\square$

Smoothing operators  $e^{-tH}$  of heat kernels are compact operators. Hence, for each  $t \in \mathbb{R}_+$ , we have a eigenvalue decomposition of  $\Gamma_{L^2}(E)$  with respect to  $e^{-tH}$ . If we defined  $e^{-tH}$  using functional calculus, we get that  $\lambda$  is an eigenvalue of  $H$  if and only if  $e^{-t\lambda}$  is an eigenvalue of  $e^{-tH}$ . We replicate this result using the heat equation. First we relate the different eigenvalue decompositions of  $e^{-tH}$ . That is, we show that there only one eigenvalue decomposition for  $e^{-tH}$  and the rest is related using the exponential map.

**Lemma 5.15.** *Let  $E$  be a vector bundle with a positive definite inner product over a compact Riemannian manifold  $(M, g)$  and let  $H$  be a generalized Laplacian. Let  $k_t^H$  be the heat kernel and let  $e^{-tH}$  be the corresponding smoothing operator. The operator  $e^{-tH}$  has only positive eigenvalues and its kernel is empty.*

*Proof.* Let  $\lambda$  be an eigenvalue of  $e^{-tH}$  and let  $\mathcal{H}_\lambda$  be the eigenspace of  $e^{-tH}$  corresponding to the eigenvalue  $\lambda$ . Because  $e^{-tH}$  is self-adjoint, we have that  $\lambda$  is real. Let  $z \in \mathcal{H}_\lambda$ . From the semi-group property of the heat kernel follows that

$$e^{-tH}(e^{-t/2 H} z) = e^{-\frac{3}{2}t H} = e^{-t/2 H} e^{-tH} z = \lambda(e^{-t/2 H} z). \quad (5.30)$$

This shows  $e^{-t/2 H}(\mathcal{H}_\lambda)$  is a subset of  $\mathcal{H}_\lambda$ . From the spectral theory of compact operators it follows that

$$\mathcal{H}_\lambda = \bigoplus_i \mathcal{H}_{\lambda, \mu_i} \quad (5.31)$$

where  $\mathcal{H}_{\lambda, \mu_i}$  is the eigenspace of  $e^{-t/2 H}$  with eigenvalue  $\mu_i$  and the eigenspace of  $e^{-tH}$  with eigenvalue  $\lambda$ . Also  $\mu_i$  is real valued, because  $e^{-t/2 H}|_{\mathcal{H}_\lambda}$  is self-adjoint. So for all  $z \in \mathcal{H}_{\lambda, \mu_i}$  we get

$$\lambda z = e^{-tH} z = (e^{-t/2 H})^2 z = \mu_i^2 z \quad (5.32)$$

We conclude that the eigenvalues of  $e^{-tH}$  are non-negative.

If  $e^{-tH}$  has a non-empty kernel, then we can construct a sequence  $\{t_i\}$  converging to zero such that  $e^{-t_i H} z = 0$  for all  $t_i$ . Hence the limit of  $e^{-tH} z$  tends to zero when  $t$  tends to zero. This is in conflict with the boundary condition of the heat kernel. Therefore, the kernel is empty.  $\square$

**Theorem 5.16.** *Let  $E$  be a vector bundle with a positive definite inner product over a compact Riemannian manifold  $(M, g)$  and let  $H$  be a generalized Laplacian. Let  $k_t^H$  be the heat kernel and let  $e^{-tH}$  be the corresponding smoothing operator. The space  $\Gamma_{L^2}(E)$  can be decomposed in an countable family of finite eigenspaces  $\mathcal{H}_\lambda$  such that*

$$e^{-tH}|_{\mathcal{H}_\lambda} = \lambda^t \cdot \text{Id}$$

*Proof.* Let  $\lambda$  be an eigenvalue of  $e^{-H}$  and let  $\mathcal{H}_\lambda$  be the eigenspace of  $e^{-H}$  corresponding to the eigenvalue  $\lambda$ . Fix  $n, m \in \mathbb{N}$ . The operators  $e^{-H}$  and  $e^{-\frac{n}{m}H}$  commute and so  $e^{-\frac{n}{m}H}(\mathcal{H}_\lambda) \subseteq \mathcal{H}_\lambda$ . From the spectral theory of compact operators it follows that

$$\mathcal{H}_\lambda = \bigoplus_i \mathcal{H}_{\lambda, \mu_i} \quad (5.33)$$

where  $\mathcal{H}_{\lambda, \mu_i}$  is also the eigenspace of  $e^{-\frac{n}{m}H}$  with eigenvalue  $\mu_i$ . Let  $z \in \mathcal{H}_{\lambda, \mu_i}$ . By comparing eigenvalues we get

$$\mu_i^m z = (e^{-\frac{n}{m}H})^m z = e^{-nH} z = (e^{-H})^n z = \lambda^n z \quad (5.34)$$

Because  $\mu_i$  and  $\lambda$  are strictly larger than zero, we conclude that  $\mu_i = \lambda^{n/m}$  and this proves the theorem for all  $t \in \mathbb{Q}_+$ . Finally recall that  $e^{-tH}$  is continuous in  $t$  and thus the theorem holds for all  $t > 0$ .  $\square$

Finally, we study the eigenvalues of  $H$ . We use the heat equation to show that there is a one to one correspondence between the spectrum of  $e^{-tH}$  and  $H$ . The proof is straight forward, but we need that  $H$  and  $e^{-tH}$  commute. This is proven in the following lemma.

**Lemma 5.17.** *Let  $E$  be a vector bundle with a positive definite inner product over a compact Riemannian manifold  $(M, g)$  and let  $H$  be a generalized Laplacian. Let  $k_t^H$  be the heat kernel and let  $e^{-tH}$  be the corresponding smoothing operator. The operators  $H$  and  $e^{-tH}$  commute for all  $t > 0$ .*

*Proof.* Because  $H$  and  $e^{-tH}$  are self-adjoint, we get

$$He^{-tH} - e^{-tH}H = He^{-tH} - (He^{-tH})^* \quad (5.35)$$

Using the heat equation, we rewrite this into

$$He^{-tH} - e^{-tH}H = -\frac{\partial}{\partial t}e^{-tH} + \left(\frac{\partial}{\partial t}e^{-tH}\right)^* = -\frac{\partial}{\partial t}(e^{-tH} - (e^{-tH})^*) \quad (5.36)$$

Because  $e^{-tH}$  is self-adjoint we conclude that  $H$  and  $e^{-tH}$  commute.  $\square$

**Theorem 5.18.** *Let  $E$  be a vector bundle with a positive definite inner product over a compact Riemannian manifold  $(M, g)$  and let  $H$  be a generalized Laplacian. Let  $k_t^H$  be the heat kernel and let  $e^{-tH}$  be the corresponding smoothing operator. A section  $s \in \Gamma_{L^2}(E)$  is an eigenvector of  $H$  with eigenvalue  $\lambda$  if and only if it is an eigenvector of  $e^{-H}$  with eigenvalue  $e^{-\lambda}$ . Therefore,  $H$  has countably many eigenvalues which are isolated.*

*Proof.* Let  $e^{-\lambda}$  be an eigenvalue of  $e^{-H}$ . Let  $s \in \Gamma_{L^2}(E)$  such that  $e^{-H}s = e^{-\lambda}s$ . From the heat equation and Theorem 5.16 it follows that

$$0 = \left(\frac{\partial}{\partial t} + H\right)e^{-tH}s = \left(\frac{\partial}{\partial t} + H\right)e^{-\lambda t}s = (-\lambda + H)e^{-\lambda t}s \quad (5.37)$$

After the limit  $t \rightarrow 0$ , one concludes that  $HS = \lambda s$ . Because  $\Gamma_{L^2}(E)$  can be fully decomposed by the eigenvalues of  $e^{-H}$ , we know that  $H$  has no other eigenvalues.  $\square$

### 5.3 The McKean-Singer formula

When we considered the Abelian anomaly using the Fujikawa method we showed that the anomaly was proportional to  $\text{Tr}(\gamma_5 e^{-tD^2})$  where  $\gamma_5$  is the grading operator and  $D$  is the Dirac operator. We already found that this is a well-defined quantity for Clifford bundles on compact Riemannian manifolds. We now study this trace using the eigenvalue decomposition we found in the previous section.

**Theorem 5.19** (McKean and Singer [1967]). *Let  $S$  be a Clifford bundle over a compact Riemannian manifold  $(M, g)$  and let  $D$  be the Dirac operator. Let  $k_t^H$  be the heat kernel of  $D^2$  and let  $e^{-tD^2}$  be the corresponding smoothing operator. Let  $\gamma_5$  be the grading operator and let  $D_\pm$  be the restriction of  $D$  to the  $\pm 1$  eigenspace of  $\gamma_5$ . Then,*

$$\text{Tr}(\gamma_5 e^{-tD^2}) = \dim \ker D_+ - \dim \ker D_- \quad (5.38)$$

*This result is called the **McKean-Singer formula***

*Proof.* By Theorem 5.18 we can decompose  $\Gamma(S)$  into the eigenspaces

$$\Gamma(S) = \bigoplus_i H_{\lambda_i} \quad (5.39)$$

where  $H_{\lambda_i}$  is the  $\lambda_i \mathbb{R}$  eigenspace of  $D$ . Denote  $\{s_j^{\lambda_i}\}$  as an orthonormal basis of  $H_{\lambda_i}$ . The trace of  $\gamma_5 e^{-tD^2}$  equals

$$\text{Tr}(\gamma_5 e^{-tD^2}) = \sum_i \sum_j \langle s_j^{\lambda_i}, \gamma_5 e^{-tD^2} s_j^{\lambda_i} \rangle = \sum_i e^{-t\lambda_i^2} \sum_j \langle s_j^{\lambda_i}, \gamma_5 s_j^{\lambda_i} \rangle \quad (5.40)$$

However,  $\langle s_j^{\lambda_i}, \gamma_5 s_j^{\lambda_i} \rangle$  will vanish if  $\lambda_i \neq 0$ . Indeed, the grading operator and the Dirac operator anti-commute and so for all  $i$  and  $j$  and so

$$D\gamma_5 s_j^{\lambda_i} = -\gamma_5 D s_j^{\lambda_i} = -\lambda_i \gamma_5 s_j^{\lambda_i} \quad (5.41)$$

This shows that  $\gamma_5 s_j^{\lambda_i}$  is an element of  $H_{-\lambda_i}$ . From the orthonormality of the eigenspaces it follows that  $\langle s_j^{\lambda_i}, \gamma_5 s_j^{\lambda_i} \rangle = 0$  if  $\lambda_i \neq 0$ . Therefore, the trace simplifies to

$$\text{Tr}(\gamma_5 e^{-tD^2}) = \sum_j \langle s_j^0, \gamma_5 s_j^0 \rangle \quad (5.42)$$

The grading operator decomposes the kernel of  $D$  into two eigenspaces  $\ker D_+ \oplus \ker D_-$ . Let  $\{s_j^\pm\}$  be an orthonormal basis of  $\ker D_\pm$ . We conclude that the trace of  $\gamma_5 e^{-tD^2}$  equals

$$\mathrm{Tr}(\gamma_5 e^{-tD^2}) = \sum_j \langle s_j^+, s_j^+ \rangle - \langle s_j^-, s_j^- \rangle = \dim \ker D_+ - \dim \ker D_-. \quad (5.43)$$

□

Notice that the adjoint of  $D_+$  is  $D_-$ . Therefore, we can write the McKean-Singer formula as  $\mathrm{Tr}(\epsilon e^{-tD^2}) = \dim \ker D_+ - \dim \mathrm{coker} D_+$ . This quantity is called the index of the operator  $D_+$ . When considering Dirac operators some writers (Roe [1998], Berline et al. [2004]) call this the index of  $D$ . Although they look similar, those definitions does not coincide. Indeed, recall that  $D$  is a formally self-adjoint operator. Hence the kernel equals the cokernel and  $\dim \ker D - \dim \mathrm{coker} D$  is zero. In this thesis we only calculate the index of Dirac operators and hence define the index as follows:

**Definition 5.20.** *Let  $S$  be a graded Clifford bundle over a compact Riemannian manifold  $(M, g)$ . Let  $D$  be the Dirac operator and let  $e^{-tD^2}$  be the smoothing operator for which the kernel is the heat kernel  $k_t^{D^2}$ . The **index of  $D$** , which we denote by  $\mathrm{Ind}(D)$ , is defined as*

$$\mathrm{Ind}(D) = \dim \ker D_+ - \dim \ker D_- \quad (5.44)$$

where  $D_\pm$  is the restriction of  $D$  on the  $\pm 1$  eigenspace of the grading operator.

One of the main results of the Atiyah-Singer index theorem is that the index of a Dirac operator is topologically determined. That is, it does not depend on the choice of metric or connection, but only depends on the topology of the vector bundle. A similar result can be deduced from the McKean-Singer formula.

**Corollary 5.21** (Roe [1998], Proposition 11.13). *Let  $D_t$  be a continuous family of graded Dirac operators. That is, Let  $g_t$  be a continuous family of Riemannian metrics on a compact manifold  $M$ ,  $\gamma_t$  be a continuous family of Clifford actions on a Clifford bundle  $E \rightarrow M$  and let  $\nabla_t$  be a continuous family of compatible connections on  $E$ .  $D_t$  is the resulting Dirac operator. Then the index of  $D_t$  is independent of  $t$ .*

*Proof.* The map  $(s, t) \mapsto \text{Tr}(\gamma_5 e^{-sD_t^2})$  is a continuous map. By the McKean-Singer formula this map equals  $(s, t) \mapsto \text{Index}(D_t)$ . Notice that the index is an integer. The only continuous maps from  $\mathbb{R}$  to  $\mathbb{Z}$  are the constant maps and hence  $\text{Index}(D_t)$  is constant.  $\square$

In theory the index of a Dirac operator  $D$  can be calculated using the formal solution of the heat equation. Indeed, we know that the index of  $D$  equals  $\int_{x \in M} \text{tr}[\gamma_5 k_t(x, x)] \text{Vol}(g)$  where  $k_t$  is the heat kernel. We approximate this using the formal solution of the heat equation  $k_t(x, y) = \frac{1}{(4\pi t)^{n/2}} e^{-d(x,y)^2/4t} \sum_i t^i \Phi_i(x, y)$ . The index is independent of  $t$  and hence

$$\text{Index}(D) = \begin{cases} \frac{1}{(4\pi)^{n/2}} \int_{x \in M} \text{tr}[\gamma_5 \Phi_{n/2}(x, x)] & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd.} \end{cases} \quad (5.45)$$

Although this calculation is straight forward, it is not easy to calculate  $\Phi_{n/2}$  by hand. To solve this equation we need to study the behavior of the heat kernel and the Clifford action with the trace. In the next section we study the interaction between the Clifford action and the trace.

## 5.4 The trace and the Clifford action

When we considered chiral anomalies using perturbation theory, we often used the 'trace identities'

$$\text{Tr}(\gamma_5) = \text{Tr}(\gamma_5 \gamma^\mu) = \text{Tr}(\gamma_5 \gamma^\mu \gamma^\nu) = \text{Tr}(\gamma_5 \gamma^\mu \gamma^\nu \gamma^\rho) = 0 \quad (5.46)$$

for indices  $\mu, \nu$  and  $\rho$  and

$$\text{Tr}(\gamma_5 \gamma^\mu \gamma^\nu \gamma^\rho \gamma^\sigma) = 4i\epsilon^{\mu\nu\rho\sigma} \quad (5.47)$$

where  $\epsilon^{\mu\nu\rho\sigma}$  is  $\pm 1$  depending on the permutation of  $\{\mu, \nu, \rho, \sigma\}$ . One can find these results by calculating the trace over the gamma matrices. However these identities can also be deduced from the Clifford algebra. We show this in the following lemma.

**Lemma 5.22** (Roe [1998], Lemma 11.5). *Let  $V$  be a  $2n$  dimensional vector space with a positive definite inner product  $g$ . Let  $\text{Cl}(V)$  be the Clifford algebra and let  $S$  be a left  $\text{Cl}(V)$ -module. Let  $\gamma_5$  be a grading operator, let  $F \in \text{End}_{\text{Cl}(V)}(S)$  and let  $\gamma: V \rightarrow \text{End}(S)$  be the Clifford action. For any orthonormal basis  $\{s_i\}$  of  $V$  there is*

$$\text{Tr}(\gamma_5 F) = 0 \quad (5.48)$$

$$\text{Tr}(\gamma_5 \gamma(s_0) \dots \gamma(s_k) F) = 0 \quad \forall k < 2n \quad (5.49)$$

If  $\gamma_5$  is the canonical grading operator then,

$$\mathrm{Tr}(\gamma_5 \gamma(s_0) \dots \gamma(s_{2n}) F) = (-i)^n \mathrm{Tr}(F) \quad (5.50)$$

*Proof.* Let  $v \in V$  with norm 1. By the anti-commutation property of the Clifford algebra it follows that  $\gamma(v)^2 = -\mathrm{Id}_S$  and hence  $\mathrm{Tr}(\gamma_5 F) = -\mathrm{Tr}(\gamma_5 \gamma(v) \gamma(v) F)$ . The Clifford action anti-commutes with the grading operator and so

$$\mathrm{Tr}(\gamma_5 F) = -\mathrm{Tr}(\gamma_5 \gamma(v) \gamma(v) F) = +\mathrm{Tr}(\gamma(v) \gamma_5 \gamma(v) F) = +\mathrm{Tr}(\gamma_5 \gamma(v) F \gamma(v)) \quad (5.51)$$

Because  $F$  commutes with the Clifford action it follows that  $\mathrm{Tr}(\gamma_5 F) = 0$ . This trick can be repeated for  $\mathrm{Tr}(\gamma_5 \gamma(s_0) \dots \gamma(s_{2k}) F)$  for all  $k < n$ . That is, we add the term  $-\gamma(s_{2k+1}) \gamma(s_{2k+1})$  and we cyclically permute. Hence,  $\mathrm{Tr}(\gamma_5 \gamma(s_0) \dots \gamma(s_{2k}) F) = 0$  for all  $k < n$ .

To show that  $\mathrm{Tr}(\gamma_5 \gamma(s_0) \dots \gamma(s_{2k+1}) F) = 0$  for all  $2k+1 \in \mathbb{N}$  we just pull  $\gamma_5$  through the other side of the trace. That is, we use that  $\gamma_5$  anti-commutes with the Clifford action and commutes with  $F$ . Hence,

$$\mathrm{Tr}(\gamma_5 \gamma(s_0) \dots \gamma(s_{2k+1}) F) = (-1)^{2k+1} \mathrm{Tr}(\gamma(s_0) \dots \gamma(s_{2k+1}) F \gamma_5) \quad (5.52)$$

By the cyclic property of the trace and  $(-1)^{2k+1} = -1$  we conclude

$$\mathrm{Tr}(\gamma_5 \gamma(s_0) \dots \gamma(s_{2k+1}) F) = 0. \quad (5.53)$$

Finally we assume that  $\gamma_5$  is the canonical grading operator and we calculate  $\mathrm{Tr}(\gamma_5 \cdot \gamma(s_0) \dots \gamma(s_{2n}) F)$ . Recall that  $\gamma_5 = i^n \gamma(s_0) \dots \gamma(s_{2n})$  and hence

$$\mathrm{Tr}(\gamma_5 \gamma(s_0) \dots \gamma(s_{2n}) F) = (-i^n) \mathrm{Tr}(\gamma_5^2 F) \quad (5.54)$$

From  $\gamma_5^2 = 1$  follows the result.  $\square$

With this lemma we characterize the trace for all elements in  $\mathrm{Cl}(V) \otimes \mathrm{End}_{\mathrm{Cl}(V)}(S)$ . Clearly these are endomorphisms on  $S$ . The next question is whether all endomorphisms on  $S$  can be decomposed into  $\mathrm{Cl}(V) \otimes \mathrm{End}_{\mathrm{Cl}(V)}(S)$ . That is, is every linear map the product of Clifford actions and a Clifford endomorphism? The following proposition shows that this is indeed true.

**Proposition 5.23.** *Let  $V$  be a  $2n$ -dimensional vector space with a positive definite inner product and let  $S$  be a left  $\text{Cl}(V)$ -module. Assume that the Clifford action is normal with respect to the norm on  $S$ . Then the map which injects  $\text{Cl}(V) \otimes \text{End}_{\text{Cl}(V)}(S)$  into  $\text{End}_{\mathbb{C}}(S)$  using the Clifford action is an isomorphism.*

*Proof.* Because elements of the Clifford algebra are invertible we have that  $\text{Cl}(V) \otimes \text{End}_{\text{Cl}(V)}(S) \rightarrow \text{End}_{\mathbb{C}}(S)$  is injective. We only need to show that this map is surjective. Let  $F \in \text{End}_{\mathbb{C}}(S)$  be a linear map on  $S$  and let  $v \in V$  be a vector of length one. We first show that  $F$  can be decomposed into two terms that (anti)-commute with  $\gamma(v)$ . Indeed, let  $P_v^\pm$  be bounded linear operators on  $S$  defined by  $P_v^\pm = \frac{1 \mp i \gamma(v)}{2}$ . These operators have the properties:

$$\begin{aligned} P_v^\pm \cdot P_v^\pm &= P_v^\pm & P_v^\pm \cdot P_v^\mp &= 0 \\ P_v^+ + P_v^- &= \text{Id}_S & P_v^\pm \gamma(v) &= \gamma(v) P_v^\pm = \pm i P_v^\pm \end{aligned} \quad (5.55)$$

Hence  $P_v^\pm$  are the orthogonal eigenspace projectors of  $\gamma(v)$ . From these properties it follows that

$$P_v^\pm F P_v^\pm \gamma(v) = \pm i P_v^\pm F P_v^\pm = +\gamma(v) P_v^\pm F P_v^\pm \quad (5.56)$$

$$P_v^\mp F P_v^\pm \gamma(v) = \pm i P_v^\mp F P_v^\pm = -\gamma(v) P_v^\mp F P_v^\pm. \quad (5.57)$$

and thus we decompose  $F$  into

$$F = (P_v^+ + P_v^-) F (P_v^+ + P_v^-) \quad (5.58)$$

$$= (P_v^+ F P_v^+ + P_v^- F P_v^-) + (P_v^+ F P_v^- + P_v^- F P_v^+) \quad (5.59)$$

We denote  $F_v^\pm$  for the operator that is defined by  $F_v^\pm = P_v^+ F P_v^\pm F + P_v^- F P_v^\mp$ . By equations 5.56 and 5.57 it follows that  $F_v^+$  commutes and  $F_v^-$  anti-commutes with  $\gamma(v)$ . Hence,  $F = F^+ + F^-$  is the sum of (anti)-commuting terms w.r.t  $\gamma(v)$ .

Assume w.l.o.g. that  $F$  (anti)-commutes with  $\gamma(v)$ . That is, suppose that  $F \gamma(v) = \epsilon \gamma(v) F$  for some  $\epsilon \in \{0, 1\}$ . Let  $w \in V$  be of length one and normal to  $v$ . By the anti-commutation property of the Clifford algebra it follows that  $P_w^\pm \gamma(v) = \gamma(v) P_w^\mp$ . However,  $F_w^\pm$  is defined such that  $F_w^\pm \gamma(v) = \epsilon \gamma(v) F_w^\pm$ . By induction we conclude that  $F$  can be decomposed into the sum of terms that (anti)-commute with a whole orthonormal basis  $\{e^\mu\}$  of  $V$ .

Now pick an orthonormal basis  $\{e_i\}$  of  $V$  and assume w.l.o.g. that  $F$  commutes with  $e_1, \dots, e_k$  and anti-commutes with  $e_{k+1}, \dots, e_{2n}$ . Let  $\tilde{F}^+, \tilde{F}^- \in \text{End}(S)$  be operators defined as  $\tilde{F}^+ = \gamma_1 \dots \gamma_k F$  and  $\tilde{F}^- = \gamma_{k+1} \dots \gamma_{2n} F$ . By the anti-commutation property of the Clifford algebra it follows that

$$\gamma_\mu \tilde{F}^+ = (-1)^{k-1} \gamma_1 \dots \gamma_k \gamma_\mu F = (-1)^{k-1} \tilde{F}^+ \gamma_\mu \quad \forall \mu \leq k \quad (5.60)$$

$$\gamma_\mu \tilde{F}^+ = (-1)^k \gamma_1 \dots \gamma_k \gamma_\mu F = (-1)^{k+1} \tilde{F}^+ \gamma_\mu \quad \forall \mu > k \quad (5.61)$$

$$\gamma_\mu \tilde{F}^- = (-1)^{2n-k} \gamma_{k+1} \dots \gamma_{2n} \gamma_\mu F = (-1)^{2n-k} \tilde{F}^- \gamma_\mu \quad \forall \mu \leq k \quad (5.62)$$

$$\gamma_\mu \tilde{F}^- = (-1)^{2n-k-1} \gamma_{k+1} \dots \gamma_{2n} \gamma_\mu F = (-1)^{2n-k} \tilde{F}^- \gamma_\mu \quad \forall \mu > k \quad (5.63)$$

This calculation shows that if  $k$  is even, then  $\tilde{F}^-$  is a Clifford endomorphism and if  $k$  is odd, then  $\tilde{F}^+$  is a Clifford bundle endomorphism. So in both cases,  $F$  has a representative in  $\text{Cl}(V) \otimes \text{End}_{\text{Cl}(V)}(S)$ . So  $\text{Cl}(V) \otimes \text{End}_{\text{Cl}(V)}(S)$  is isomorphic to  $\text{End}_{\mathbb{C}} S$ .  $\square$

In the next chapter we decompose the heat kernel  $k_t$  into  $\text{Cl}(V) \otimes \text{End}_{\text{Cl}(V)}(S)$  and we will calculate the index of a Dirac operator.

## 6 Characteristic classes

In this chapter we consider the following question: "How can we distinguish two different vector bundles?" The answer to this question is studied in the theory of characteristic classes. We introduce this theory by following the Chern-Weil method.

### 6.1 Invariant polynomials

**Definition 6.1.** Let  $M_n(\mathbb{C})$  be the algebra of  $n \times n$  complex matrices. An **invariant polynomial** on  $M_n(\mathbb{C})$  is a polynomial function

$$P: M_n(\mathbb{C}) \rightarrow \mathbb{C} \quad (6.1)$$

such that for all  $X, Y \in M_n(\mathbb{C})$

$$P(XY) = P(YX) \quad (6.2)$$

Clearly the trace is an invariant polynomial over  $\mathbb{C}$ . Another example of an invariant polynomial is the determinant. Recall that the determinant and the trace are independent on the choice of basis. We show that it is also true for invariant polynomials.

**Lemma 6.2.** Let  $V$  be an  $n$ -dimensional vector space and let  $P$  be a polynomial. Then  $P$  is an invariant polynomial if and only if it descends to a map from  $\text{End}(V)$  to  $\mathbb{C}$ .

*Proof.* Assume that  $P$  is invariant and let  $E$  be an endomorphism on  $V$ . Any choice of basis for  $V$  induces an isomorphism between  $\text{End}(V)$  and  $M_n(\mathbb{C})$ . Hence in a basis  $\{e^i\}$  the endomorphism  $E$  has a matrix representation  $M$ . If we consider another basis  $\{\tilde{e}^i\}$  of  $V$  there is a matrix  $S \in GL_n(\mathbb{C})$  that represents the basis transformation from  $\{e^i\}$  to  $\{\tilde{e}^i\}$ . In this new basis  $E$  is given by  $SMS^{-1}$ . Because  $P$  is invariant we have

$$P(SMS^{-1}) = P(MS^{-1}S) = P(M) \quad (6.3)$$

Therefore,  $P$  is independent of basis representation and  $P$  descends to a map from  $\text{End}(V)$  to  $\mathbb{C}$ .

Now assume that  $P$  is a polynomial which descends to a map from  $\text{End}(V)$  to  $\mathbb{C}$ . Let  $X, Y \in M_n(\mathbb{C})$  and pick a basis  $\{v^i\}$  of  $V$ . Then  $X$  and  $Y$  represents endomorphisms on  $V$ . Assume that  $Y$  is invertible and let  $M = XY$ . Viewing  $Y$  as a basis transformation we have

$$P(XY) = P(M) = P(YMY^{-1}) = P(YX) \quad (6.4)$$

Because  $GL_n(\mathbb{C})$  is dense in  $M_n(\mathbb{C})$  and polynomials are continuous we conclude that  $P(XY) = P(YX)$  for all  $X, Y \in M_n(\mathbb{C})$ .  $\square$

Formally an invariant polynomial  $P$  over a vector space  $V$  is an element of  $\text{Sym}^\bullet(\text{End}_{\mathbb{C}} V)^*$ . For all  $M \in \text{End}_{\mathbb{C}}(V)$  we consider the linear map  $\iota_M: (\text{End}_{\mathbb{C}} V)^* \rightarrow \mathbb{C}$  which is defined as  $\iota_M(v) = v(M)$ . Using the universal property of the symmetric algebra  $\iota_M$  extends to a unique algebra homomorphism  $\iota_M: \text{Sym}^\bullet(\text{End}_{\mathbb{C}} V)^* \rightarrow \mathbb{C}$ . By construction  $\iota_M(P)$  equals the polynomial  $P$  evaluated at  $M$ .

**Example 6.3.** Consider the vector space  $\mathbb{C}^2$  and let  $M$  be a complex  $2 \times 2$  matrix. The components of  $M$  we denote as  $M_{ij}$ . Let  $p_{ij}: M_2(\mathbb{C}) \rightarrow \mathbb{C}$  be the projection onto the  $(i, j)^{\text{th}}$  component. In this case the map  $\iota_M: M_2(\mathbb{C})^* \rightarrow \mathbb{C}$  equals  $\iota_M(p_{ij}) = M_{ij}$  for all  $i$  and  $j$ . We now consider the determinant which is formally the element  $p_{00} \otimes p_{11} - p_{01} \otimes p_{10}$  in  $\text{Sym}^\bullet M_2(\mathbb{C})^*$ . Calculating  $\iota_M(\det)$  we get

$$\iota_M(\det) = \iota_M(p_{00} \otimes p_{11} - p_{01} \otimes p_{10}) \quad (6.5)$$

$$= \iota_M(p_{00}) \cdot \iota_M(p_{11}) - \iota_M(p_{01}) \cdot \iota_M(p_{10}) \quad (6.6)$$

$$= M_{00}M_{11} - M_{01}M_{10} \quad (6.7)$$

Hence  $\iota_M(\det)$  equals the determinant of  $M$ .

Let  $A$  be a commutative algebra,  $V$  be a vector space and consider ~~matrices over~~  $V$  with values in  $A$ . So let  $M \in \text{End}_{\mathbb{C}}(V) \otimes A$  and define the map  $\iota_M: (\text{End}_{\mathbb{C}} V)^* \rightarrow A$  by  $\iota_M(v) = v(M)$ . Using the universal property of the symmetric algebra we extend  $\iota_M$  to a map from  $\text{Sym}^\bullet(\text{End}_{\mathbb{C}} V)^*$  to  $A$ . Informally we interpret  $\iota_M(P)$  as the polynomial  $P$  evaluated over  $M$ , but we replaced the multiplication over  $\mathbb{C}$  with the multiplication over  $A$ . Hence we write  $\iota_M(P)$  as  $P(M)$ .

**Example 6.4.** Again consider  $\mathbb{C}^2$  and let  $A$  be a commutative algebra. We denote the matrix multiplication with  $*$ . Let  $M$  be a matrix on  $\mathbb{R}^2$  with values in  $A$ . We want to interpret the determinant of  $M$  as

$$\det(M) = M_{00} * M_{11} - M_{01} * M_{10} \quad (6.8)$$

With the above construction this is indeed possible. Indeed, the map  $\iota_M$  is given by  $\iota_M(p_{ij}) = M_{ij} \in A$  for all  $i, j$ . We conclude that

$$\iota_M(\det) = \iota_M(p_{00} \otimes p_{11} - p_{01} \otimes p_{10}) \quad (6.9)$$

$$= \iota_M(p_{00}) * \iota_M(p_{11}) - \iota_M(p_{01}) * \iota_M(p_{10}) \quad (6.10)$$

$$= M_{00} * M_{11} - M_{01} * M_{10} \quad (6.11)$$

Hence we define the determinant of an element  $M \in M_2(A)$  as  $\iota_M(\det)$ .

Let  $E \rightarrow M$  be a  $k$ -dimensional complex vector bundle over an  $n$  dimensional manifold. We consider the vector bundle  $\text{End}(E) \otimes (\oplus_k \Lambda^{2k} T^*M)$  which we denote as  $\text{End}(E) \otimes \Lambda^{ev} T^*M$ . In a local trivialization, sections of  $\text{End}(E) \otimes \Lambda^{ev} T^*M$  are maps from  $M$  to  $\mathbb{R}^{k \times k} \otimes \Lambda^{ev} \mathbb{R}^n$ . The algebra  $\Lambda^{ev} \mathbb{R}^n$  is commutative and hence we can use the construction described before. Because invariant polynomials are independent on the choice of basis, they behave well on overlapping charts. Hence, it follows that invariant polynomials induce smooth maps from  $\Gamma(\text{End}_{\mathbb{C}}(E) \otimes \Lambda^{ev} T^*M)$  to  $\Gamma(\Lambda^{ev} T^*M)$ .

Let  $\nabla$  be a connection on a vector bundle  $E$  and let  $P$  be an invariant polynomial. The curvature  $K$  w.r.t  $\nabla$  is a section of  $\text{End}(S) \otimes \Lambda^2 T^*M$ . We show that  $P(K)$  is a useful measure to distinguish vector bundles. For this we need the curvature expressed in a local basis. These calculations are straight forward and are done in section 9.3. Here we just state the results.

**Definition 6.5.** Let  $E \rightarrow M$  be a  $k$  dimensional vector bundle with a positive definite inner product and let  $\{s^i\}$  be a local orthonormal frame of  $E$ . Let  $\nabla$  be a compatible connection on  $E$ . The **connection 1-form**  $\omega$  is an element of  $\Gamma(M_k(E) \otimes \Lambda^1 T^*M)$  defined by the components

$$\omega_{ij} = s^i \lrcorner \nabla s^j \quad (6.12)$$

**Lemma 6.6.** Let  $E \rightarrow M$  be a vector bundle and let  $\{s^i\}$  be a local frame of  $E$ . Let  $\nabla$  be a connection on  $E$  and let  $\omega$  be the connection 1-form. Equip  $\Gamma(\text{End}(E)) \otimes \Omega^\bullet(M)$  with the multiplication induced by the matrix multiplication on  $\text{End}(E)$  and the wedge product on  $\Omega^\bullet(M)$ . The curvature  $K$  is a matrix on  $E$  with values in  $\Omega^2(M)$  and its components are given by

$$K = d\omega - \omega^2 \quad (6.13)$$

**Lemma 6.7** (Second Bianchi identity). *Let  $E \rightarrow M$  be a vector bundle and let  $\{s^i\}$  be a local frame of  $E$ . Let  $\nabla$  be a connection on  $E$  and let  $\omega$  be the connection 1-form. Equip  $\Gamma(\text{End}(E)) \otimes \Omega^\bullet(M)$  with the multiplication induced by the matrix multiplication on  $\text{End}(E)$  and the wedge product on  $\Omega^\bullet(M)$ . Then the exterior derivative of  $K$  equals*

$$dK = \omega K - K\omega \quad (6.14)$$

**Example 6.8.** For the Riemann curvature  $R$  on a Riemannian manifold  $M$  the second Bianchi identity equals  $dR = 0$ . Indeed, for all  $x \in M$  we can pick the Riemannian normal coordinate system centered at  $x$ . It has the property that  $\nabla \frac{\partial}{\partial x^\mu} = 0$  for all  $\frac{\partial}{\partial x^\mu}$ . Hence, the connection 1-form at  $x$  is zero and so  $dR = 0$  at  $x$ . Because we can pick  $x \in M$  randomly we get that  $dR = 0$  everywhere.

In general the exterior derivative of a curvature  $K$  is not zero. However,  $P(K)$  is closed if  $P$  is an invariant polynomial. We show this in the next lemmas, but for this we need to define the directional derivative of an invariant polynomial.

**Definition 6.9.** *Let  $A$  be an algebra, let  $M_n(A)$  be the matrix algebra of complex  $n \times n$  matrices with values in  $A$  and let  $P: M_n(A) \rightarrow A$  be an invariant polynomial. The directional derivative  $P'$  of  $P$  is a map from  $M_n(A)$  to  $M_n(A)^* \otimes A$  that is given by*

$$P'(M)(N) = \lim_{t \rightarrow 0} \frac{1}{t} [P(M + tN) - P(M)] \quad (6.15)$$

for all  $M, N \in M_n(A)$ .

**Lemma 6.10** (Milnor and Stasheff [1974], Page 297). *Let  $E \rightarrow M$  be a vector bundle and let  $\{s^i\}$  be a local frame of  $E$ . Let  $K$  be a section of  $\text{End}(E) \otimes \Lambda^{ev} T^*M$  and let  $P: \Gamma(\text{End}(E) \otimes \Lambda^{ev} T^*M) \rightarrow \Gamma(\Lambda^{ev} T^*M)$  be an invariant polynomial. Then the exterior derivative of  $P(K)$  equals*

$$dP(K) = P'(K)(dK) \quad (6.16)$$

where  $P'$  is the directional derivative of  $P$ .

*Proof.* We show this by induction. Assume that  $P$  is constant. Then  $P(K)$  is constant and so  $dP(K)$  vanishes. At the same time  $P'(K)$  is zero and so  $dP(K) = P'(K)(dK)$ .

Now assume that for some invariant polynomial  $Q$  that  $dQ(K) = Q'(K)(dK)$ . Let  $v \in \Gamma(\text{End}^*(E))$  be a locally constant section. That is,  $dv(K) = v(dK)$  and consider the polynomial  $P = v \cdot Q$ . The exterior derivative of  $P(K)$  is  $dP(K) = dv(K) \wedge Q(K) + v(K) \wedge dQ(K)$ . By assumption this equals

$$dP(K) = v(dK) \wedge Q(K) + v(K) \wedge Q'(K)(dK) \quad (6.17)$$

Now we calculate  $P'(K)(dK)$ . From the definition of the directional derivative it follows that

$$P'(K)(dK) = \lim_{t \rightarrow 0} \frac{1}{t} [v(K + t \cdot dK) \wedge Q(K + t \cdot dK) - v(K) \wedge Q(K)] \quad (6.18)$$

Recall that  $v$  is a linear map and so  $P'(K)(dK)$  simplifies into

$$\begin{aligned} P'(K)(dK) = & v(K) \wedge \left( \lim_{t \rightarrow 0} \frac{1}{t} [Q(K + t \cdot dK) - Q(K)] \right) + \\ & + \lim_{t \rightarrow 0} v(dK) \wedge Q(K + t \cdot dK) \end{aligned} \quad (6.19)$$

We conclude that  $P'(K)(dK) = v(K) \wedge Q'(K)(dK) + v(dK) \wedge Q(K)$  and so  $dP(K) = P'(K)(dK)$ . By induction we conclude the statement.  $\square$

**Theorem 6.11** (Milnor and Stasheff [1974], Page 296). *Let  $E \rightarrow M$  be a vector bundle, let  $\nabla$  be a connection on  $E$  and let  $K$  be its curvature. For every invariant polynomial  $P$ , the differential form  $P(K)$  is closed.*

*Proof.* Let  $\omega$  be the connection 1-form. Because  $P$  is an invariant polynomial we have

$$P((1 + t\omega)K) = P(K(1 + t\omega)) \quad (6.20)$$

Taking the derivative of  $t$  and set  $t = 0$  we conclude

$$P'(K)(\omega K) = P'(K)(K\omega) \quad (6.21)$$

By linearity of  $P'(K)$  we have  $P'(K)(\omega K - K\omega) = 0$  and by the second Bianchi identity it follows that  $P'(K)(dK)$  vanishes. Using the previous lemma we conclude that  $P(K)$  is closed.  $\square$

**Theorem 6.12** (Milnor and Stasheff [1974], Page 298). *Let  $E \rightarrow M$  be a vector bundle and let  $\nabla$  be a connection on  $E$ . Let  $K$  be its curvature and let  $P$  be an invariant polynomial. Then the cohomology class of  $P(K)$  does not depend on the choice of  $\nabla$ .*

*Proof.* Let  $\pi: M \times \mathbb{R} \rightarrow M$  be defined as  $\pi(x, t) = x$  and let for all  $t \in \mathbb{R}$ ,  $\iota_t: M \rightarrow M \times \mathbb{R}$  be defined as  $\iota_t(x) = (x, t)$ . Let  $\pi^*E$  and  $\iota_t^*\pi^*E$  be the pull-back bundles. Because  $\pi \circ \iota$  is the identity map we have that the pull-back bundle  $\iota_t^*\pi^*E \rightarrow M$  is equal to  $E \rightarrow M$ . Now consider the following diagram

$$\begin{array}{ccccc} E \cong \iota_t^*\pi^*E & & \pi^*E & & E \\ \downarrow & & \downarrow & & \downarrow \\ M & \xrightarrow{\iota_t} & M \times \mathbb{R} & \xrightarrow{\pi} & M \end{array} \quad (6.22)$$

Let  $\nabla^0$  and  $\nabla^1$  be two connections on  $E \rightarrow M$ . Using the pull-back<sup>14</sup> we get two connections  $\pi^*\nabla^0$  and  $\pi^*\nabla^1$  on  $\pi^*E$ . Also the linear combination

$$\nabla' = t\pi^*\nabla^1 + (1-t)\pi^*\nabla^0 \quad (6.23)$$

is a connection on  $\pi^*E$ . We pull-back  $\nabla'$  on  $\iota_t^*\pi^*E$  and we calculate  $\iota_0^*\pi^*\nabla'$ . Let  $s \in \Gamma(E)$  and note that  $s = \iota_t^*\pi^*s$  for all  $t \in \mathbb{R}$ . Hence for all  $u \in \Gamma(T^*M)$  we have

$$(\iota_0^*\nabla')_u s = (\iota_0^*\nabla')_u (\iota_0^*\pi^*s) = \iota_0^*(\nabla'_{d\iota_t(u)} \pi^*s) \quad (6.24)$$

Note that for  $\iota_0^*(\nabla'_{d\iota_t(u)} \pi^*s)$  we only consider the fibers of  $(\pi^*E)_{(\cdot, 0)}$  and so  $\iota_0^*(\nabla'_{d\iota_t(u)} \pi^*s)$  equals  $\iota_0^*((\pi^*\nabla^0)_{d\iota_t(u)} \pi^*s)$ . By the definition of the pull-back connection  $\pi^*\nabla^0$  ( $\iota_0^*\nabla'$ )<sub>u</sub>  $s$  simplifies to

$$(\iota_0^*\nabla')_u s = \iota_0^*\pi^*\nabla^0_{d\pi \circ d\iota_t(u)} s = \nabla^0 s \quad (6.25)$$

Using a similar argument we can show that  $\iota_1^*\nabla' = \nabla^1$ .

Let  $K^0$ ,  $K^1$  and  $K'$  be the curvature with respect to the curvatures  $\nabla^0$ ,  $\nabla^1$  and  $\nabla'$ . Note that  $\iota_t$  is a homotopy between  $\iota_0$  and  $\iota_1$ . Hence the induced maps  $\iota_0^*, \iota_1^*: H^\bullet(M \times \mathbb{R}) \rightarrow H^\bullet(M)$  are equal. Therefore, we have

$$[P(K^0)] = \iota_0^*[P(K')] = \iota_1^*[P(K')] = [P(K^1)] \in H^\bullet(M) \quad (6.26)$$

So the cohomology class of  $P(K)$  does not depend on the choice of connection.  $\square$

<sup>14</sup>Recall that for a vector bundle  $E \rightarrow M$  and a smooth map between manifolds  $\phi: N \rightarrow M$  the pull-back connection is the unique connection on  $\phi^*E$  such that  $(\phi^*\nabla)_X(\phi^*s) = \phi^*(\nabla_{d\phi(X)}s)$  for all  $X \in T^*N$  and  $s \in \Gamma(\phi^*E)$

Given two vector bundles  $E^0 \rightarrow M$  and  $E^1 \rightarrow M$  of same dimension we can distinguish them using an invariant polynomial  $P$ . Indeed, we can pick two connections  $\nabla^0$  on  $E^0$  and  $\nabla^1$  on  $E^1$  and this induces two curvature tensors  $K^0$  and  $K^1$ . By Theorem 6.11 we know that  $P(K^0)$  and  $P(K^1)$  represents two cohomology classes in  $H^\bullet(M)$  and by theorem 6.12 we know that these cohomology classes does not depend on the choice of  $\nabla^0$  and  $\nabla^1$ . Hence, if  $P(K^0)$  and  $p(K^1)$  represent different cohomology classes, we know that they poses different topologies. We call the cohomology classes of  $P(K^0)$  and  $P(K^1)$  **characteristic classes**.

## 6.2 Examples of characteristic classes

We now revisit some examples of characteristic classes. Let  $E \rightarrow M$  be a complex vector bundle and let  $P$  be an invariant polynomial. If  $f$  is a holomorphic map on  $\mathbb{C}$  then  $f$  has a power series. So the composition  $P \circ f$  is an formal power series. Because  $\Lambda^{ev} T^*M$  is nilpotent the power series of  $P \circ f$  only contains finitely many non-zero terms and so it is an invariant polynomial.

**Example 6.13.** Let  $E \rightarrow M$  be a complex vector bundle, let  $K$  be a curvature tensor on  $E$  and let  $f$  be a holomorphic map on  $\mathbb{C}$ . The **Chern  $f$ -genus** of  $E$  is the complex cohomology class of  $\Pi_f(K)$  where  $\Pi_k$  is the invariant polynomial

$$\Pi_k(X) = \det \left[ f \left( \frac{-X}{2\pi i} \right) \right] \quad (6.27)$$

**Example 6.14.** Let  $E \rightarrow M$  be a complex vector bundle. The **total Chern class**  $c(E) \in H^{ev}(M)$  is the Chern  $f$ -genus with respect to the holomorphic map  $f(z) = 1 + z$ .

The  $i^{\text{th}}$  **Chern class** is the projection of the total Chern class into  $H^{2i}(M) \otimes \mathbb{C}$ .

**Example 6.15.** Let  $E \rightarrow M$  be a complex vector bundle and let  $K$  be a curvature tensor on  $E$ . The **Chern character**  $\text{Ch}(E) \in H^{ev}(M) \otimes \mathbb{C}$  is the cohomology class of  $P(K)$  where  $P$  is the invariant polynomial

$$P(X) = \text{tr} \left[ \exp \left( \frac{-X}{2\pi i} \right) \right] \quad (6.28)$$

For the Atiyah-Singer index theorem we mainly focus on the Chern Character and the Chern  $f$ -genus of  $TM$  where  $f(z) = \sqrt{\frac{z/2}{\sinh(z/2)}}$ . The latter is a cohomology class in  $H^\bullet(M)$  and not in  $H^\bullet(M) \otimes \mathbb{C}$ . Indeed, if  $R$  is the curvature tensor on  $TM$ , then  $R$  is a skew-symmetric matrix. Hence, it has eigenvalues of the form  $\pm i\lambda$  where  $\lambda \in \Omega^2(M)$ . Hence, the Chern  $f$ -genus is

$$\Pi_f(TM \otimes \mathbb{C}) = \prod_j \sqrt{\frac{\lambda_j/2}{\sinh \lambda_j/2}} \cdot \sqrt{\frac{-\lambda_j/2}{\sinh -\lambda_j/2}} = \prod_j \frac{\lambda_j/2}{\sinh \lambda_j/2} \quad (6.29)$$

This shows that  $\Pi_f(TM \otimes \mathbb{C})$  is indeed a representative of  $H^\bullet(M)$ . Also it does not depend on the choice of branch of the square root. This is not a special property of  $TM$  and so we define:

**Example 6.16.** Let  $E \rightarrow M$  be a real vector bundle. The  $\hat{A}$ -genus  $\hat{A}(E)$  is the Chern  $f$ -genus of  $E \otimes \mathbb{C}$  where  $f$  is the holomorphic map  $z \mapsto \sqrt{\frac{z/2}{\sinh z/2}}$ .

### 6.3 Characteristic classes of Clifford bundles

When we consider a Clifford bundle  $S \rightarrow (M, g)$  the curvature  $K$  is not the only section of  $\text{End}(S) \otimes \Lambda^{ev} T^*M$  we study. Namely, for a Clifford bundle we also have the twisting curvature  $F^S$  and the Riemann endomorphism  $R^S$ . We study if invariant polynomials over  $F^S$  and  $R^S$  also define characteristic classes.

**Lemma 6.17.** Let  $A$  be an algebra and let  $V, W$  be two vector spaces. Let  $P$  be an element of  $\text{Sym}^\bullet(V \oplus W)^*$ . Let  $\tilde{P}$  be the projection of  $P$  onto  $\text{Sym}^\bullet W^*$ . Then for all  $w \in W \otimes A$ ,  $v \in V \otimes A$  the polynomials  $P$  and  $\tilde{P}$  satisfies

$$P(w) = \tilde{P}(v + w) \quad (6.30)$$

*Proof.* Assume w.l.o.g. that  $P$  has a unique representation as  $\sum_i \prod_{j=0}^i (f_{ij} + g_{ij})$  where  $f_{ij} \in V^*$  and  $g_{ij} \in W^*$  for all  $i$  and  $j$ . Then  $\tilde{p}$  is the sum  $\sum_i \prod_{j=0}^i g_{ij}$ . Let

$v \in V \otimes A$  and  $w \in W \otimes A$ . Because  $f_{ij}(w) = 0$  and  $g_{ij}(v) = 0$  there is

$$P(w) = \sum_i \prod_{j=0}^i (f_{ij}(w) + g_{ij}(w)) = \sum_i \prod_{j=0}^i g_{ij}(w) \quad (6.31)$$

$$\tilde{P}(v + w) = \sum_i \prod_{j=0}^i (g_{ij}(v) + g_{ij}(w)) = \sum_i \prod_{j=0}^i g_{ij}(w) \quad (6.32)$$

This concludes that  $P(w) = \tilde{P}(v + w)$ .  $\square$

**Theorem 6.18.** *Let  $S \rightarrow (M, g)$  be a Clifford bundle, let  $R^S$  be the Riemann endomorphism and let  $F^S$  be the twisting curvature. For all invariant polynomials  $P$ , the differential forms  $P(R^S)$  and  $P(F^S)$  are closed.*

*Proof.* Let  $K \in \Gamma(\text{End}(S) \otimes \Lambda^2 T^*M)$  be the curvature on  $S$ . Recall that  $\text{End}_{\mathbb{C}}(S)$  is isomorphic to  $\text{Cl}(TM) \otimes \text{End}_{\text{Cl}}(S)$  where  $\text{Cl}(TM)$  is the Clifford action and  $\text{End}_{\text{Cl}}(S)$  is the bundle of Clifford endomorphisms. Because  $\text{Cl}(TM)$  is unital we can decompose  $\text{End}(S)$  into  $V \oplus W$  where  $W = \{\mathbb{C} \cdot \text{Id}\} \otimes \text{End}_{\text{Cl}}(S)$  and  $V$  is the rest. Let  $\tilde{P}$  be the projection of  $P$  onto  $W$ . By the previous lemma we know that  $P(F^S) = \tilde{P}(F^S + R^S) = \tilde{P}(K)$ . Notice that  $\tilde{P}$  is also an invariant polynomial. Hence,  $d\tilde{P}(K) = dP(F^S) = 0$ . Using the same argument we can show that  $dP(R^S) = 0$ .  $\square$

For Clifford bundles we can also use  $F^S$  and  $R^S$  to calculate characteristic classes. For the Atiyah-Singer theorem we only use the following:

**Definition 6.19.** *The relative Chern character  $\text{Ch}^{rel}(S) \in H^{ev}(M) \otimes \mathbb{C}$  of a Clifford bundle  $S \rightarrow M$  on an  $n$  dimensional manifold is the cohomology class of  $P(F^S)$  where  $P$  is the invariant polynomial*

$$P(X) = (-2)^{n/2} \text{tr} \left[ \exp \left( \frac{-X}{2\pi i} \right) \right] \quad (6.33)$$

## 7 Symbol Calculus

In chapter 3 we saw how the Fujikawa method relates the chiral anomaly to the index of a Dirac operator. In one of the steps we considered a Taylor approximation and we showed that only a single order contributed. We formalize this approach by introducing graded and filtered algebras. We show that the Taylor approximation used by Fujikawa is related to the Getzler filtration. Secondly we investigate how the heat equation changes the Getzler filtration. The result is related to the heat equation for Mehlers kernel and this proves the Atiyah-Singer index theorem.

### 7.1 Definitions

**Definition 7.1.** A **graded algebra** is an algebra  $G$  with a direct sum decomposition  $G = \bigoplus_{i \in \mathbb{Z}} G_i$  such that  $G_i \cdot G_j \subseteq G_{i+j}$  for all  $i, j \in \mathbb{Z}$ . An element  $g \in G$  is of **degree**  $k$  if  $g \in G_k$

**Example 7.2** (Trivial grading). We can turn any algebra  $G$  into a graded algebra. For this let  $G_0 = G$  and  $G_i = \{0\}$  if  $i \neq 0$ . Clearly  $G$  has the decomposition  $\bigoplus_i G_i$ . Because  $G$  is an algebra  $G_0 \cdot G_0 \subseteq G_0$ . Also  $\{0\} \cdot G = G \cdot \{0\} = \{0\}$  and hence  $G_i \cdot G_j \subseteq G_{i+j}$ . We call this the trivial grading.

**Example 7.3** (Taylor series). Let  $V$  be a vector space let  $G = \mathbb{C}[[V]]$  be the space of formal power series of maps on  $V$ . Homogeneous polynomials span this space and it even induces a direct sum decomposition. Indeed, we say that a formal series is of degree<sup>15</sup>  $-k$  and only if it is a homogeneous polynomial of degree  $k$ . Under change of coordinates a homogeneous polynomial stays homogeneous and the degree stays the same. Hence the homogeneous polynomials induce a direct sum decomposition  $G$ . Also the multiplication between a degree  $i$  and  $j$  homogeneous polynomial induces a degree  $i + j$  homogeneous polynomial. Hence the property  $G_i \cdot G_j \subseteq G_{i+j}$  is also satisfied.

**Example 7.4** (Exterior algebra). Let  $V$  be a vector space and let  $G_k = \Lambda^k V$  be the  $k$ -times wedge product of elements in  $V$ . The exterior algebra is the direct sum  $G = \bigoplus_{k \in \mathbb{N}} \Lambda^k V$ . The wedge product between an  $i$ - and  $j$ -form is a  $(i + j)$ -form. Hence  $G_i \wedge G_j \subseteq G_{i+j}$  and the exterior algebra is a graded algebra.

<sup>15</sup>Although the minus sign is not necessarily, it will be useful in later calculations.

Let  $V$  be a vector space with inner product  $g$  and let  $\text{Cl}_k(V, g)$  be the linear span of  $k$  products of pairwise orthonormal Clifford actions. The space  $\text{Cl}(V, g)$  has the direct sum decomposition  $\bigoplus_k \text{Cl}_k(V, g)$  but is not a graded algebra. Indeed, for all  $v \in V$  the Clifford action  $\gamma(v)$  is an element of  $\text{Cl}_1(V, g)$ . However  $\gamma(v)^2$  is an element of  $\text{Cl}_0(V, g)$  instead of  $\text{Cl}_2(V, g)$ , because  $\gamma(v)^2 = -g(v, v) \cdot \text{Id}$ . To assign degrees for elements in the Clifford algebra we use filtered algebras:

**Definition 7.5.** A **filtered algebra** is an algebra  $A$  with a family of subspaces  $A_i$ ,  $i \in \mathbb{Z}$  such that  $A_i \subseteq A_{i+1}$  and  $A_i \cdot A_j \subseteq A_{i+j}$ . An element  $a \in A$  is of **degree**  $k \in \mathbb{Z}$  if  $a \in A_k$ , but  $a \notin A_{k-1}$ .

**Example 7.6.** (Clifford algebra) Consider the Clifford algebra  $A = \text{Cl}(V, g)$  and let  $A_i = \bigoplus_{j \leq i} \text{Cl}_j(V, g)$ . Clearly  $A$  is the union of all  $A_i$  and for all  $i$  and  $j$  the product of an element in  $A_i$  and in  $A_j$  is an element of  $A_{i+j}$ . Hence the Clifford algebra is a filtered algebra.

**Example 7.7.** (Differential operators) Recall that for a manifold  $M$  a differential operator is a smooth operator on  $C^\infty(M)$  such that in a local coordinate frame  $\{x^\mu\}$  the differential operator is given by

$$\sum c_{\mu_1, \dots, \mu_k} \frac{\partial}{\partial x^{\mu_1}} \frac{\partial}{\partial x^{\mu_2}} \cdots \frac{\partial}{\partial x^{\mu_k}} \quad (7.1)$$

Here  $c_{\mu_1, \dots, \mu_k}$  is a smooth maps on  $M$ . If we use multi-index notation we say that a differential must locally be of the form  $\sum_I c_I \frac{\partial}{\partial x^I}$ . Let  $|I|$  be the degree of the multi index  $I$  and let  $A_k$  be the vector space generated by the differential operators that are locally of the form  $\sum_{|I| \leq k} c_I \frac{\partial}{\partial x^I}$ . Notice that  $A_k$  is independent of choice of basis and  $A_i \cdot A_j \subseteq A_{i+j}$  for all  $i, j$ . Hence  $A_k$  induce a filtration on the space of differential operators. We denote differential operators as  $\mathcal{D}(M)$ .

**Lemma 7.8.** Let  $G = \bigoplus_k G_k$  be a graded algebra and let  $A_k = \bigoplus_{j \leq k} G_j$ . If there exists an  $M \in \mathbb{Z}$  such that  $G_k = \{0\}$  for all  $k > M$ , then  $\bigcup_k A_k$  is a filtration of  $G$ .

*Proof.* Clearly  $G$  is the union of all  $A_k$  and by definition  $A_k \subseteq A_{k+1}$  for all  $k$ . Because  $G$  is graded we conclude that  $A_i \cdot A_j \subseteq A_{i+j}$ . Hence  $A_i$  is a filtration of  $G$ .  $\square$

From the previous lemma we conclude that ~~all~~ our examples of graded algebras are also filtered algebras. Not every filtered algebra is a graded algebra. However for every filtered algebra we can construct a graded algebra by considering quotient spaces.

**Definition 7.9.** Let  $A$  be a filtered algebra. The **associated graded algebra** is the graded algebra  $G(A)$  that is defined by

$$G(A) = \bigoplus_k A_k / A_{k-1} \quad (7.2)$$

Given a filtered algebra  $A = \bigcup_i A_i$  consider the associated graded algebra  $G(A)$ . Notice that the quotient map  $\pi_k: A_k \rightarrow A_k / A_{k-1}$  describes the relation between the filtration of  $A$  and the grading of  $G(A)$ . Also notice that for all  $a \in A_i$  and  $a' \in A_j$

$$\pi_i(a) \cdot \pi_j(a') = \pi_{i+j}(a \cdot a'). \quad (7.3)$$

This is an example of a symbol map.

**Definition 7.10.** Let  $A = \bigcup_{k \in I} A_k$  be a filtered algebra and let  $G = \bigoplus_{k \in I} G_k$  be a graded algebra. A **symbol map** is a family of linear maps  $\sigma_k: A_k \rightarrow G_k$  such that

1. for all  $k \in I$  and  $a \in A_{k-1}$  the map  $\sigma_k$  satisfies  $\sigma_k(a) = 0$ .
2. for all  $i, j \in I$ ,  $a \in A_i$  and  $a' \in A_j$  the symbol map satisfies  $\sigma_i(a) \cdot \sigma_j(a') = \sigma_{i+j}(a \cdot a')$ .

If  $G$  is the associated graded algebra of  $A$ , then the **associated symbol map** is the family of projection maps  $\pi_k: A_k \rightarrow A_k / A_{k-1}$ , which is a symbol map between  $A$  and  $G(A)$ .

**Example 7.11.** Consider the trivially filtered algebra  $A$ . That is,  $A_i = A$  if  $i \geq 0$  and  $A_i = \{0\}$  else. The associated graded algebra  $G(A)$  is isomorphic to the trivially graded algebra of  $A$ . Hence, the associated symbol map equals  $\sigma_k = \text{Id}_A$  if  $k = 0$  and  $\sigma_k = 0$  else.

**Example 7.12.** (Clifford algebra) Let  $V$  be a vector space with symmetric 2-form  $g$ . Consider the associated graded algebra of  $\text{Cl}(V, g)$ . By the definition of a Clifford algebra the projection map  $\pi_i$  satisfies

$$\pi_1(v_1) \cdot \pi_1(v_2) + \pi_1(v_2) \cdot \pi_1(v_1) = \pi_2(v_1 \cdot v_2 + v_2 \cdot v_1) = -g(v_1, v_2) \cdot \pi_2(\text{Id}) = 0 \quad (7.4)$$

for all  $v_1, v_2 \in V$ . This shows that the associated graded algebra of  $\text{Cl}(V, g)$  is isomorphic to the exterior algebra  $\Lambda^\bullet V^*$ . Equation 7.4 also shows that the associated symbol map interchanges the Clifford multiplication with the wedge product. That is, for all  $v_1 \dots v_k \in V$ , the map  $\pi_k(v_1 \dots v_k)$  equals  $v_1^b \wedge \dots \wedge v_k^b$ .

**Example 7.13** (Endomorphisms of algebras). Given an filtered algebra  $A = \bigcup_i A_i$  we can create a filtered subalgebra on  $\text{End}(A)$ . Indeed,  $\text{End}_{\sigma_n}(A)$  be the subspace of  $\text{End}(A)$  such that for all  $k \in \mathbb{Z}$  the map  $f|_{A_k}$  maps into  $A_{k+n}$ . Because  $A_{k+n}$  is a subspace of  $A_{k+n+1}$  we conclude that  $\text{End}_{\sigma_n}(A) \subseteq \text{End}_{\sigma_{n+1}}(A)$ . Using the composition of maps we notice that  $\text{End}_{\sigma_n}(A) \circ \text{End}_{\sigma_m}(A) \subseteq \text{End}_{\sigma_{n+m}}(A)$ . Hence,  $\bigcup_n \text{End}_{\sigma_n}(A)$  is a filtered algebra and a subalgebra of  $\text{End}(A)$ . This property can also be shown for graded algebras  $G$ . That is, the subalgebra  $\bigoplus_n \text{End}_{\sigma_n}(G)$  of  $\text{End}(G)$  is also a graded algebra.

We construct a symbol map between  $\bigcup_n \text{End}_{\sigma_n}(A)$  and  $\bigoplus_n \text{End}_{\sigma_n}(G(A))$ . Fix  $n \in \mathbb{Z}$  and let  $f \in \text{End}_{\sigma_n}(A)$ . For any  $k \in \mathbb{Z}$  and  $a \in A_{k-1}$  we have  $f(a) \in A_{k+n-1}$ . This shows that  $\pi_{n+k} \circ f(a) = 0$ . We conclude that  $\pi_{n+k} \circ f$  factors over  $A_k/A_{k-1}$  and we can define  $\sigma_n(f): G(A) \rightarrow G(A)$  as the unique map such that for all  $k \in \mathbb{Z}$  the following diagram commutes

$$\begin{array}{ccc} A_k & \xrightarrow{f} & A_{k+n} \\ \downarrow \pi_k & & \downarrow \pi_{k+n} \\ A_k/A_{k-1} & \xrightarrow{\sigma_n(f)} & A_{k+n}/A_{k+n-1} \end{array}$$

We see that  $\sigma_n$  is a family of linear maps from  $\text{End}_{\sigma_n}(A)$  to  $\text{End}_{\sigma_n}(G(A))$ . It is also a symbol map. We already showed that  $\sigma_n(\text{End}_{\sigma_{n-1}}(A)) = 0$  for all  $n \in \mathbb{Z}$ . To show the homomorphism property let  $n, m \in \mathbb{Z}$ ,  $f \in \text{End}_{\sigma_n}(A)$  and let  $g \in \text{End}_{\sigma_m}(A)$ . For all  $k \in \mathbb{Z}$  we get the commuting diagram

$$\begin{array}{ccccc} A_k & \xrightarrow{f} & A_{k+n} & \xrightarrow{g} & A_{k+n+m} \\ \downarrow \pi_k & & \downarrow \pi_{k+n} & & \downarrow \pi_{k+n+m} \\ A_k/A_{k-1} & \xrightarrow{\sigma_n(f)} & A_{k+n}/A_{k+n-1} & \xrightarrow{\sigma_m(g)} & A_{k+n+m}/A_{k+n+m-1} \\ & & \searrow \sigma_{n+m}(g \circ f) & & \end{array}$$

By uniqueness we conclude that  $\sigma_{n+m}(g \circ f) = \sigma_m(g) \circ \sigma_n(f)$  and so  $\sigma$  is a symbol map.

**Example 7.14** (Taylor series). Consider formal power series  $\mathbb{C}[[\mathbb{R}]]$  over  $\mathbb{R}$ . Note that for all  $\alpha_i, \beta_j \in \mathbb{C}$  the differential operator  $\sum_{i=0}^n \alpha_i \left(\frac{\partial}{\partial x}\right)^i$  maps the formal power series  $\sum_{j=m}^{\infty} \beta_j x^j$  to  $\sum_{i=0}^n \sum_{j=m}^{\infty} \alpha_i \beta_j \cdot \frac{j!}{i!} x^{j-i}$ . Hence, the space  $\mathcal{D}_n(\mathbb{R})$  is a subalgebra of  $\text{End}_{\sigma_n}(\mathbb{C}[[\mathbb{R}]])$  for all  $n \in \mathbb{Z}$ . However the filtrations of  $\mathcal{D}_n(\mathbb{R})$  and  $\text{End}_{\sigma_n}(\mathbb{C}[[\mathbb{R}]])$  differs. To see this note that the operator  $p \mapsto x \cdot p$  is a differential operator of degree zero, but this operator is an element of  $\text{End}_{\sigma_{-1}}(\mathbb{C}[[\mathbb{R}]])$ .

## 7.2 Getzler filtration

In paragraph 3.1 we calculated the trace over  $\gamma_5 e^{-tD^2}$  by considering a Taylor approximation in  $t^{1/2}$  for equation 3.18.

$$\int \frac{d^4 k}{(2\pi)^4} t^{-2} \text{tr} \left[ \gamma_5 f \left( -(t^{1/2} \nabla_\mu + i k_\mu)^2 - \frac{it}{4} [\gamma^\mu, \gamma^\nu] F_{\mu\nu} \right) \right] \quad (3.18)$$

Notice that we have a factor  $t^{1/2}$  for every  $k_\mu$  and for every  $\gamma^\mu$ . Now we rephrase this in terms of filtrations. The element  $k_\mu$  is the Fourier transform for a degree one differential operator. Also,  $\gamma^\mu$  is a degree one Clifford action. By applying a Taylor approximation we combine the filtration of the Clifford action with the filtration **induced by differential operators**. In general we always can combine multiple filtrations using the tensor algebra.

**Lemma 7.15.** *Let  $B = \bigcup_i B_i$  and  $C = \bigcup_i C_i$  be filtered algebras and let  $A$  be the tensor algebra of  $B$  and  $C$ . Then  $A$  is a filtered algebra spanned by  $A_k = \sum_{i=0}^k B_i \otimes C_{k-i}$ .*

*Proof.* Let  $b \in B$ ,  $c \in C$  and consider  $b \otimes c \in A$ . Because  $B$  and  $C$  are filtrations there exists an  $i, j \in \mathbb{Z}$  such that  $b \in B_i$  and  $c \in C_j$ . Hence,  $b \otimes c \in B_i \otimes C_j$ . The space  $B_i \otimes C_j$  is a subspace of  $\sum_{i=0}^{i+j} B_i \otimes C_j = \sum_{i=0}^{i+j} B_i \otimes C_{(i+j)-i}$  which equals  $A_{i+j}$ . Hence,  $A$  is the union of the family of spaces  $A_k$ .

Now let  $k \in \mathbb{Z}$  and consider  $A_k$ . By definition it equals  $\sum_{i=0}^k B_i \otimes C_{k-i}$ . The space  $A_k$  is a subspace of  $A_{k+1}$ , because

$$A_k = \sum_{i=0}^k B_i \otimes C_{k-i} \subseteq \sum_{i=0}^k B_i \otimes C_{(k+1)-i} \subseteq \sum_{i=0}^{k+1} B_i \otimes C_{(k+1)-i} = A_{k+1} \quad (7.5)$$

Finally let  $i, j, k, l \in \mathbb{Z}$  and let  $b_i \in B_i$ ,  $c_j \in C_j$ ,  $b_k \in B_k$  and  $c_l \in C_l$ . Then  $b_i \otimes c_j \in A_{i+j}$  and  $b_k \otimes c_l \in A_{k+l}$ . The tensor product between  $b_i \otimes c_j$  and  $b_k \otimes c_l$  equals  $(b_i \cdot b_j) \otimes (c_k \cdot c_l)$ . Because  $B$  and  $C$  are filtrations  $(b_i \cdot b_j) \otimes (c_k \cdot c_l)$  is an element of  $B_{i+j} \otimes C_{k+l}$ . This proves that  $b_i \otimes c_j \cdot b_k \otimes c_l \in A_{i+j+k+l}$  and so  $A_{i+j} \cdot A_{j+k} \in A_{i+j+k+l}$ . We conclude that  $A$  is a filtered algebra.  $\square$

We want to create a filtration on the space of smoothing operators. For this let  $(M, g)$  be an even dimensional compact Riemannian manifold and let  $y \in M$ . Consider a neighborhood  $U_y \subseteq M$  such that  $\exp_y^{-1}: U_y \rightarrow T_y M$  is a chart of  $M$ . Let  $S \rightarrow U_y$

be a Clifford bundle and let  $p \in \Gamma(S \boxtimes S)$  be a kernel. We fix the second component of  $p$  at  $y$  and so we only consider  $p(\cdot, y) \in \Gamma(S \otimes S_y^*)$ . Using parallel transport and the exponential map we trivialize  $S$  into  $T_y M \times S_y$ . Then locally  $p(\cdot, y)$  is a smooth map from  $T_y M$  to  $S_y \otimes S_y^*$ . Notice that  $S_y \otimes S_y^*$  is isomorphic to  $\text{End}(S_y)$ . By Lemma 5.23 the space  $S_y \otimes S_y^*$  is isomorphic to  $\text{Cl}(T_y M) \otimes \text{End}_{\text{Cl}}(S_y)$  where  $\text{End}_{\text{Cl}}(S_y)$  is the space of Clifford endomorphisms on  $S_y$ . We conclude that  $p(\cdot, y)$  is locally a map from  $T_y M$  to  $\text{Cl}(T_y M) \otimes \text{End}_{\text{Cl}}(S_y)$ . That is,  $p(\cdot, y)$  is an element of  $C^\infty(T_y M) \otimes \text{Cl}(T_y M) \otimes \text{End}_{\text{Cl}}(S_y)$ . Notice that this space is the tensor product of filtered algebras. Using Lemma 7.15 we can create a filtration on  $\Gamma(S \otimes S_y^*)$ .

**Definition 7.16.** The tensor **filtration** induced from the Taylor **filtration** on  $C^\infty(T_y M)$ , the Clifford **filtration** on  $\text{Cl}(T_y M)$  and the trivial **filtration** on  $\text{End}_{\text{Cl}}(S_y)$  is the **Getzler filtration** for kernels on  $S$ . For operators on  $\Gamma(S \otimes S_y^*)$  we define the **Getzler filtration** for operators on  $\Gamma(S)$  and the **Getzler symbol** as the **filtration** and symbol described in Example 7.13.

In the rest of this section we refer to the trivialization  $T_y M \times \text{End}_{\mathbb{C}}(S_y)$  of  $S$  as the trivialization induced by parallel transport. Without further introducing we also use  $\{s_k\}$  for an orthonormal frame of  $T_y M$  and  $\{\tilde{s}_k\}$  as the local frame of  $S$  induced from  $\{s_k\}$  by parallel transport. Note that  $\{\tilde{s}_k\}$  is also orthonormal, because the connection on  $S$  is metric compatible.

Our goal is to calculate the Getzler symbol of the operator  $D^2$ . For this we need to calculate the symbol of the Clifford action and the covariant derivative. We show this in the next lemmas.

**Lemma 7.17.** Let  $v_y \in T_y M$  and consider the vector field  $v \in \Gamma(TM)$  induced by parallel transport. The Clifford action  $\gamma(v)$  along  $v$  is locally given by  $\text{Id}_{T_y M} \otimes \gamma(v_y) \otimes \text{Id}_{S_y} \in C^\infty(T_y M) \otimes \text{Cl}(T_y M) \otimes \text{End}_{\text{Cl}}(S_y)$ . The Clifford action  $\gamma(v)$  has Getzler degree one and the Getzler symbol  $\sigma_1(\gamma(v))$  equals  $v_y^\flat$ .

*Proof.* Let  $x \in M$  and consider a radial path  $p: [0, 1] \rightarrow T_y M$  from  $y$  to  $x$ . The local trivialization induced by parallel transport of  $\gamma(v)$  is given as

$$\sum_{i,j} \langle \tilde{s}_i, \gamma(v) \tilde{s}_j \rangle_x \otimes s_i \cdot s_j^\flat \in C^\infty(T_y M) \otimes \text{End}_{\mathbb{C}}(S_y) \quad (7.6)$$

The metric on  $S$  is compatible with the covariant derivative  $\nabla$  and so

$$\frac{\partial}{\partial t} \langle \tilde{s}_i, \gamma(v) \tilde{s}_j \rangle_{p(t)} = \langle \nabla_{\dot{p}(t)} \tilde{s}_i, \gamma(v) \tilde{s}_j \rangle_{p(t)} + \langle \tilde{s}_i, \nabla_{\dot{p}(t)} \gamma(v) \tilde{s}_j \rangle_{p(t)} \quad (7.7)$$

Because  $\tilde{s}_i$  is a vector field induced by parallel transport, we have the identity  $\nabla_{\dot{p}(t)} \tilde{s}_i = 0$  for all  $t$ . So equation 7.7 simplifies to

$$\frac{\partial}{\partial t} \langle \tilde{s}_i, \gamma(v) \tilde{s}_j \rangle_{p(t)} = \langle \tilde{s}_i, \nabla_{\dot{p}(t)} \gamma(v) \tilde{s}_j \rangle_{p(t)} \quad (7.8)$$

$$= \langle \tilde{s}_i, \gamma(\nabla_{\dot{p}(t)} v) \tilde{s}_j \rangle_{p(t)} + \langle \tilde{s}_i, \gamma(v) \nabla_{\dot{p}(t)} \tilde{s}_j \rangle_{p(t)} \quad (7.9)$$

$$= \langle \tilde{s}_i, \gamma(\nabla_{\dot{p}(t)} v) \tilde{s}_j \rangle_{p(t)} \quad (7.10)$$

Also the vector field  $v$  is induced from parallel transport and so  $\langle \tilde{s}_i, \gamma(v) \tilde{s}_j \rangle_{p(t)}$  is constant. We calculate this constant by evaluating at  $y$  and we conclude that  $\gamma(v)$  locally equals

$$\sum_{i,j} \langle s_i, \gamma(v_y) s_j \rangle \cdot \text{Id}_{U_y} \otimes s_i \cdot s_j^\flat \in C^\infty(T_y M) \otimes \text{End}_{\mathbb{C}}(S_y) \quad (7.11)$$

This simplifies to  $\text{Id}_{T_y M} \otimes \gamma(v_y) \in C^\infty(T_y M) \otimes \text{End}_{\mathbb{C}}(S_y)$  and we conclude the result.  $\square$

**Lemma 7.18.** *Any endomorphism  $F \in \Gamma(\text{End}_{\mathbb{C}}(S))$  is a Clifford endomorphism if and only if there exists some  $\alpha_i \in C^\infty(T_y M)$  and  $f^i \in \text{End}_{\text{Cl}}(S_y)$  such that in the trivialization induced by parallel transport  $F$  equals*

$$\sum_i \alpha_i \otimes \text{Id} \otimes f^i \in C^\infty(T_y M) \otimes \text{Cl}(T_y M) \otimes \text{End}_{\text{Cl}}(S_y) \quad (7.12)$$

*The Clifford bundle endomorphism  $F$  have Getzler degree zero and the Getzler symbol  $\sigma_0(F)$  equals  $F(y)$ .*

*Proof.* Let  $f \in \text{End}_{\text{Cl}}(S_y)$  and let  $\alpha \in C^\infty(T_y M)$ . Let  $F$  be a section of  $\text{End}_{\mathbb{C}}(S)$  such that  $F$  is locally given by  $\alpha \otimes \text{Id} \otimes f \in C^\infty(T_y M) \otimes \text{Cl}(T_y M) \otimes \text{End}_{\text{Cl}}(S_y)$ . Using the orthonormality of  $\{\tilde{s}_k\}$  we see that  $F$  equals

$$F = \sum_{ij} \alpha \cdot \langle s_i, f s_j \rangle \tilde{s}_i \cdot \tilde{s}_j^\flat \quad (7.13)$$

Let  $v \in \Gamma(TM)$  be the parallel transport of  $v_y \in T_y M$ . By the identity  $\sum_k \tilde{s}_k \tilde{s}_k^b = \text{Id}$  we conclude

$$F\gamma(v) = \sum_{ijk} \alpha \cdot \langle s_i, f s_j \rangle \langle \tilde{s}_j, \gamma(v) \tilde{s}_k \rangle \tilde{s}_i \cdot \tilde{s}_k^b \quad (7.14)$$

By Lemma 7.17 we know that  $\langle \tilde{s}_j, \gamma(v) \tilde{s}_k \rangle$  is constant and hence equal to  $\langle s_j, \gamma(v_y) s_k \rangle$ . This simplifies  $F\gamma(v)$  into

$$F\gamma(v) = \sum_{ijk} \alpha \cdot \langle s_i, f \gamma(v_y) s_k \rangle \tilde{s}_i \cdot \tilde{s}_k^b \quad (7.15)$$

By definition  $f$  must commute with  $\gamma(v_y)$  and so  $F$  commutes with  $\gamma(v)$ . Hence  $F$  is a Clifford bundle endomorphism.

At last consider the subbundle of  $\text{End}(S)$  spanned by the Clifford action and all Clifford endomorphisms that are locally of the form  $\sum_i \alpha_i \otimes \text{Id} \otimes f^i \in C^\infty(T_y M) \otimes \text{Cl}(T_y M) \otimes \text{End}_{\text{Cl}}(S_y)$ . By dimension counting we conclude it spans the whole bundle and so all Clifford bundle endomorphisms are of the form  $\sum_i \alpha_i \otimes \text{Id} \otimes f^i \in C^\infty(U_y) \otimes \text{Cl}(T_y M) \otimes \text{End}_{\text{Cl}}(S_y)$ .  $\square$

**Lemma 7.19** (Roe [1998], Proposition 12.22). *Let  $\{x^\mu\}$  be an orthonormal coordinate frame of  $T_y M$  and extend this to an orthonormal coordinate frame on  $TM$  using parallel transport. Let  $R$  be the Riemann curvature tensor. Then for all  $\mu$  the covariant derivative  $\nabla_\mu$  is a differential operator of Getzler degree one and the Getzler symbol of  $\nabla_\mu$  equals*

$$\sigma_1(\nabla_\mu) = \frac{\partial}{\partial x^\mu} + \frac{1}{4} \sum_\nu \left\langle \frac{\partial}{\partial x^\mu}, R(\cdot, \cdot) \frac{\partial}{\partial x^\nu} \right\rangle x^\nu \quad (7.16)$$

*Proof.* Let  $p$  be a section of  $S \otimes S_y^*$  and assume without loss of generality that  $p = \sum_{ij} p_{ij} \tilde{s}_i \cdot s_j^b$  where  $p_{ij} \in C^\infty(U_y)$ . In the trivialization induced by the parallel transport  $p$  equals  $\sum_{ij} p_{ij} \cdot s_i \cdot s_j^b$ . We show that in this trivialization  $\nabla_\mu p$  equals  $\sigma_1(\nabla_\mu) p$  plus lower order terms. From the Leibniz rule it follows that

$$\nabla_\mu p = \sum_{ij} \nabla_\mu (p_{ij} \tilde{s}_i \cdot s_j^b) = \sum_{ij} \frac{\partial p_{ij}}{\partial x^\mu} \tilde{s}_i \cdot s_j^b + \sum_{ijk} \langle \tilde{s}_k, \nabla_\mu \tilde{s}_i \rangle \cdot p_{ij} \tilde{s}_k \cdot s_j^b \quad (7.17)$$

Locally we can write  $\nabla_\mu p$  as  $(\frac{\partial}{\partial x^\mu} + \omega)p$  where  $\omega$  is a linear endomorphism on  $S_y$  with the elements  $\omega_{ki} = \langle \tilde{s}_k, \nabla_\mu \tilde{s}_i \rangle \in C^\infty(T_y M)$ . Clearly  $\frac{\partial}{\partial x^\mu}$  is a degree one operator and so we only need to show that  $\omega$  has Getzler degree one. For this we use a trick. Let  $K$  be the curvature tensor w.r.t. the connection  $\nabla$  of  $S$  and let  $\mathcal{R}$  be the radial vector field. That is,  $\mathcal{R} = \sum_\nu x^\nu \frac{\partial}{\partial x^\nu}$ . Note that the Lie bracket between  $\frac{\partial}{\partial x^\mu}$  and  $\mathcal{R}$  equals  $\frac{\partial}{\partial x^\mu}$ . So the matrix element  $\omega_{ki}$  equals  $\langle \tilde{s}_k, \nabla_{[\frac{\partial}{\partial x^\mu}, \mathcal{R}] \tilde{s}_i} \rangle$ . We write this in terms of the curvature.

$$\omega_{ki} = \langle \tilde{s}_k, \nabla_\mu \tilde{s}_i \rangle = \left\langle \tilde{s}_k, \left( -K \left( \frac{\partial}{\partial x^\mu}, \mathcal{R} \right) + \nabla_\mu \nabla_{\mathcal{R}} - \nabla_{\mathcal{R}} \nabla_\mu \right) \tilde{s}_i \right\rangle \quad (7.18)$$

By the definition of the parallel transport it follows that  $\nabla_{\mathcal{R}} \tilde{s}_i = 0$ . Using that  $\nabla$  is a metric connection equation 7.18 simplifies into

$$\langle \tilde{s}_k, \nabla_\mu \tilde{s}_i \rangle = - \left\langle \tilde{s}_k, K \left( \frac{\partial}{\partial x^\mu}, \mathcal{R} \right) \tilde{s}_i \right\rangle - \mathcal{L}_{\mathcal{R}} \langle \tilde{s}_k, \nabla_\mu \tilde{s}_i \rangle \quad (7.19)$$

Moving  $\mathcal{L}_{\mathcal{R}}$  to the left hand side, we get  $(1 + \mathcal{L}_{\mathcal{R}})\omega = -K \left( \frac{\partial}{\partial x^\mu}, \mathcal{R} \right)$ . The Weitzenböck formula states that  $K = R^S + F^S$  where  $F^S$  is a Clifford endomorphism and  $R^S = \frac{1}{4} \sum_{\nu\rho} \gamma_\nu \gamma_\rho \langle R(\cdot, \cdot) \frac{\partial}{\partial x^\nu}, \frac{\partial}{\partial x^\rho} \rangle$ . So the matrix  $\omega$  satisfies

$$\begin{aligned} (1 + \mathcal{L}_{\mathcal{R}})\omega &= -R^S \left( \frac{\partial}{\partial x^\mu}, \mathcal{R} \right) - F^S \left( \frac{\partial}{\partial x^\mu}, \mathcal{R} \right) \\ &= \frac{1}{4} \sum_{\nu\rho\sigma} \gamma_\nu \gamma_\rho \left\langle \frac{\partial}{\partial x^\mu}, R \left( \frac{\partial}{\partial x^\nu}, \frac{\partial}{\partial x^\rho} \right) \frac{\partial}{\partial x^\sigma} \right\rangle x^\sigma - F^S \left( \frac{\partial}{\partial x^\mu}, \mathcal{R} \right) \end{aligned} \quad (7.20)$$

Now we write  $\omega$  in local coordinates. Let  $I, J$  be multi-indices over  $\{x^\mu\}$  with degree  $|I|$  and  $|J|$  and write  $\omega$  as  $\sum_{IJ} x^I \otimes c^J \otimes \omega_{IJ} \in C^\infty(T_y M) \otimes \text{Cl}(T_y M) \otimes \text{End}_{\text{Cl}}(S_y)$ . The left hand side of equation 7.20 equals

$$\begin{aligned} (1 + \mathcal{L}_{\mathcal{R}})\omega &= \sum_{IJ} \left( 1 + \sum_\tau x^\tau \frac{\partial}{\partial x^\tau} \right) x^I \otimes c^J \otimes \omega_{IJ} \\ &= \sum_{IJ} \left( 1 + \sum_\tau x^\tau \frac{I(\tau)}{x^\tau} \right) x^I \otimes c^J \otimes \omega_{IJ} \\ &= \sum_{IJ} (1 + |I|) x^I \otimes c^J \otimes \omega_{IJ} \end{aligned} \quad (7.21)$$

We compare this to the right hand side of equation 7.20. Because  $F^S$  is a Clifford endomorphism, we conclude that  $F^S \left( \frac{\partial}{\partial x^\mu}, \mathcal{R} \right)$  is a Getzler degree zero operator. Also,

$R^S\left(\frac{\partial}{\partial x^\mu}, \mathcal{R}\right)$  has Getzler degree one and so

$$\begin{aligned} & \sum_{IJ} (1 + |I|) x^I \otimes c^J \otimes \omega_{IJ} \\ &= \frac{1}{4} \sum_{\nu\rho\sigma} \gamma_\nu \gamma_\rho \left\langle \frac{\partial}{\partial x^\mu}, R\left(\frac{\partial}{\partial x^\nu}, \frac{\partial}{\partial x^\rho}\right) \frac{\partial}{\partial x^\sigma} \right\rangle x^\sigma + \text{lower order terms} \end{aligned} \quad (7.22)$$

We conclude for the top order that  $X^I = x^\sigma$ ,  $C^J = \gamma_\nu \gamma_\rho$  and  $\omega_{IJ} = \left\langle \frac{\partial}{\partial x^\mu}, R\left(\frac{\partial}{\partial x^\nu}, \frac{\partial}{\partial x^\rho}\right) \frac{\partial}{\partial x^\sigma} \right\rangle$ . Hence  $\omega$  has Getzler degree one and is given by

$$2\omega = \frac{1}{4} \sum_{\nu\rho\sigma} \gamma_\nu \gamma_\rho \left\langle \frac{\partial}{\partial x^\mu}, R\left(\frac{\partial}{\partial x^\nu}, \frac{\partial}{\partial x^\rho}\right) \frac{\partial}{\partial x^\sigma} \right\rangle x^\sigma + \text{lower order terms} \quad (7.23)$$

The Getzler symbol of  $\omega$  equals

$$\sigma_1(\omega) = \frac{1}{8} \sum_{\nu\rho\sigma} \left\langle \frac{\partial}{\partial x^\mu}, R\left(\frac{\partial}{\partial x^\nu}, \frac{\partial}{\partial x^\rho}\right) \frac{\partial}{\partial x^\sigma} \right\rangle x^\sigma \cdot dx^\nu \wedge dx^\rho \quad (7.24)$$

$$= \frac{1}{4} \sum_{\sigma} \left\langle \frac{\partial}{\partial x^\mu}, R(\cdot, \cdot) \frac{\partial}{\partial x^\sigma} \right\rangle x^\sigma \quad (7.25)$$

and we finish the proof.  $\square$

**Corollary 7.20** (Roe [1998], Example 12.16). *The Dirac operator  $D$  has Getzler degree two and the Getzler symbol of  $D$  equals*

$$\sigma_2(D) = \sum_{\mu} \frac{\partial}{\partial x^\mu} \cdot dx^\mu + \frac{1}{8} \sum_{\mu\nu\rho\sigma} \left\langle \frac{\partial}{\partial x^\mu}, R\left(\frac{\partial}{\partial x^\nu}, \frac{\partial}{\partial x^\rho}\right) \frac{\partial}{\partial x^\sigma} \right\rangle x^\sigma \cdot dx^\mu \wedge dx^\rho \wedge dx^\sigma \quad (7.26)$$

where  $\{x^\mu\}$  is the Riemannian normal coordinate frame.

*Proof.* Recall that in the Riemannian normal coordinate frame  $D = \sum_{\mu} \gamma_{\mu} \nabla_{\mu}$  and use Lemmas 7.17 and 7.19.  $\square$

**Proposition 7.21** (Roe [1998], Proposition 12.17). *Let  $S \rightarrow (M, g)$  be a Clifford bundle over a compact even dimensional Riemannian manifold and let  $D$  be the Dirac operator. Let  $R$  be the Riemann curvature and  $F^s$  be the twisting curvature. Let  $\{x^\mu\}$  be the Riemannian normal coordinate frame and denote*

$R_{\mu\nu} = \left\langle \frac{\partial}{\partial x_\mu}, R(\cdot, \cdot) \frac{\partial}{\partial x_\nu} \right\rangle \in \Omega^2(M)$ . The operator  $D^2$  has Getzler degree two and its symbol equals

$$\sigma_2(D^2) = - \sum_{\mu} \left( \frac{\partial}{\partial x^\mu} + \frac{1}{4} \sum_{\nu} R_{\mu\nu} x^\nu \right)^2 + F^S \quad (7.27)$$

*Proof.* Proposition 1.27 states that  $D^2 = \nabla^* \nabla + F^S + \frac{1}{4} \kappa$  where  $F^S$  is the Clifford contraction of the twisting curvature and  $\kappa$  is the scalar curvature. In Riemannian normal coordinates we write this identity as

$$D^2 = - \sum_{\mu} \nabla_{\mu} \nabla_{\mu} + \frac{1}{2} \sum_{\mu, \nu} \gamma_{\mu} \gamma_{\nu} F^S \left( \frac{\partial}{\partial x^\mu}, \frac{\partial}{\partial x^\nu} \right) + \frac{1}{4} \kappa \quad (7.28)$$

Recall that the twisting curvature is a Clifford endomorphism. Hence by Lemmas 7.17, 7.18 and 7.19 we conclude that  $D^2$  has Getzler degree 2 and the symbol is given by

$$\sigma_2(D^2) = - \sum_{\mu} \sigma_2(\nabla_{\mu}^2) + \frac{1}{2} \sum_{\mu, \nu} \sigma_2 \left( \gamma_{\mu} \gamma_{\nu} F^S \left( \frac{\partial}{\partial x^\mu}, \frac{\partial}{\partial x^\nu} \right) \right) + \sigma_2 \left( \frac{1}{4} \kappa \right) \quad (7.29)$$

The scalar curvature and  $F^S \left( \frac{\partial}{\partial x^\mu}, \frac{\partial}{\partial x^\nu} \right)$  are Clifford endomorphisms and hence it has at most degree zero. So we simplify equation 7.29 into

$$\sigma_2(D^2) = - \sum_{\mu} \sigma_1(\nabla_{\mu})^2 + \frac{1}{2} \sum_{\mu, \nu} \sigma_1(\gamma_{\mu}) \sigma_1(\gamma_{\nu}) \sigma_0 \left( F^S \left( \frac{\partial}{\partial x^\mu}, \frac{\partial}{\partial x^\nu} \right) \right) \quad (7.30)$$

$$= - \sum_{\mu} \sigma_1(\nabla_{\mu})^2 + \frac{1}{2} \sum_{\mu, \nu} F^S \left( \frac{\partial}{\partial x^\mu}, \frac{\partial}{\partial x^\nu} \right) dx^{\mu} \wedge dx^{\nu} \quad (7.31)$$

$$= - \sum_{\mu} \sigma_1(\nabla_{\mu})^2 + F^S \quad (7.32)$$

Using Lemma 7.19 we conclude the result.  $\square$

### 7.3 The symbol of the heat kernel

In chapter 4 we showed that each Dirac operator on a compact Riemannian manifold has a heat kernel  $k_t$ . Using the heat equation we calculate the Getzler symbol of the heat kernel.

**Proposition 7.22** (Roe [1998], Proposition 12.24). *Let  $S \rightarrow (M, g)$  be a Clifford bundle over an  $n$  dimensional compact Riemannian manifold and let  $D$  be the Dirac operator. Assume that  $n$  is even. Let  $k_t$  be the heat kernel w.r.t. the generalized Laplacian  $D^2$  and denote the formal power series of  $k_t$  as  $\frac{1}{(4\pi t)^{n/2}} e^{-r^2/4t} \sum_{i=0}^{\infty} t^i \Phi_i$ . Then for all  $i$  the kernel  $\Phi_i$  has at most Getzler degree  $2i$  and the symbol of  $\Phi_i$  satisfies*

$$r \frac{\partial}{\partial r} \sigma_0(\Phi_0) = 0 \quad (7.33)$$

$$\left( r \frac{\partial}{\partial r} + i \right) \sigma_{2i}(\Phi_i) = - \sigma_2(D^2) \sigma_{2i-2}(\Phi_{i-1}) \quad \forall i \in \mathbb{N} \quad (7.34)$$

*Proof.* Let  $y \in M$ . In Theorem 4.13 we showed that  $k_t$  has a formal solution to the heat equation by proving that  $\Phi_i$  is the unique solution of the differential equations

$$\left( \frac{r}{4g} \frac{\partial g}{\partial r} + \nabla_{\mathcal{R}} \right) \Phi_0(\cdot, y) = 0 \quad (4.66)$$

$$\left( i + \frac{r}{4g} \frac{\partial g}{\partial r} + \nabla_{\mathcal{R}} \right) \Phi_i(\cdot, y) = - D^2 \Phi_{i-1}(\cdot, y) \quad \forall i > 0 \quad (4.67)$$

Here  $r$  is the distance from  $y$ ,  $\mathcal{R} = r \partial/\partial r$  is the radial vector field originating in  $y$  and  $g$  is the determinant of the metric. Suppose that  $\Phi_i(\cdot, y)$  has Getzler degree  $k_i$  for some  $k_i \in \mathbb{N}$ . The symbol of  $\Phi_i$  equals

$$\sigma_{k_0} \left( \left( \frac{r}{4g} \frac{\partial g}{\partial r} + \nabla_{\mathcal{R}} \right) \Phi_0(\cdot, y) \right) = 0 \quad (7.35)$$

$$\sigma_{k_i} \left( \left( i + \frac{r}{4g} \frac{\partial g}{\partial r} + \nabla_{\mathcal{R}} \right) \Phi_i(\cdot, y) \right) = \sigma_{k_i}(-D^2 \Phi_{i-1}(\cdot, y)) \quad \forall i > 0 \quad (7.36)$$

Multiplication by  $i$  is a degree zero operator and  $\frac{r}{4g} \frac{\partial g}{\partial r}$  has degree -1. Also  $\nabla_{\mathcal{R}}$  is a degree zero operator. Indeed, let  $\{x^\mu\}$  be the Riemannian orthonormal coordinate and notice that  $\nabla_{\mathcal{R}} = \sum_{\mu} x^\mu \cdot \nabla_{\mu}$ . The symbol of  $\nabla_{\mathcal{R}}$  equals

$$\sigma_0(\nabla_{\mathcal{R}}) = \sum_{\mu} x^\mu \frac{\partial}{\partial x^\mu} + \frac{1}{4} \sum_{\nu} R_{\mu\nu} x^\mu x^\nu. \quad (7.37)$$

Using the anti-symmetry in  $R$  we simplify  $\sigma_0(\nabla_{\mathcal{R}})$  into  $x^\mu \frac{\partial}{\partial x^\mu}$  and this equals  $r \frac{\partial}{\partial r}$ .

Therefore the symbol of  $\Phi_k$  equals

$$r \frac{\partial}{\partial r} (\sigma_{k_0} \Phi_0(\cdot, y)) = 0 \quad (7.38)$$

$$\left( r \frac{\partial}{\partial r} + i \right) \sigma_{k_i} \Phi_i(\cdot, y) = -\sigma_2(D^2) \sigma_{k_i-2}(\Phi_{i-1}) \quad \forall i > 0 \quad (7.39)$$

Inductively we conclude that  $\Phi_i$  has Getzler degree  $2i$ .  $\square$

Comparing equations 4.66 and 4.67 with 7.38 and 7.39 suggest we need to investigate the differential equation  $\frac{\partial}{\partial t} + \sigma_2(D^2) = 0$ . Even more, this is a heat equation. Indeed, it is sufficient to show that  $\sigma_2(D^2)$  is a generalized Laplacian. For this consider the trivial bundle  $T_y M \times \Lambda^\bullet T_y M \otimes \text{End}_{\text{Cl}}(S_y) \rightarrow T_y M$ . Equip  $\Lambda^\bullet T_y M$  with the induced metric from  $T_y M$  and equip  $\text{End}_{\text{Cl}}(S_y)$  with the Hilbert-Schmidt metric. The operator  $\frac{\partial}{\partial x^\mu} + \frac{1}{4} \sum_\nu R_{\mu\nu} x^\nu$  is a covariant derivative for all  $x^\mu$ . Denote this connection as  $\nabla$ . This connection is metric compatible, because  $R_{\mu\nu}$  is skew-symmetric. Hence the operator  $\sigma_2(D^2)$ , which equals  $\nabla^* \nabla + F^S$ , is a generalized Laplacian. Next we explicitly calculate the heat kernel.

**Proposition 7.23** (Roe [1998], Proposition 12.24). *Let  $S \rightarrow (M, g)$  be a Clifford bundle over an  $n$  dimensional compact Riemannian manifold. Let  $D$  be the Dirac operator and let  $k_t = \frac{1}{(4\pi t)^{n/2}} \exp(-r^2/4t) \sum_i t^i \Phi_i$  be the formal solution of the heat equation with respect to the generalized Laplacian  $D^2$ . Let  $y \in M$  and consider the trivial vector bundle  $T_y M \times \Lambda^\bullet T_y M \otimes \text{End}_{\text{Cl}}(S_y) \rightarrow T_y M$ . Let  $W_t = \frac{1}{(4\pi t)^{n/2}} \exp(-r^2/4t) \sum_i t^i \Theta_i$  be the formal solution of the heat equation with respect to the generalized Laplacian  $\sigma_2(D^2)$ . Then for all  $x \in T_y M$  and  $i \in \mathbb{N}$  the components  $\Theta_i$  satisfy*

$$\Theta_i(x, y) = \sigma_{2i}(\Phi_i) \quad (7.40)$$

*Proof.* In Theorem 4.13 we showed that  $\Theta_i$  is the solution of the differential equation

$$\left( \frac{r}{4g} \frac{\partial g}{\partial r} + \nabla_{\mathcal{R}} \right) \Phi_0(\cdot, y) = 0 \quad (4.66)$$

$$\left( i + \frac{r}{4g} \frac{\partial g}{\partial r} + \nabla_{\mathcal{R}} \right) \Phi_i(\cdot, y) = -D^2 \Phi_{i-1}(\cdot, y) \quad \forall i > 0 \quad (4.67)$$

Comparing this to Equations 7.38 and 7.39 we conclude that  $\sigma_{2i} \Phi_i$  satisfies the same set of differential equations. The only possible difference is in the initial conditions.

This is not the case. For  $i = 0$ , the initial condition is that  $\Phi_0(y, y)$  equals the identity map. The symbol map doesn't alter the identity map and so  $\Phi_0(\cdot, y) = \sigma_0(\Phi_0)$ . For  $i > 0$  the initial condition is determined by the requirement that  $\Phi_i$  is differentiable at the origin. Because  $\sigma_{2i}(\Phi_i)$  is differentiable by construction, we conclude the result.  $\square$

If we replace  $T_y M \times \Lambda^\bullet T_y M \otimes \text{End}_{\text{Cl}}(S_y) \rightarrow T_y M$  with the vector bundle  $\mathbb{R} \times T_y M \rightarrow T_y M$  we already have a heat kernel w.r.t.  $\sigma_2(D^2)$ , namely Mehlers kernel. Consider the Taylor series of this kernel in  $R$  and  $F$ . If replace  $R \in M_{n \times n}(\mathbb{R})$  with the Riemann curvature and  $F$  with the twisting curvature we created a kernel on  $\Lambda^\bullet T_y M \otimes \text{End}_{\text{Cl}}$ . Note that the algebra generated by the Riemann curvature and the Twisting curvature is a nilpotent commutative algebra. Hence the Taylor series converges on  $\Lambda^\bullet T_y M \otimes \text{End}_{\text{Cl}}$  and satisfies the heat equation. So Mehlers kernels is also the heat kernel w.r.t.  $\sigma_2(D^2)$  and by uniqueness we conclude

**Proposition 7.24** (Roe [1998], Proposition 12.26). *Let  $S \rightarrow (M, g)$  be a Clifford bundle over an even dimension compact Riemannian manifold. Let  $D$  be the Dirac operator and let  $k_t = \frac{1}{(4\pi t)^{n/2}} \exp(-r^2/4t) \sum_i t^i \Phi_i$  be the formal solution of the heat equation w.r.t.  $D^2$ . Let  $y \in M$ ,  $U_y \subseteq M$  be a neighborhood of  $y$  such that  $\exp_y^{-1}: U_y \rightarrow T_y M$  is a chart of  $M$ , let  $x \in U_y$  and let  $r$  be the distance between  $x$  and  $y$ . Then for all  $i$  the symbol  $\sigma_{2i}(\Phi_i)$  satisfies*

$$\det^{1/2} \left( \frac{tR/2}{\sinh(tR/2)} \right) \exp \left[ -\frac{1}{4t} \left\langle \frac{tR}{2} \coth \left( \frac{tR}{2} \right) x, x \right\rangle - tF \right] = \exp(-r^2/4t) \sum_i t^i \sigma_{2i}(\Phi_i) \quad (7.41)$$

## 7.4 Atiyah-Singer Index theorem

At last we consider the interaction between the symbol map and the trace and we prove the Atiyah-Singer index theorem. From the trace identities we know that only top Clifford degree part are non-vanishing in the trace. In the next definition and lemma we translate this in terms of Getzler symbols.

**Definition 7.25.** *Let  $S \rightarrow (M, g)$  be a Clifford bundle on a compact even dimensional Riemannian manifold and let  $y \in M$ . The **constant part of the Getzler***

**symbol**  $\sigma_k^0$  is the projection of the Getzler symbol  $\sigma_k$  to  $C_0^\infty(T_y M) \otimes \Lambda^\bullet T_y M \otimes \text{End}_{\text{Cl}}(S_y)$  where  $C_0^\infty(T_y M)$  denotes the constant part of the Taylor grading.

**Lemma 7.26.** Let  $S \rightarrow (M, g)$  be a Clifford bundle on a compact ~~even- $n$~~   <sup>$n$</sup>  dimensional Riemannian manifold, let  $y \in M$  and let  $p$  be a section of  $S \otimes S_y$ . Let  $\gamma_5$  be the canonical grading operator. If  $S$  is canonically graded, then

$$\text{tr}(\gamma_5 p(y)) \cdot \text{Vol}(g) = (-i)^{n/2} \text{tr}(\sigma_n^0(p)) \quad (7.42)$$

*Proof.* In this proof we use the letters  $I$  and  $J$  for multi-indices and we denote their degrees with  $|I|$  resp.  $|J|$ . Locally  $p$  is represented by an element  $\sum_{IJ} x^I \otimes c^J \otimes p_{IJ}$  in  $C^\infty(T_y M) \otimes \text{Cl}(T_y M) \otimes \text{End}_{\text{Cl}}(S_y)$ . The trace of  $\gamma_5 p(y)$  equals

$$\text{tr}(\gamma_5 p(y)) = \sum_{IJ} x^I \cdot \text{tr}(\gamma_5 c^J \cdot p_{IJ}) \quad (7.43)$$

Note that if  $|I| \neq 0$ , then  $x^I = 0$  because we evaluate  $x$  at zero. Hence the trace of  $p(y)$  is

$$\text{tr}(\gamma_5 p(y)) = \sum_J \text{tr}(\gamma_5 c^J \cdot p_{\emptyset, J}) \quad (7.44)$$

By lemma 5.22 it follows that  $\text{tr}(\gamma_5 c^J \cdot p_{IJ}) = 0$  if  $|J| \neq n$ . The only non-vanishing multi-index of  $J$  is  $(1, 2, \dots, n)$ . The same lemma states that  $\text{tr}(\gamma_5 c^{(1, \dots, n)} \cdot p_{I, (1, \dots, n)}) = (-i)^{n/2} \text{tr}(p_{I, (1, \dots, n)})$  and we conclude

$$\text{tr}(\gamma_5 p(y)) = (-i)^{n/2} \text{tr}(p_{\emptyset, (1, \dots, n)}) \quad (7.45)$$

The trace over the constant part of the Getzler symbol equals

$$\text{tr}(\sigma_n^0(p)) = \sum_{\substack{I, J \\ |I|=0 \\ |I|+|J|=n}} \text{tr}(x^I \otimes dx^J \otimes p_{IJ}) \quad (7.46)$$

$$= \text{tr}(p_{\emptyset, (1, 2, \dots, n)}) dx^1 \wedge dx^2 \wedge \dots \wedge dx^n \quad (7.47)$$

and so  $\text{tr}(\gamma_5 p(y)) = (-i)^{n/2} \text{tr}(\sigma_n^0(p))$   $\square$

We finally have all the ingredients to prove the Atiyah-Singer index theorem.

**Theorem 7.27** (Atiyah and Singer [1968]). *Let  $S \rightarrow (M, g)$  be a Clifford bundle on a compact  $n$ -dimensional Riemannian manifold  $M$ . Let  $D$  be the Dirac operator. If  $S$  is canonically graded, then the index of  $D$  is the integral over the  $n$ -form part of  $\hat{A}(TM) \wedge \text{ch}^{rel}(S)$ . That is,*

$$\text{Index}(D) = \int_M \hat{A}(TM) \wedge \text{ch}^{rel}(S) \quad (7.48)$$

*Proof.* By Theorem 4.17 each Dirac operator has a unique heat kernel  $k_t^H \in \Gamma(S \boxtimes S)$  such that the corresponding smoothing operator  $e^{-tD^2}$  satisfies  $(\frac{\partial}{\partial t} + D^2)e^{-tD^2} = 0$ . From Proposition 5.10 we know that the composition between the canonical grading operator  $\gamma_5$  and the operator  $e^{-tD^2}$  is traceclass and its trace equals

$$\text{Tr}(\gamma_5 e^{-tD^2}) = \int_{y \in M} \text{tr}(\gamma_5 k_t(y, y)) \text{Vol}(M) \quad (7.49)$$

The McKean-Singer formula states that the trace of  $\gamma_5 e^{-tD^2}$  equals the index of  $D$  and we conclude

$$\text{Index}(D) = \int_{y \in M} \text{tr}(\gamma_5 k_t(x, x)) \text{Vol}(M) \quad (7.50)$$

Now Theorem 4.13 states that  $k_t$  has a formal solution  $\frac{1}{(4\pi t)^{n/2}} e^{-r^2/4t} \sum_i t^i \Phi_i$ . Because  $\text{Index}(D)$  is independent of the choice of  $t$ , all  $t$ -dependent terms in  $\frac{1}{(4\pi t)^{n/2}} e^{-r^2/4t} \sum_i t^i \Phi_i$  will vanish and so

$$\text{Index}(D) = \begin{cases} \frac{1}{(4\pi)^{n/2}} \int_{y \in M} \text{tr}(\gamma_5 \Phi_{n/2}(x, x)) \text{Vol}(M) & \text{if } n \text{ is even} \\ 0 & \text{if } n \text{ is odd} \end{cases} \quad (7.51)$$

By definition  $\hat{A}(TM) \wedge \text{ch}^{rel}(S)$  is an element of  $H^{ev}(M) \otimes \mathbb{C}$ . Hence if  $n$  is odd, then the  $n$ -form part of  $\hat{A}(TM) \wedge \text{ch}^{rel}(S)$  is zero and we conclude

$$\text{Index}(D) = \int_M \hat{A}(TM) \wedge \text{ch}^{rel}(S) \quad (7.52)$$

Now assume that  $n$  is even. By Lemma 7.26 the index of  $D$  equals

$$\text{Index}(D) = \frac{1}{(4\pi i)^{n/2}} \int_M \text{tr}(\sigma_n^0(\Phi_{n/2})) \quad (7.53)$$

According to Proposition 7.24 the constant part of the Getzler symbol  $\sigma_n^0(\Phi_{n/2})$  equals the  $n$ -form part of

$$\det^{1/2} \left( \frac{R/2}{\sinh(R/2)} \right) \exp(-F) \quad (7.54)$$

Up to the constant  $2^{n/2} \times (2\pi i)^{n/2}$  this equals the  $n$ -form part of  $\hat{A}(TM) \wedge \text{ch}^{rel}(S)$  and we conclude

$$\text{Index}(D) = \int_M \hat{A} \wedge \text{ch}^{rel}(S) \quad (7.55)$$

for all  $n \in \mathbb{N}$ .

□

## 8 Summary and Outlook

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

## 9 Glossary: Differential geometry for physicists

There is a joke that "*differential geometry is the study of properties that are invariant under change of notation*" (Lee [2013]). This is especially true when we compare the notation in Riemannian geometry and gauge theory. If one is familiar with general relativity and field theory, then one has already seen many of the constructions used by mathematicians. So in this chapter we will translate the language used by mathematicians into the language used by physicists. In this chapter we assume that  $M$  is a manifold, which is a space-time which the existence of a metric is not required. We also denote  $x$  as a point on  $M$ .

If one is already familiar with differential geometry, one can safely skip this chapter.

### 9.1 Sections and fields

### 9.2 Constructing vector bundles

Type	Field	Element of	Remark
Real vector field	$\phi^\mu$	$\Gamma(TM)$	Also used: $\Omega^1(M)$
	$\phi_\mu$	$\Gamma(T^*M)$	
Complex vector field	$\phi^\mu$	$\Gamma(TM \otimes C)$	Also used: $\Omega^1(M, \mathbb{C})$
	$\phi_\mu$	$\Gamma(T^*M \otimes \mathbb{C})$	
Real gauge Field	$A_\mu^a$	$\Gamma(TM \otimes \mathfrak{g})$	Here $\mathfrak{g}$ is a Lie algebra.
Tensor field	$T^{\mu\nu}$	$\Gamma(TM \otimes TM)$	Also used: $\Gamma(\text{End}(TM))$
	$T^\mu{}_\nu$	$\Gamma(TM \otimes T^*M)$	
	$T_{\mu\nu}$	$\Gamma(T^*M \otimes T^*M)$	
Antisymmetric tensor field	$F^{\mu\nu}$	$\Gamma(TM \wedge TM)$	Also used: $\Gamma(\Lambda^2 TM)$
	$F_{\mu\nu}$	$\Gamma(\Lambda^2 T^*M)$	Also used: $\Omega^2(M)$ .
	$\epsilon_{\mu\nu\rho\sigma}$	$\Gamma(\Lambda^4 T^*M)$	Also used: $\Omega^4(M)$

Table 6: Examples of different fields and the common used notation.

### 9.3 Derivatives and curvature

In calculus we defined the derivative of a function  $f(x)$  at  $a \in \mathbb{R}$  as the limit  $\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$ . For functions there is a canonical way to extend this definition to manifolds. That is, we take the flow  $\phi_t$  in a given direction  $v \in T_a M$  and we follow

it for a given time  $t$ . This gives us a point  $x = \phi_t(a) \in M$ . The derivative  $\frac{\partial f}{\partial v}$  is defined as  $\lim_{t \rightarrow 0} \frac{f(x) - f(a)}{t}$ . However, on curved manifolds there is no canonical method to extend the directional derivative for fields. This is because there is no canonical way to compare different fibers. See figure 5. In this paragraph we study different generalizations of derivatives on fields and we study which extra structure they require.




Figure 5: *In this picture we see a manifold(blue line) and different vectors(red) in a vector bundle. We cannot compare the vectors, because they are in a different fibre(dotted line). So there is no canonical method to generalize the derivative for sections on curved manifolds, because there is no canonical method to compare different fibers.*

For vector fields on the tangent bundle there is a canonical derivative and it is called the **Lie derivative**. For the tangent bundle we use the flow  $\phi_t$  to identify different fibers. Under this identification the Lie derivative of a field  $\psi \in \Gamma(TM)$  in the direction  $v \in \Gamma(TM)$  at  $a \in M$  is  $\lim_{t \rightarrow 0} \frac{\psi(\phi_t(a)) - \psi(a)}{t}$ . Usually we write this as  $\mathcal{L}_v(\psi)$ .

Note that  $v$  and  $\psi$  are the same type of field. To show this equal footing, one also writes  $L_v \psi$  as  $[v, \psi]$ . This bracket is called the **Lie bracket**. It also behaves like a bracket, because one can show that  $[v, \psi] = -[\psi, v]$ . Another usefull identity is  $\mathcal{L}_{[u, v]} w = \mathcal{L}_u \mathcal{L}_v w - \mathcal{L}_v \mathcal{L}_u w$ . In general the flow identifies different fibers of the cotangent bundle. It also works for tensor bundles of  $TM$ . So for all these cases we can define an derivative and for all these cases we call it the Lie derivative.

Another method to define the derivative on fields uses the Leibniz rule. Recall that for multi-variable calculus, the Leibniz rule reads

$$\frac{\partial}{\partial v}(f \cdot \vec{g}) = \frac{\partial f}{\partial v} \cdot \vec{g} + f \cdot \frac{\partial \vec{g}}{\partial v} \quad (9.1)$$

for all  $f \in \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $\vec{g} \in \mathbb{R}^n \rightarrow \mathbb{R}^m$  and  $v \in \mathbb{R}^n$ . A **covariant derivative** is a linear map  $\nabla: \Gamma(TM) \times \Gamma(E) \rightarrow \Gamma(E)$  on a vector bundle  $E \rightarrow M$  that satisfies the Leibniz rule

$$\nabla_\mu(f \cdot \psi) = f \cdot \nabla_\mu \psi + \frac{\partial f}{\partial x^\mu} \psi \quad (9.2)$$

It must also be linear in the following way:

$$\nabla_v(\psi_1 + \psi_2) = \nabla_v(\psi_1) + \nabla_v(\psi_2) \quad (9.3)$$

$$\nabla_{v_1 + f(x)v_2} \psi = \nabla_{v_1} \psi + f(x) \cdot \nabla_{v_2} \psi \quad (9.4)$$

In gauge theory the covariant derivative is defined such that it acts covariant under gauge transformations. However, it also satisfies the properties listed above. Therefore, the covariant derivative in gauge theory is an example of a general covariant derivative. Another example is the Levi-Civita connection. It is the unique covariant derivative on  $TM$  such that it is compatible<sup>16</sup> with the metric and torsion free<sup>17</sup>. A non-example is the Lie-derivative on vector fields. Indeed, for any smooth map  $f \in C^\infty(M)$  and vector fields  $u, v \in \Gamma(M)$ , we have

$$\mathcal{L}_{f_u} v = -\mathcal{L}_v(f \cdot u) \quad (9.5)$$

$$= -\partial_v f \cdot u - f \cdot \mathcal{L}_v u \quad (9.6)$$

$$= -\partial_v f \cdot u + f \cdot \mathcal{L}_u v \quad (9.7)$$

This does not satisfy condition 9.4. Another word for the covariant derivative is a **connection**. Both words are frequently used in this thesis.

For a given connection  $\nabla: \Gamma(E) \rightarrow \Gamma(E \otimes T^*M)$  we define the **curvature tensor**  $K \in \Gamma(T^*M \otimes T^*M \otimes \text{End}(E))$  as

$$K(u, v) = \nabla_u \nabla_v - \nabla_v \nabla_u - \nabla_{[u, v]} \quad (9.8)$$

In local coordinates the term  $\nabla_{[u, v]}$  vanishes and is usually omitted in physics literature. However due to this term, the curvature tensor doesn't acts like a differential operator but acts like a tensor. Indeed, using the Leibniz rule we have for all smooth maps  $f \in C^\infty(M)$  and sections  $\psi \in \Gamma(E)$

$$K(u, v)f \cdot \psi = \nabla_\mu \nabla_\nu f \cdot \psi - \nabla_\nu \nabla_\mu f \cdot \psi - \nabla_{[\mu, \nu]} f \cdot \psi \quad (9.9)$$

$$= +f \cdot \nabla_\mu \nabla_\nu \psi - f \cdot \nabla_\nu \nabla_\mu \psi - f \cdot \nabla_{[\mu, \nu]} \psi \quad (9.10)$$

$$+ \mathcal{L}_\mu \mathcal{L}_\nu f \cdot \psi - \mathcal{L}_\nu \mathcal{L}_\mu f \cdot \psi - \mathcal{L}_{[\mu, \nu]} f \cdot \psi \quad (9.11)$$

$$= f \cdot K(u, v)\psi$$

<sup>16</sup>That is  $\mathcal{L}_u g(v, w) = g(\nabla_u v, w) + g(v, \nabla_u w)$  for all  $u, v, w \in \Gamma(TM)$

<sup>17</sup>That is  $\nabla_u v - \nabla_v u = [u, v]$  for all  $u, v \in \Gamma(TM)$ .

At last we introduce the exterior derivative. This derivative makes use of the fact that antisymmetric tensor fields forms a Grassmann algebra where fields on  $T^*M$  are the Grassmann numbers. In mathematics this algebra is more commonly known as the **exterior algebra on forms** and it is defined as follows: Let  $\phi$  be an antisymmetric  $(0, k)$ -tensor field and let  $\psi$  be an antisymmetric  $(0, m)$ -tensor field. The **wedge product**  $\phi \wedge \psi$  is an antisymmetric  $(0, k + m)$ -tensor field. Up to a constant it is the graded sum of all permutation of the indices. For example if  $\phi$  is a  $(0, 2)$ -tensor field and  $\psi$  is a  $(0, 1)$ -tensor field, then

$$\begin{aligned} (\phi \wedge \psi)_{\mu\nu\rho} \propto & \phi_{\mu\nu}\psi_\rho + \phi_{\nu\rho}\psi_\mu + \phi_{\rho\mu}\psi_\nu - \\ & - \phi_{\nu\mu}\psi_\rho - \phi_{\rho\nu}\psi_\mu - \phi_{\mu\rho}\psi_\nu \end{aligned} \quad (9.12)$$

By direct calculation we can check that  $\phi \wedge \psi$  is antisymmetric in its indices and that  $\phi \wedge \psi = (-1)^k \psi \wedge \phi$ . So the wedge product defines a Grassmann algebra on the antisymmetric tensor fields. Usually the space of antisymmetric  $(0, k)$ -tensors is denoted by  $\Gamma(\Lambda^k T^*M)$  or  $\Omega^k(M)$ . Antisymmetric  $(0, k)$ -tensors are also called  **$k$ -forms**.

Notice that  $\Omega^0(M)$  is just the space of smooth functions and  $\Omega^1(M)$  is the space of vector fields. The **exterior derivative** for a function  $f \in \Omega^0(M)$  is the unique dual vector field  $df: \Gamma(TM) \rightarrow C^\infty(M)$  such that  $df(v) = \frac{\partial f}{\partial v}$  (Compare this to the definition of the variational derivative). Hence,  $d$  is a map from  $C^\infty(M) = \Omega^0(M)$  to  $\Gamma(T^*M) = \Omega^1(M)$ . We extend  $d$  to a map from  $\Omega^k(M)$  to  $\Omega^{k+1}(M)$  using the following rules:

- $d \circ d: \Omega^k(M) \rightarrow \Omega^{k+2}(M) = 0 \quad \forall k \in \mathbb{N}$
- $d(\alpha \wedge \beta) = (d\alpha) \wedge \beta + (-1)^k \alpha \wedge (d\beta) \quad \forall \alpha \in \Omega^k(M), \beta \in \Omega^m(M)$

The exterior derivative is related to the Lie derivative by the use of the **interior product**. This is a map  $\iota: \Gamma(TM) \times \Omega^k(M) \rightarrow \Omega^{k-1}(M)$  which reduces a form with one degree. For a  $(0, k)$ -tensor field  $T$  it is defined as

$$(\iota_\mu T)_{\nu\rho\sigma\dots} = T_{\mu\nu\rho\sigma\dots} \quad (9.13)$$

Cartan has showed that the Lie derivative can be expressed as

$$\mathcal{L}_\mu = \iota_\mu d + d\iota_\mu \quad (9.14)$$

This result is called **Cartans magic formula**. Another usefull identities of the interior product are

$$\iota_u \iota_v = -\iota_v \iota_u \quad (9.15)$$

$$\iota_u(\alpha \wedge \beta) = (\iota_u \alpha) \wedge \beta + (-1)^k \alpha \wedge (\iota_u \beta) \quad \forall \alpha \in \Omega^k(M) \quad (9.16)$$

$$\iota_{[u,v]} = \mathcal{L}_u \iota_v - \iota_v \mathcal{L}_u \quad (9.17)$$

To show the interaction between these three derivatives we calculate the curvature in local coordinates. We denote  $\{e_\mu\}$  for a basis on  $TM$  and  $\{s_i\}$  for a basis a vector bundle  $E \rightarrow M$ . Let  $\nabla$  be the connection on  $E$  and let  $K$  be the curvature tensor. Using the basis we create a matrix of one-forms

$$\omega_{ij} = \langle s_i | \nabla_\bullet | s_j \rangle \quad (9.18)$$

This matrix is called the **connection 1-form**. The curvature tensor is a matrix of two forms. For  $u, v \in \Gamma(TM)$ , the components of this matrix are given by

$$\langle s_i | K(u, v) | s_j \rangle = \langle s_i | \nabla_u \nabla_v - \nabla_v \nabla_u - \nabla_{[u, v]} | s_j \rangle \quad (9.19)$$

To use the Leibniz rule we introduce the identity  $1 = \sum_k |s_k\rangle \langle s_k|$

$$\langle s_i | K(u, v) | s_j \rangle = \langle s_i | \nabla_u (|s_k\rangle \langle s_k| \nabla_v | s_j \rangle) - \langle s_i | \nabla_v (|s_k\rangle \langle s_k| \nabla_u | s_j \rangle) - \langle s_i | \nabla_{[u, v]} | s_j \rangle \quad (9.20)$$

$$= \langle s_i | \nabla_u (\omega_{kj}(v) \cdot |s_k\rangle) - \langle s_i | \nabla_v (\omega_{kj}(u) \cdot |s_k\rangle) - \omega_{ij}([u, v]) \quad (9.21)$$

For each component  $\omega_{ij}(v)$  is a smooth function and hence we apply the Leibniz rule:

$$\begin{aligned} \langle s_i | K(u, v) | s_j \rangle &= \mathcal{L}_u(\omega_{kj}(v)) \langle s_i | s_k \rangle + \omega_{ik}(u) \omega_{kj}(v) \\ &\quad - \mathcal{L}_v(\omega_{kj}(u)) \langle s_i | s_k \rangle - \omega_{ik}(v) \omega_{kj}(u) - \omega_{ij}([u, v]) \end{aligned} \quad (9.22)$$

$$\begin{aligned} &= \mathcal{L}_u(\omega_{ij}(v)) - \mathcal{L}_v(\omega_{ij}(u)) - \omega_{ij}([u, v]) \\ &\quad + \omega_{ik}(u) \omega_{kj}(v) - \omega_{ik}(v) \omega_{kj}(u) \end{aligned} \quad (9.23)$$

In physics this equation is usually written as  $K_{\mu\nu} = \partial_\mu \omega_\nu - \partial_\nu \omega_\mu + [\omega_\mu, \omega_\nu]$ . So we view  $\omega$  as a gauge potential and  $K$  as the field strength. To reformulate this result in terms of forms we write  $\omega_{ij}(u)$  as  $\iota_u \omega_{ij}$ . Note that the wedge product between functions coincide with the normal product.

$$\begin{aligned} \langle s_i | K(u, v) | s_j \rangle &= \mathcal{L}_u(\iota_v \omega_{ij}) - \mathcal{L}_v(\iota_u \omega_{ij}) - \iota_{[u, v]} \omega_{ij} \\ &\quad + (\iota_u \omega_{ik}) \wedge (\iota_v \omega_{kj}) - (\iota_v \omega_{ik}) \wedge (\iota_u \omega_{kj}) \end{aligned} \quad (9.24)$$

Using equation 9.17 we get that  $\mathcal{L}_u(\iota_v \omega_{ij}) - \iota_{[u, v]} \omega_{ij} = \iota_v \mathcal{L}_u \omega_{ij}$ . From Cartans magic formula follows that

$$\begin{aligned} \langle s_i | K(u, v) | s_j \rangle &= \iota_v (\iota_u d + d \iota_u) \omega_{ij} - (\iota_v d + d \iota_v) (\iota_u \omega_{ij}) \\ &\quad + (\iota_u \omega_{ik}) \wedge (\iota_v \omega_{kj}) - (\iota_v \omega_{ik}) \wedge (\iota_u \omega_{kj}) \end{aligned} \quad (9.25)$$

Forms of degree -1 doesn't exists. Therefore  $\iota_v \iota_u \omega_{ij} = 0$  and

$$\langle s_i | K(u, v) | s_j \rangle = \iota_v \iota_u d \omega_{ij} + (\iota_u \omega_{ik}) \wedge (\iota_v \omega_{kj}) - (\iota_v \omega_{ik}) \wedge (\iota_u \omega_{kj}) \quad (9.26)$$

We use equation 9.16 twice to get

$$\langle s_i | K(u, v) | s_j \rangle = \iota_v \iota_u d\omega_{ij} + \iota_v ((\iota_u \omega_{ik}) \wedge \omega_{kj} - \omega_{ik} \wedge (\iota_u \omega_{kj})) \quad (9.27)$$

$$= \iota_v \iota_u (d\omega_{ij} + \omega_{ik} \wedge \omega_{kj}) \quad (9.28)$$

Viewing  $K$  as a matrix of 2-forms, we conclude that

$$K = d\omega + \omega \wedge \omega \quad (9.29)$$

Using equation 9.29 we can derive the Bianchi identity. Indeed, denote  $K_{ij} = \langle s_i | K(\cdot, \cdot) | s_j \rangle \in \Omega^2(M)$ . From equation 9.28 follows that

$$K_{ij} = d w_{ij} - w_{ik} \wedge w_{kj} \quad (9.30)$$

We take the exterior derivative of  $K_{ij}$ . Recall that  $d^2 = 0$  and so

$$d K_{ij} = d^2 w_{ij} - d(w_{ik} \wedge w_{kj}) = -d(w_{ik} \wedge w_{kj}) \quad (9.31)$$

By definition  $d(w_{ik} \wedge w_{kj}) = (d w_{ik}) \wedge w_{kj} - w_{ik} \wedge (d w_{kj})$ . Using equation 9.30 again we conclude

$$d K_{ij} = w_{ik} \wedge (K_{kj} + w_{kl} \wedge w_{lj}) - (K_{ik} + w_{il} \wedge w_{lk}) \wedge w_{kj} \quad (9.32)$$

$$= w_{ik} \wedge (K_{kj} - K_{ik} \wedge w_{kj}) \quad (9.33)$$

Viewing  $K$  as a matrix, we conclude

$$d K = \omega \wedge K - K \wedge \omega \quad (9.34)$$

This is the **second Bianchi identity**.

## 9.4 Topology of vector bundles

As mathematicians we often ask the question: "How can we tell two spaces apart?". One of the main theorems in differential geometry is that the exterior derivative can be used to discriminate different manifolds. First we recall this result and then we explain how this is relevant for physics.

Before we know the difference between manifold we need to know when they are the same. There is a saying that for a mathematician a donut and a coffee mug are equal. How weird as this sounds but if the mug was not baked then the mag can be continuously molded into a donut. That is, it can be molded *without tearing, creating*




Figure 6: How to transform a coffee mug into a donut (Commons [2007])

or *closing holes*. This is shown in figure 6. We say that if two spaces are equivalent up to continuous deformation they are **homotopy equivalent**.

Look at figure 7. We added a subscript to the exterior derivative to keep track of the vector spaces it acts on. So  $d_{k-1}$  is a map from  $\Omega^{k-1}(M)$  into  $\Omega^k(M)$ . It doesn't necessary mean that the whole space  $\Omega^k(M)$  is reached. We call the subspace in  $\Omega^k(M)$  that is reached the **image** of  $d_{k-1}$  and is denoted by  $\text{Im } d_{k-1}$ . In figure 7 the images are depicted as red squares. The blue square represent the **kernel** of  $d_k$ . It is the subspace in  $\Omega^k(M)$  that is mapped to zero by  $d_k$  and we denote it by  $\ker d_k$ . Because  $d_k d_{k-1} = 0$ , the image always lies inside of the kernel. That is, the red square lies inside the blue square. The part of the blue that is not covered by the red square is called the  $k$ -th cohomology class. We denote this by  $H_{dR}^k(M)$ . The collection of all  $H_{dR}^k(M)$  is called the **de Rham cohomology**. Formally  $H_{dR}^k(M)$  is a subspace of  $\ker d_k$  that is orthogonal to  $\text{Im } d_{k-1}$ .




Figure 7: In this image the red squares represents the image of the exterior derivative. The blue squares represent the kernel of  $d$ . Because  $d^2 = 0$ , the image always lies inside the kernel.

One of the main theorem in differential geometry is that cohomology is invariant under continuous deformations. That means that if two spaces are homotopy equivalent,

then they must have the same cohomology classes. So if the cohomology classes differ, then the manifolds must differ.

Also vector bundles can be characterized using differential operators. For this we don't use the exterior derivative but the curvature tensor. The idea is that from the curvature tensor we create elements in  $H_{dR}^\bullet(M)$  called **characteristic classes**. These classes don't depend on the choice of connection and so it only depends on the shape of the vector bundle. This plays an important role in the study of anomalies. Indeed, anomalies are given in terms of characteristic classes. For example, the abelian anomaly is proportional to

$$\epsilon^{\mu\nu\rho\sigma} \text{Tr}(F_{\mu\nu} F_{\rho\sigma}) \quad (9.35)$$

This is proportional to the Second Chern class. So the shape of the vector bundle breaks the classical conservation laws. More details on characteristic classes can be found in chapter 6

## **10 Field theory in terms of differential geometry**

## 11 Appendix: Source codes

In this chapter we add the source codes we used to do the nasty calculations. We performed the calculations using FORM from Vermaseren [2000].

### 11.1 Calculation of Bardeen Anomaly

```

*#-;

*
* Define global variables
*

Indices m1, ..., m10;
Indices x1, ..., x10;

Functions g5, g;
CFunction ep(a);

Symbol i;
Symbol z1,..., z10;

.global

*
* Give an explicit expression for  $D^2$ . The results are hardcoded
*    $\rightarrow$  later.
*

Symbol M;
Functions D1, D2, d;
Functions Pp, Pm;

Functions V,A, dA, dV;
CFunctions k;

* D
Local expr1 = 1/M^2 * (g(m1) * D1(m1) * g(m2) * D1(m2));

```

```

* Dbar
Local expr2 = 1/M^2 * (g(m1) * D2(m1) * g(m2) * D2(m2));

* Give definition of D and D Bar
Id D1(x1?) = d(x1) - i * V(x1) + A(x1) * g5;
Id D2(x1?) = d(x1) - i * V(x1) - A(x1) * g5;

* Reorder Clifford Multiplication
repeat;
Id d(x1?) * g(x2?) = g(x2) * d(x1);
Id V(x1?) * g(x2?) = g(x2) * V(x1);
Id A(x1?) * g(x2?) = g(x2) * A(x1);

Id g(x1?) * g5 = - g5 * g(x1);
Id A(x1?) * g5 = g5 * A(x1);
Id V(x1?) * g5 = g5 * V(x1);
Id d(x1?) * g5 = g5 * d(x1);

Id g5 * g5 = 1;
endrepeat;

* Pull e^{ikx} through expression
Id d(x1?) = d(x1) + i * k(x1) * M;
Id i^2 = -1;

* Write clifford multiplication using commutator and anticommutator
Id g(x1?)*g(x2?) = d_(x1,x2) + 1/2 * (g(x1)*g(x2) - g(x2)*g(x1));
Id g(x1?)*g(x2?)*d(x2?)*d(x1?) = g(x1)*g(x2)*d(x1)*d(x2);

* Take derivatives
repeat;
Id d(x1?) * A(x2?) = dA(x1,x2) + A(x2) * d(x1);
Id d(x1?) * V(x2?) = dV(x1,x2) + V(x2) * d(x1);
endrepeat;

* Print answer
Sum m1, ..., m10;

```

```

Bracket M,g, g5;
Print;

.store
Symbols s,t M;

*
* Calculate the Bardeen Anomaly
*

Functions Xa, Xb, Ya, Yb;

Set spaceIndOdd: m1,m3,m5,m7,m9;
Set spaceIndEven: m2,m4,m6,m8,m10;

Functions Xa1, ..., Xa4;
Functions Ya1, ..., Ya4;
Functions Xb1, ..., Xb4;
Functions Yb1, ..., Yb4;

Set gFields: Xa1, ..., Xa4,Xb1, ..., Xb4, Ya1, ..., Ya4, Yb1, ...,
    ↪ Yb4;
Functions gField;

Functions V, A, d;
Functions dA, dV;

Functions F,G;
CFunctions k;

*
* Expand Exp (Xa/M^2 + Ya/M) + Exp (Xb/M^2 + Yb/M)
*

Global expr = t/2 * g5 * M^4 * (1+
#do a = 1, 5
+ 1
#do b = 1, 'a'

```

```

* (Xa(spaceIndOdd['b'], spaceIndEven['b'])/M^2 + Ya(spaceIndOdd['b'
  ↳ ], spaceIndEven['b'])/M)
#enddo
/ fac_('a')
#enddo
) +
t/2 * g5 * M^4 * (1+
#do a = 1, 5
+ 1
#do b = 1, 'a'
* (Xb(spaceIndOdd['b'], spaceIndEven['b'])/M^2 + Yb(spaceIndOdd['b'
  ↳ ], spaceIndEven['b'])/M)
#enddo
/ fac_('a')
#enddo
);

* Consider only constant order in M
Id M^{-1} = 0;

* Write X = X1 + g5 * X2 + g(mu) g(nu) X3(mu,nu) + g5 g(mu)g(nu) X4
  ↳ (mu,nu). Same for Y
Id Xa(x1?, x2?) = Xa1(x1,x2) + g5 * Xa2(x1,x2) + g(x1) * g(x2) *
  ↳ Xa3(x1,x2) + g5 * g(x1) * g(x2) * Xa4(x1, x2);
Id Ya(x1?, x2?) = Ya1(x1,x2) + g5 * Ya2(x1,x2) + g(x1) * g(x2) *
  ↳ Ya3(x1,x2) + g5 * g(x1) * g(x2) * Ya4(x1, x2);

Id Xb(x1?, x2?) = Xb1(x1,x2) + g5 * Xb2(x1,x2) + g(x1) * g(x2) *
  ↳ Xb3(x1,x2) + g5 * g(x1) * g(x2) * Xb4(x1, x2);
Id Yb(x1?, x2?) = Yb1(x1,x2) + g5 * Yb2(x1,x2) + g(x1) * g(x2) *
  ↳ Yb3(x1,x2) + g5 * g(x1) * g(x2) * Yb4(x1, x2);

* Order gamma matrices
repeat;
Id gField?gFields * g5 = g5 * gField;
Id gField?gFields(x1?) * g5 = g5 * gField(x1);
Id gField?gFields(x1?, x2?) * g5 = g5 * gField(x1, x2);

```

```

Id g(x1?) * g5 = - g5 * g(x1);
Id g5 * g5 = 1;

Id gField?gFields * g(x1?) = g(x1) * gField;
Id gField?gFields(x1?) * g(x2?) = g(x2) * gField(x1);
Id gField?gFields(x1?, x2?) * g(x3?) = g(x3) * gField(x1, x2);
endrepeat;

* Fill in earlier results for Y
Id Ya1(x1?,x2?) = 2*d(x1)*k(x1)*i + 2*V(x1)*k(x1);
Id Ya2(x1?,x2?) = 0;
Id Ya3(x1?,x2?) = 0;
Id Ya4(x1?,x2?) = - A(x1)*k(x2)*i + A(x2)*k(x1)*i;

Id Yb1(x1?,x2?) = 2*d(x1)*k(x1)*i + 2*V(x1)*k(x1);
Id Yb2(x1?,x2?) = 0;
Id Yb3(x1?,x2?) = 0;
Id Yb4(x1?,x2?) = A(x1)*k(x2)*i - A(x2)*k(x1)*i;

Id i^2 = -1;

* Perform K integration
Id k(x1?) * k(x2?) * k(x3?) * k(x4?) = (d_(x1,x2)* d_(x3,x4) + d_(
  ↪ x1,x3)* d_(x2,x4) + d_(x1,x4)* d_(x2,x3))/4;
Id k(x1?) * k(x2?) * k(x3?) = 0;
Id k(x1?) * k(x2?) = d_(x1,x2)/2;
Id k(x1?) = 0;

* Group gamma functions and remove odd gamma fields
Id g5 * g(x1?) * g(x2?) * g(x3?) * g(x4?) * g(x5?) * g(x6?) * g(x7
  ↪ ?) * g(x8?) = g5(x1,x2,x3,x4,x5,x6,x7,x8);;
Id g(x1?) * g(x2?) * g(x3?) * g(x4?) * g(x5?) * g(x6?) * g(x7?) * g
  ↪ (x8?) = g(x1,x2,x3,x4,x5,x6,x7,x8);
Id g5 * g(x1?) * g(x2?) * g(x3?) * g(x4?) * g(x5?) * g(x6?) * g(x7
  ↪ ?) = 0;
Id g(x1?) * g(x2?) * g(x3?) * g(x4?) * g(x5?) * g(x6?) * g(x7?) =
  ↪ 0;

```

```

Id g5 * g(x1?) * g(x2?) * g(x3?) * g(x4?) * g(x5?) * g(x6?) = g5(x1
  ↪ ,x2,x3,x4,x5,x6);;
Id g(x1?) * g(x2?) * g(x3?) * g(x4?) * g(x5?) * g(x6?) = g(x1,x2,x3
  ↪ ,x4,x5,x6);
Id g5 * g(x1?) * g(x2?) * g(x3?) * g(x4?) * g(x5?) = 0;
Id g(x1?) * g(x2?) * g(x3?) * g(x4?) * g(x5?) = 0;
Id g5 * g(x1?) * g(x2?) * g(x3?) * g(x4?) = g5(x1,x2,x3,x4);
Id g(x1?) * g(x2?) * g(x3?) * g(x4?) = g(x1,x2,x3,x4);
Id g5 * g(x1?) * g(x2?) * g(x3?) = 0;
Id g(x1?) * g(x2?) * g(x3?) = 0;
Id g5 * g(x1?) * g(x2?) = 0;
Id g(x1?) * g(x2?) = g(x1,x2);
Id g5 * g(x1?) = 0;
Id g(x1?) = 0;
Id g5 = 0;

* Now simplify gamma expressions
repeat;
Id g(?p11, x1?,x1?, ?p12) = 4 * g(?p11, ?p12);
Id g5(?p11, x1?,x1?, ?p12) = 4 * g5(?p11, ?p12);
Id g(?p11, x1?,?p12,x2?,x1?, ?p13) = - g(?p11, x1,?p12,x1,x2, ?p13)
  ↪ + 2 * g(?p11, x2,?p12,?p13);
Id g5(?p11, x1?,?p12,x2?,x1?, ?p13) = - g5(?p11, x1,?p12,x1,x2, ?
  ↪ p13) + 2 * g5(?p11, x2,?p12,?p13);
endrepeat;

repeat;
Id g5(x1?,x2?) = 0;
Id g5(x1?,x2?,x3?,x4?) = 4 * ep(x1,x2,x3,x4)/t;
endrepeat;

* Fill in earlier results for X
Id Xa1(x1?,x2?) = d(x1)*d(x1) - 2*V(x1)*d(x1)*i - V(x1)*V(x1) - A(
  ↪ x1)*A(x1) - dV(x1,x1)*i;
Id Xa2(x1?,x2?) = - V(x1)*A(x1)*i + A(x1)*V(x1)*i + dA(x1,x1);
Id Xa3(x1?,x2?) = - 1/2*V(x1)*V(x2) + 1/2*V(x2)*V(x1) - 1/2*A(x1)*A
  ↪ (x2) + 1/2*A(x2)*A(x1) - 1/2*dV(x1,x2)*i + 1/2*dV(x2,x1)*i;

```

```

Id Xa4(x1?,x2?) = - 1/2*V(x1)*A(x2)*i + 1/2*V(x2)*A(x1)*i - A(x1)*d
  ↪ (x2) + 1/2*A(x1)*V(x2)*i + A(x2)*d(x1) - 1/2*A(x2)*V(x1)*i +
  ↪ 1/2*dA(x1,x2) - 1/2*dA(x2,x1);

Id Xb1(x1?,x2?) = d(x1)*d(x1) - 2*V(x1)*d(x1)*i - V(x1)*V(x1) - A(
  ↪ x1)*A(x1) - dV(x1,x1)*i;
Id Xb2(x1?,x2?) = V(x1)*A(x1)*i - A(x1)*V(x1)*i - dA(x1,x1);
Id Xb3(x1?,x2?) = - 1/2*V(x1)*V(x2) + 1/2*V(x2)*V(x1) - 1/2*A(x1)*A
  ↪ (x2) + 1/2*A(x2)*A(x1) - 1/2*dV(x1,x2)*i + 1/2*dV(x2,x1)*i ;
Id Xb4(x1?,x2?) = 1/2*V(x1)*A(x2)*i - 1/2*V(x2)*A(x1)*i + A(x1)*d(
  ↪ x2) - 1/2*A(x1)*V(x2)*i - A(x2)*d(x1) + 1/2*A(x2)*V(x1)*i -
  ↪ 1/2*dA(x1,x2) + 1/2*dA(x2,x1);

* Take derivatives
repeat;
Id d(x1?) * A(x2?) = dA(x1,x2) + A(x2) * d(x1);
Id d(x1?) * V(x2?) = dV(x1,x2) + V(x2) * d(x1);
Id d(x1?) * dA(x2?, x3?) = dA(x2,x3) * d(x1);
Id d(x1?) * dV(x2?, x3?) = dV(x2,x3) * d(x1);
endrepeat;
Id d(x1?) = 0;

* Rewrite in terms of Bardeen Curvatures
Id dA(x1?,x2?) = 1/2 * (G(x1,x2) + i * V(x1)*A(x2) - i *A(x2) * V(
  ↪ x1) + i*A(x1)*V(x2) - i *V(x2) * A(x1));
Id dV(x1?,x2?) = 1/2 * (F(x1,x2) + i *V(x1)*V(x2) - i * V(x2) * V(
  ↪ x1) - i * A(x1)*A(x2) + i *A(x2) * A(x1));

Id i^2 = -1;

* Return result
Sum m1,...,m10;
Bracket ep, g, g5;
Print;

.store;
.end

```

## Index

- $\hat{A}$ -genus, 103
- $d$ -symbol, 39
- $i^{\text{th}}$  Chern class, 102
  
- Abelian chiral symmetry, 23
- Anomalies, 22
- Anomalous Ward identity, 54
- Associated graded algebra, 107
- Associated symbol map, 107
  
- Bundle of Clifford modules, 8
  
- Canonical grading operator, 11
- Cartans magic formula, 123
- Characteristic class, 102
- Chern  $f$ -genus, 102
- Chern character, 103
- Chiral current, 24
- Clifford action, 8
- Clifford algebra, 6
- Clifford bundle, 9
- Clifford contraction, 17
- Clifford endomorphism, 20
- Compact operators, 86
- Connection, 122
- Connection 1-form, 98, 124
- Consistent anomaly, 54
- Covariant anomaly, 48
- Covariant derivative, 122
- Curvature tensor, 122
  
- De Rham cohomology, 126
- Degree of a filtered algebra, 106
- Degree of a graded algebra, 105
- Dimensional regularization, 26
- Dirac operator, 10
  
- Directional derivative of an invariant polynomial, 99
  
- Exterior algebra, 123
- Exterior derivative, 123
  
- Feynman-slash notation, 9
- Filtered algebra, 106
- Formal solution of the heat equation, 67
- Forms, 123
  
- Generalized Laplacian, 58
- Getzler filtration for kernels, 110
- Getzler filtration for operators, 110
- Getzler symbol, 110
- Graded algebra, 105
- Graded Clifford bundle, 10
- Grading operator, 11
  
- Heat kernel, 59
- Hilbert-Schmidt, 79
- Hilbert-Schmidt inner product, 82
- Hilbert-Schmidt norm, 81
- Homotopy equivalent, 126
  
- Image of a linear operator, 126
- Index of a Dirac operator, 91
- Interior product, 123
- Invariant polynomial, 96
  
- Kernel of a linear operator, 126
- Kernel of an operator, 59
  
- Laplacian, 15
- Lichnerowicz formula, 20
- Lie bracket, 121
- Lie derivative, 121

McKean-Singer formula, 90  
Mehlers kernel, 61  
  
Noethers Current, 23  
Noethers theorem, 22  
Non-Abelian chiral symmetry, 24  
  
Relative Chern character, 104  
Riemann endomorphism, 18  
Riemannian normal coordinate system, 13  
  
Second Bianchi identity, 125  
Smoothing operator, 59  
Spectrum, 86  
Structure constant, 40  
Symbol map, 107  
  
Total Chern class, 102  
Trace, 83  
Traceclass, 83  
Twisting curvature, 20  
  
Wedge product, 123  
Weitzenbock formula, 17

## References

- Stephen L. Adler and William A. Bardeen. Absence of higher-order corrections in the anomalous axial-vector divergence equation. *Phys. Rev.*, 182:1517–1536, Jun 1969. doi: 10.1103/PhysRev.182.1517. URL <https://link.aps.org/doi/10.1103/PhysRev.182.1517>.
- L. Alvarez-Gaumé and P. Ginsparg. The topological meaning of non-abelian anomalies. *Nuclear Physics B*, 243(3):449 – 474, 1984. ISSN 0550-3213. doi: [https://doi.org/10.1016/0550-3213\(84\)90487-5](https://doi.org/10.1016/0550-3213(84)90487-5). URL <http://www.sciencedirect.com/science/article/pii/0550321384904875>.
- Luis Alvarez-Gaumé and Paul Ginsparg. The structure of gauge and gravitational anomalies. *Annals of Physics*, 161(2):423 – 490, 1985. ISSN 0003-4916. doi: [https://doi.org/10.1016/0003-4916\(85\)90087-9](https://doi.org/10.1016/0003-4916(85)90087-9). URL <http://www.sciencedirect.com/science/article/pii/0003491685900879>.
- A. Andrianov and L. Bonora. Finite-mode regularization of the fermion functional integral. *Nuclear Physics B*, 233(2):232 – 246, 1984. ISSN 0550-3213. doi: [https://doi.org/10.1016/0550-3213\(84\)90413-9](https://doi.org/10.1016/0550-3213(84)90413-9). URL <http://www.sciencedirect.com/science/article/pii/0550321384904139>.
- M. F. Atiyah and I. M. Singer. The index of elliptic operators. I. *Ann. of Math. (2)*, 87:484–530, 1968. ISSN 0003-486X. doi: 10.2307/1970715. URL <https://doi.org/10.2307/1970715>.
- William A. Bardeen. Anomalous Ward identities in spinor field theories. *Phys. Rev.*, 184:1848–1857, 1969. doi: 10.1103/PhysRev.184.1848.
- Nicole Berline, Ezra Getzler, and Michèle Vergne. *Heat kernels and Dirac operators*. Grundlehren Text Editions. Springer-Verlag, Berlin, 2004. ISBN 3-540-20062-2. Corrected reprint of the 1992 original.
- R. A. Bertlmann. *Anomalies in quantum field theory*. 1996.
- C. G. Bollini and J. J. Giambiagi. Dimensional renormalization : The number of dimensions as a regularizing parameter. *Il Nuovo Cimento B (1971-1996)*, 12(1):20–26, Nov 1972. ISSN 1826-9877. doi: 10.1007/BF02895558. URL <https://doi.org/10.1007/BF02895558>.
- Wikimedia Commons. A coffee mug morphing into a torus, a popular example in topology, 2007. URL

- [https://en.wikipedia.org/wiki/File:Mug\\_and\\_Torus\\_morph.gif](https://en.wikipedia.org/wiki/File:Mug_and_Torus_morph.gif). File: Mug and Torus morph.gif.
- K. Fujikawa and H. Suzuki. *Path integrals and quantum anomalies*. 2004. doi: 10.1093/acprof:oso/9780198529132.001.0001.
- Kazuo Fujikawa. Path Integral for Gauge Theories with Fermions. *Phys. Rev.*, D21:2848, 1980. doi: 10.1103/PhysRevD.21.2848,10.1103/PhysRevD.22.1499. [Erratum: *Phys. Rev.*D22,1499(1980)].
- Howard Georgi and Sheldon L. Glashow. Gauge theories without anomalies. *Phys. Rev. D*, 6:429--431, Jul 1972. doi: 10.1103/PhysRevD.6.429. URL <https://link.aps.org/doi/10.1103/PhysRevD.6.429>.
- Ezra Getzler. Pseudodifferential operators on supermanifolds and the atiyah-singer index theorem. *Comm. Math. Phys.*, 92(2):163--178, 1983. URL <https://projecteuclid.org/443/euclid.cmp/1103940796>.
- Shi-ke Hu, Bing-Lin Young, and Douglas W. McKay. Functional Integral and Minimal Anomalies in Theories With  $S$ ,  $P$ ,  $V$  and a Currents. *Phys. Rev.*, D30:836, 1984. doi: 10.1103/PhysRevD.30.836.
- John M. Lee. *Introduction to smooth manifolds*, volume 218 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2013. ISBN 978-1-4419-9981-8.
- André Lichnerowicz. Spineurs harmoniques. *C. R. Acad. Sci. Paris*, 257:7--9, 1963.
- H. P. McKean, Jr. and I. M. Singer. Curvature and the eigenvalues of the laplacian. *J. Differential Geom.*, 1 (1-2):43--69, 1967. doi: 10.4310/jdg/1214427880. URL <https://doi.org/10.4310/jdg/1214427880>.
- F. G. Mehler. Ueber die Entwicklung einer Function von beliebig vielen Variablen nach Laplaceschen Functionen höherer Ordnung. *J. Reine Angew. Math.*, 66:161--176, 1866. ISSN 0075-4102. doi: 10.1515/crll.1866.66.161. URL <https://doi.org/10.1515/crll.1866.66.161>.

- John W. Milnor and James D. Stasheff. *Characteristic classes*. Princeton University Press, Princeton, N. J.; University of Tokyo Press, Tokyo, 1974. Annals of Mathematics Studies, No. 76.
- Gerard J. Murphy. *C\*-algebras and operator theory*. Academic Press, San Diego, 1990. ISBN 978-0-08-092496-0. doi: <https://doi.org/10.1016/B978-0-08-092496-0.50005-3>. URL <https://www.sciencedirect.com/science/article/pii/B9780080924960500053>.
- John Roe. *Elliptic operators, topology and asymptotic methods*, volume 395 of *Pitman Research Notes in Mathematics Series*. Longman, Harlow, second edition, 1998. ISBN 0-582-32502-1.
- G. 't Hooft and M. Veltman. Regularization and renormalization of gauge fields. *Nuclear Physics B*, 44(1):189 -- 213, 1972. ISSN 0550-3213. doi: [https://doi.org/10.1016/0550-3213\(72\)90279-9](https://doi.org/10.1016/0550-3213(72)90279-9). URL <http://www.sciencedirect.com/science/article/pii/0550321372902799>.
- P. van Nieuwenhuizen. *Anomalies in quantum field theory: Cancellation of anomalies in  $d = 10$  supergravity*, volume B3 of *Leuven notes in mathematical and theoretical physics*. Leuven Univ. Pr., Leuven, Belgium, 1989.
- J. A. M. Vermaseren. New features of FORM. 2000.
- J. Wess and B. Zumino. Consequences of anomalous Ward identities. *Phys. Lett.*, 37B:95--97, 1971. doi: 10.1016/0370-2693(71)90582-X.