# **INLDS Practicum 7**

Study the following two-parameter planar systems exhibiting local codim 2 bifurcations by combining analytical and numerical methods.

## Exercises

#### Ex.1 Takens normal form

$$\begin{cases} \dot{x} = \beta x + y + x^2, \\ \dot{y} = \alpha - 4x^2. \end{cases}$$
(1)

- (a) Derive equations for the fold and Hopf bifurcations in the system.
- (b) Prove that a Bogdanov–Takens (BT) bifurcation occurs in the system and find the corresponding parameter values.
- (c) Compute the normal form coefficients a and b for the BT-bifurcation and verify that  $ab \neq 0$ .
- (d) Use **pplane9** to produce all representative phase portraits of the system near the Bogdanov-Takens point. Sketch the bifurcation set of the system and give phase portraits.
- (e) For  $\alpha = 0.25$  find numerically the value of  $\beta$  corresponding to the saddle homoclinic bifurcation.

### Ex.2 A prey-predator model by Bazykin and Khibnik

$$\begin{cases} \dot{x} = \frac{x^2(1-x)}{n+x} - xy, \\ \dot{y} = -y(m-x), \end{cases}$$
(2)

where  $x, y \ge 0$  and 0 < m < 1.

- (a) Using **pplane9**, produce several phase portraits of model (2) for different values of m and fixed  $n = \frac{1}{4}$  and  $n = \frac{1}{16}$ . *Hint*: The most interesting phase portrait occurs at  $(m, n) = (\frac{1}{5}, \frac{1}{16})$ .
- (b) For  $m = \frac{1}{5}$  approximate by simulations the value of n corresponding to the collision and disappearance of two periodic orbits.
- (c) Derive an equation for the Hopf bifurcation in the model. *Hint*: Consider the orbitally equivalent (why?) polynomial system

$$\begin{cases} \dot{x} = x^2(1-x) - xy(n+x), \\ \dot{y} = -y(m-x)(n+x). \end{cases}$$
(3)

- (d) Compute the 1st Lyapunov coefficient  $l_1$  along the Hopf bifurcation line in (3) and prove that this coefficient vanishes at  $(m, n) = (\frac{1}{4}, \frac{1}{8})$ . Conclude that a Bautin bifurcation happens in the model.
- (e) Sketch the bifurcation set of the model and give phase portraits.
- (f) **Challenge**: Prove that the Bautin bifurcation is nondegenerate by computing the corresponding 2nd Lyapunov coefficient  $l_2$ .

## Homework

Hand-in is exercise 2.