

# On Steinhilber's conjecture

**Peter Koymans**  
University of Michigan



*UC Irvine Number Theory Seminar*

21 April 2022

# History of Pell's equation

For a fixed squarefree integer  $d > 0$ , the equation

$$x^2 - dy^2 = 1 \text{ to be solved in } x, y \in \mathbb{Z}$$

has been studied since at least the ancient Greeks.

# History of Pell's equation

For a fixed squarefree integer  $d > 0$ , the equation

$$x^2 - dy^2 = 1 \text{ to be solved in } x, y \in \mathbb{Z}$$

has been studied since at least the ancient Greeks.

Bhaskara II (12th century) gave an algorithm to find non-trivial solutions of this equation.

# History of Pell's equation

For a fixed squarefree integer  $d > 0$ , the equation

$$x^2 - dy^2 = 1 \text{ to be solved in } x, y \in \mathbb{Z}$$

has been studied since at least the ancient Greeks.

Bhaskara II (12th century) gave an algorithm to find non-trivial solutions of this equation.

Unbeknownst of Bhaskara's work, Fermat challenged English mathematicians Brouncker and Wallis to solve the notorious case  $d = 61$ .

# History of Pell's equation

For a fixed squarefree integer  $d > 0$ , the equation

$$x^2 - dy^2 = 1 \text{ to be solved in } x, y \in \mathbb{Z}$$

has been studied since at least the ancient Greeks.

Bhaskara II (12th century) gave an algorithm to find non-trivial solutions of this equation.

Unbeknownst of Bhaskara's work, Fermat challenged English mathematicians Brouncker and Wallis to solve the notorious case  $d = 61$ .

The smallest non-trivial solution is

$$1766319049^2 - 61 \cdot 226153980^2 = 1.$$

# History of Pell's equation

For a fixed squarefree integer  $d > 0$ , the equation

$$x^2 - dy^2 = 1 \text{ to be solved in } x, y \in \mathbb{Z}$$

has been studied since at least the ancient Greeks.

Bhaskara II (12th century) gave an algorithm to find non-trivial solutions of this equation.

Unbeknownst of Bhaskara's work, Fermat challenged English mathematicians Brouncker and Wallis to solve the notorious case  $d = 61$ .

The smallest non-trivial solution is

$$1766319049^2 - 61 \cdot 226153980^2 = 1.$$

Lagrange was the first to give an algorithm with proof of correctness.

# A modern interpretation of Pell's equation

In modern terms, we know that

$$x^2 - dy^2 = 1$$

always has a non-trivial solution by Dirichlet's Unit Theorem.

# A modern interpretation of Pell's equation

In modern terms, we know that

$$x^2 - dy^2 = 1$$

always has a non-trivial solution by Dirichlet's Unit Theorem.

Indeed,  $\mathcal{O}_K^* = \{\pm\epsilon^k\}$  for a real quadratic field  $K$ .



# A modern interpretation of Pell's equation

In modern terms, we know that

$$x^2 - dy^2 = 1$$

always has a non-trivial solution by Dirichlet's Unit Theorem.

Indeed,  $\mathcal{O}_K^* = \{\pm\epsilon^k\}$  for a real quadratic field  $K$ .

The equation

$$x^2 - dy^2 = -1$$

is known as the negative Pell equation and is not always soluble.

# A modern interpretation of Pell's equation

In modern terms, we know that

$$x^2 - dy^2 = 1$$

always has a non-trivial solution by Dirichlet's Unit Theorem.

Indeed,  $\mathcal{O}_K^* = \{\pm\epsilon^k\}$  for a real quadratic field  $K$ .

The equation

$$x^2 - dy^2 = -1$$

is known as the negative Pell equation and is not always soluble.

Question: as we vary  $d$ , how often is the negative Pell equation soluble?

# Solubility over the rationals

Define  $\mathcal{D}$  to be the set of squarefree integers having as odd prime divisors only primes  $p \equiv 1 \pmod{4}$  and define  $\mathcal{D}^-$  to be the set of squarefree integers for which negative Pell is soluble.

# Solubility over the rationals

Define  $\mathcal{D}$  to be the set of squarefree integers having as odd prime divisors only primes  $p \equiv 1 \pmod{4}$  and define  $\mathcal{D}^-$  to be the set of squarefree integers for which negative Pell is soluble.

By the Hasse-Minkowski Theorem we have for all squarefree  $d$

$$d \in \mathcal{D} \iff x^2 - dy^2 = -1 \text{ is soluble with } x, y \in \mathbb{Q},$$

so in particular  $\mathcal{D}^-$  is a subset of  $\mathcal{D}$ .

# Solubility over the rationals

Define  $\mathcal{D}$  to be the set of squarefree integers having as odd prime divisors only primes  $p \equiv 1 \pmod{4}$  and define  $\mathcal{D}^-$  to be the set of squarefree integers for which negative Pell is soluble.

By the Hasse-Minkowski Theorem we have for all squarefree  $d$

$$d \in \mathcal{D} \iff x^2 - dy^2 = -1 \text{ is soluble with } x, y \in \mathbb{Q},$$

so in particular  $\mathcal{D}^-$  is a subset of  $\mathcal{D}$ .

Classical techniques in analytic number theory give a constant  $C > 0$  such that

$$\#\{d \leq X : d \in \mathcal{D}\} \sim C \cdot \frac{X}{\sqrt{\log X}}.$$

# Solubility over the rationals

Define  $\mathcal{D}$  to be the set of squarefree integers having as odd prime divisors only primes  $p \equiv 1 \pmod{4}$  and define  $\mathcal{D}^-$  to be the set of squarefree integers for which negative Pell is soluble.

By the Hasse-Minkowski Theorem we have for all squarefree  $d$

$$d \in \mathcal{D} \iff x^2 - dy^2 = -1 \text{ is soluble with } x, y \in \mathbb{Q},$$

so in particular  $\mathcal{D}^-$  is a subset of  $\mathcal{D}$ .

Classical techniques in analytic number theory give a constant  $C > 0$  such that

$$\#\{d \leq X : d \in \mathcal{D}\} \sim C \cdot \frac{X}{\sqrt{\log X}}.$$

Refined question: what is the density of  $\mathcal{D}^-$  inside  $\mathcal{D}$ ?

# Conjectures on the negative Pell equation

Nagell (1930s) conjectured that

$$\lim_{X \rightarrow \infty} \frac{\#\{d \leq X : d \in \mathcal{D}^-\}}{\#\{d \leq X : d \in \mathcal{D}\}}$$

exists and lies in  $(0, 1)$ .

# Conjectures on the negative Pell equation

Nagell (1930s) conjectured that

$$\lim_{X \rightarrow \infty} \frac{\#\{d \leq X : d \in \mathcal{D}^-\}}{\#\{d \leq X : d \in \mathcal{D}\}}$$

exists and lies in  $(0, 1)$ .

Stevenhagen (1995) conjectured that

$$\lim_{X \rightarrow \infty} \frac{\#\{d \leq X : d \in \mathcal{D}^-\}}{\#\{d \leq X : d \in \mathcal{D}\}} = 1 - \alpha,$$

where

$$\alpha = \prod_{j=1}^{\infty} (1 + 2^{-j})^{-1} \approx 0.41942.$$



# Progress towards Stevenhagen's conjecture

Dirichlet showed that all primes  $p \equiv 1 \pmod{4}$  are in  $\mathcal{D}^-$ .

# Progress towards Stevenhagen's conjecture

Dirichlet showed that all primes  $p \equiv 1 \pmod{4}$  are in  $\mathcal{D}^-$ .

Blomer showed that

$$\#\{d \leq X : d \in \mathcal{D}^-\} \gg \frac{X}{(\log X)^{0.62}}.$$

# Progress towards Stevenhagen's conjecture

Dirichlet showed that all primes  $p \equiv 1 \pmod{4}$  are in  $\mathcal{D}^-$ .

Blomer showed that

$$\#\{d \leq X : d \in \mathcal{D}^-\} \gg \frac{X}{(\log X)^{0.62}}.$$

Fouvry and Klüners (2010) proved that

$$\frac{5\alpha}{4} \leq \liminf_{X \rightarrow \infty} \frac{\#\{d \leq X : d \in \mathcal{D}^-\}}{\#\{d \leq X : d \in \mathcal{D}\}} \leq \limsup_{X \rightarrow \infty} \frac{\#\{d \leq X : d \in \mathcal{D}^-\}}{\#\{d \leq X : d \in \mathcal{D}\}} \leq \frac{2}{3}.$$

# Progress towards Stevenhagen's conjecture

Dirichlet showed that all primes  $p \equiv 1 \pmod{4}$  are in  $\mathcal{D}^-$ .

Blomer showed that

$$\#\{d \leq X : d \in \mathcal{D}^-\} \gg \frac{X}{(\log X)^{0.62}}.$$

Fouvry and Klüners (2010) proved that

$$\frac{5\alpha}{4} \leq \liminf_{X \rightarrow \infty} \frac{\#\{d \leq X : d \in \mathcal{D}^-\}}{\#\{d \leq X : d \in \mathcal{D}\}} \leq \limsup_{X \rightarrow \infty} \frac{\#\{d \leq X : d \in \mathcal{D}^-\}}{\#\{d \leq X : d \in \mathcal{D}\}} \leq \frac{2}{3}.$$

CKMP (2019) improved the lower bound to

$$\alpha \cdot \sum_{n=0}^{\infty} 2^{-n(n+3)/2} \approx \alpha \cdot 1.28325.$$

# Stevenhagen's conjecture is true

## Theorem 1 (K., Pagano (2021))

We have

$$\lim_{X \rightarrow \infty} \frac{\#\{d \leq X : d \in \mathcal{D}^-\}}{\#\{d \leq X : d \in \mathcal{D}\}} = 1 - \alpha$$

in accordance with Stevenhagen's conjecture.

# Stevenhagen's conjecture is true

## Theorem 1 (K., Pagano (2021))

We have

$$\lim_{X \rightarrow \infty} \frac{\#\{d \leq X : d \in \mathcal{D}^-\}}{\#\{d \leq X : d \in \mathcal{D}\}} = 1 - \alpha$$

in accordance with Stevenhagen's conjecture.

## Corollary 2

We have

$$\#\{d \leq X : d \in \mathcal{D}^-\} \sim C \cdot (1 - \alpha) \cdot \frac{X}{\sqrt{\log X}}.$$

## A criterion for solubility

Recall that the narrow class group  $\text{Cl}^+(K)$  is defined as the quotient of the ideal group  $I_K$  by the principal ideals  $P_K^+$  admitting a totally positive generator, while the class group is the quotient by the principal ideals  $P_K$ .

# A criterion for solubility

Recall that the narrow class group  $\text{Cl}^+(K)$  is defined as the quotient of the ideal group  $I_K$  by the principal ideals  $P_K^+$  admitting a totally positive generator, while the class group is the quotient by the principal ideals  $P_K$ .

We have

$$\begin{aligned}x^2 - dy^2 = -1 \text{ is soluble} &\iff \text{fundamental unit } \epsilon \text{ has negative norm} \\ &\iff (\sqrt{d}) \text{ is trivial in } \text{Cl}^+(\mathbb{Q}(\sqrt{d})).\end{aligned}$$



# A criterion for solubility

Recall that the narrow class group  $\text{Cl}^+(K)$  is defined as the quotient of the ideal group  $I_K$  by the principal ideals  $P_K^+$  admitting a totally positive generator, while the class group is the quotient by the principal ideals  $P_K$ .

We have

$$\begin{aligned}x^2 - dy^2 = -1 \text{ is soluble} &\iff \text{fundamental unit } \epsilon \text{ has negative norm} \\ &\iff (\sqrt{d}) \text{ is trivial in } \text{Cl}^+(\mathbb{Q}(\sqrt{d})).\end{aligned}$$

There is a fundamental exact sequence

$$1 \rightarrow \frac{P_K}{P_K^+} \rightarrow \text{Cl}^+(K) \rightarrow \text{Cl}(K) \rightarrow 1$$

with  $\# \frac{P_K}{P_K^+} \in \{1, 2\}$  and  $\frac{P_K}{P_K^+}$  generated by  $(\sqrt{d})$ .

# A criterion for solubility

Recall that the narrow class group  $\text{Cl}^+(K)$  is defined as the quotient of the ideal group  $I_K$  by the principal ideals  $P_K^+$  admitting a totally positive generator, while the class group is the quotient by the principal ideals  $P_K$ .

We have

$$\begin{aligned}x^2 - dy^2 = -1 \text{ is soluble} &\iff \text{fundamental unit } \epsilon \text{ has negative norm} \\ &\iff (\sqrt{d}) \text{ is trivial in } \text{Cl}^+(\mathbb{Q}(\sqrt{d})).\end{aligned}$$

There is a fundamental exact sequence

$$1 \rightarrow \frac{P_K}{P_K^+} \rightarrow \text{Cl}^+(K) \rightarrow \text{Cl}(K) \rightarrow 1$$

with  $\# \frac{P_K}{P_K^+} \in \{1, 2\}$  and  $\frac{P_K}{P_K^+}$  generated by  $(\sqrt{d})$ .

Goal: study joint distribution of  $(\text{Cl}^+(K)[2^\infty], \text{Cl}(K)[2^\infty])$ .

# Genus theory

The group  $\text{Cl}^+(K)[2]$  has a very predictable behavior unlike  $\text{Cl}^+(K)[p]$  for  $p$  odd.

# Genus theory

The group  $\text{Cl}^+(K)[2]$  has a very predictable behavior unlike  $\text{Cl}^+(K)[p]$  for  $p$  odd.

The description of  $\text{Cl}^+(K)[2]$  is due to Gauss and is known as genus theory. We have that

$$\#\text{Cl}^+(K)[2] = 2^{\omega(D_K)-1}$$

and  $\text{Cl}^+(K)[2]$  is generated by the ramified prime ideals of  $\mathcal{O}_K$ .

# Genus theory

The group  $\text{Cl}^+(K)[2]$  has a very predictable behavior unlike  $\text{Cl}^+(K)[p]$  for  $p$  odd.

The description of  $\text{Cl}^+(K)[2]$  is due to Gauss and is known as genus theory. We have that

$$\#\text{Cl}^+(K)[2] = 2^{\omega(D_K)-1}$$

and  $\text{Cl}^+(K)[2]$  is generated by the ramified prime ideals of  $\mathcal{O}_K$ .

If  $p$  divides the discriminant of  $\mathbb{Q}(\sqrt{d})$ , then  $p$  ramifies, so

$$\begin{array}{ccc} \mathbb{Q}(\sqrt{d}) & \mathfrak{p} & \mathfrak{p}^2 = (p). \\ | & | & \\ \mathbb{Q} & p & \end{array}$$

There is precisely one relation between the ramified primes.

Let  $p$  be an odd prime. The group  $\text{Cl}(K)[p^\infty]$  is believed to behave as a random finite, abelian  $p$ -group.

Let  $p$  be an odd prime. The group  $\text{Cl}(K)[p^\infty]$  is believed to behave as a random finite, abelian  $p$ -group.

More formally, Cohen and Lenstra conjectured that

$$\lim_{X \rightarrow \infty} \frac{\#\{K \text{ im. quadr.} : |D_K| < X \text{ and } \text{Cl}(K)[p^\infty] \cong A\}}{\#\{K \text{ im. quadr.} : |D_K| < X\}} = \frac{\prod_{i=1}^{\infty} \left(1 - \frac{1}{p^i}\right)}{\#\text{Aut}(A)}$$

for every finite, abelian  $p$ -group  $A$ .

# Gerth's adaptation

Gerth adapted the conjecture of Cohen and Lenstra to  $p = 2$

$$\lim_{X \rightarrow \infty} \frac{\#\{K \text{ im. quadr.} : |D_K| < X, 2\text{Cl}(K)[2^\infty] \cong A\}}{\#\{K \text{ im. quadr.} : |D_K| < X\}} = \frac{\prod_{i=1}^{\infty} (1 - \frac{1}{2^i})}{\#\text{Aut}(A)}$$

for every finite, abelian 2-group  $A$ .



# Gerth's adaptation

Gerth adapted the conjecture of Cohen and Lenstra to  $p = 2$

$$\lim_{X \rightarrow \infty} \frac{\#\{K \text{ im. quadr.} : |D_K| < X, 2\text{Cl}(K)[2^\infty] \cong A\}}{\#\{K \text{ im. quadr.} : |D_K| < X\}} = \frac{\prod_{i=1}^{\infty} (1 - \frac{1}{2^i})}{\#\text{Aut}(A)}$$

for every finite, abelian 2-group  $A$ .

## Theorem 3 (Alexander Smith (2017))

*Gerth's conjecture is true.*

# Gerth's adaptation

Gerth adapted the conjecture of Cohen and Lenstra to  $p = 2$

$$\lim_{X \rightarrow \infty} \frac{\#\{K \text{ im. quadr.} : |D_K| < X, 2\text{Cl}(K)[2^\infty] \cong A\}}{\#\{K \text{ im. quadr.} : |D_K| < X\}} = \frac{\prod_{i=1}^{\infty} (1 - \frac{1}{2^i})}{\#\text{Aut}(A)}$$

for every finite, abelian 2-group  $A$ .

## Theorem 3 (Alexander Smith (2017))

*Gerth's conjecture is true.*

Strategy: adapt Smith's method to the family  $\mathcal{D}$ .

# Gerth's adaptation

Gerth adapted the conjecture of Cohen and Lenstra to  $p = 2$

$$\lim_{X \rightarrow \infty} \frac{\#\{K \text{ im. quadr.} : |D_K| < X, 2\text{Cl}(K)[2^\infty] \cong A\}}{\#\{K \text{ im. quadr.} : |D_K| < X\}} = \frac{\prod_{i=1}^{\infty} (1 - \frac{1}{2^i})}{\#\text{Aut}(A)}$$

for every finite, abelian 2-group  $A$ .

## Theorem 3 (Alexander Smith (2017))

*Gerth's conjecture is true.*

Strategy: adapt Smith's method to the family  $\mathcal{D}$ .

Two major difficulties:  $\mathcal{D}$  has density 0 in the squarefree integers, and  $\mathcal{D}$  naturally ends up in the error term in Smith's proof!

# Strategy for Stevenhagen's conjecture

## Example 1 (Definition of $2^k$ -rank)

Take

$$A = \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/8\mathbb{Z} \oplus \mathbb{Z}/3\mathbb{Z} \quad .$$

Then  $\text{rk}_2 A = 3$ ,  $\text{rk}_4 A = \text{rk}_8 A = 1$ ,  $\text{rk}_{2^k} A = 0$  for  $k > 3$ .

# Strategy for Stevenhagen's conjecture

## Example 1 (Definition of $2^k$ -rank)

Take

$$A = \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/8\mathbb{Z} \oplus \mathbb{Z}/3\mathbb{Z} \quad (\text{rk}_{2^k} A := \dim_{\mathbb{F}_2} 2^{k-1}A/2^k A).$$

Then  $\text{rk}_2 A = 3$ ,  $\text{rk}_4 A = \text{rk}_8 A = 1$ ,  $\text{rk}_{2^k} A = 0$  for  $k > 3$ .

Find for every integer  $m \geq 1$ , the density of  $d \in \mathcal{D}$  for which

$$\text{rk}_{2^k} \text{Cl}^+(\mathbb{Q}(\sqrt{d})) = \text{rk}_{2^k} \text{Cl}(\mathbb{Q}(\sqrt{d})) > 0 \text{ for } 1 \leq k \leq m \text{ and}$$

$$\text{rk}_{2^{m+1}} \text{Cl}^+(\mathbb{Q}(\sqrt{d})) = 0.$$

This gives better and better lower bounds for negative Pell.

# Strategy for Stevenhagen's conjecture

## Example 1 (Definition of $2^k$ -rank)

Take

$$A = \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/8\mathbb{Z} \oplus \mathbb{Z}/3\mathbb{Z} \quad (\text{rk}_{2^k} A := \dim_{\mathbb{F}_2} 2^{k-1}A/2^k A).$$

Then  $\text{rk}_2 A = 3$ ,  $\text{rk}_4 A = \text{rk}_8 A = 1$ ,  $\text{rk}_{2^k} A = 0$  for  $k > 3$ .

Find for every integer  $m \geq 1$ , the density of  $d \in \mathcal{D}$  for which

$$\begin{aligned} \text{rk}_{2^k} \text{Cl}^+(\mathbb{Q}(\sqrt{d})) = \text{rk}_{2^k} \text{Cl}(\mathbb{Q}(\sqrt{d})) > 0 \text{ for } 1 \leq k \leq m \text{ and} \\ \text{rk}_{2^{m+1}} \text{Cl}^+(\mathbb{Q}(\sqrt{d})) = 0. \end{aligned}$$

This gives better and better lower bounds for negative Pell. Similarly, find for every integer  $m \geq 1$ , the density of  $d \in \mathcal{D}$  for which

$$\begin{aligned} \text{rk}_{2^k} \text{Cl}^+(\mathbb{Q}(\sqrt{d})) = \text{rk}_{2^k} \text{Cl}(\mathbb{Q}(\sqrt{d})) > 0 \text{ for } 1 \leq k \leq m \text{ and} \\ \text{rk}_{2^{m+1}} \text{Cl}^+(\mathbb{Q}(\sqrt{d})) = \text{rk}_{2^{m+1}} \text{Cl}(\mathbb{Q}(\sqrt{d})) + 1. \end{aligned}$$

This gives better and better upper bounds for negative Pell.

# Strategy for Stevenhagen's conjecture

## Example 1 (Definition of $2^k$ -rank)

Take

$$A = \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/8\mathbb{Z} \oplus \mathbb{Z}/3\mathbb{Z} \quad (\text{rk}_{2^k} A := \dim_{\mathbb{F}_2} 2^{k-1}A/2^k A).$$

Then  $\text{rk}_2 A = 3$ ,  $\text{rk}_4 A = \text{rk}_8 A = 1$ ,  $\text{rk}_{2^k} A = 0$  for  $k > 3$ .

Find for every integer  $m \geq 1$ , the density of  $d \in \mathcal{D}$  for which

$$\begin{aligned} \text{rk}_{2^k} \text{Cl}^+(\mathbb{Q}(\sqrt{d})) = \text{rk}_{2^k} \text{Cl}(\mathbb{Q}(\sqrt{d})) > 0 \text{ for } 1 \leq k \leq m \text{ and} \\ \text{rk}_{2^{m+1}} \text{Cl}^+(\mathbb{Q}(\sqrt{d})) = 0. \end{aligned}$$

This gives better and better lower bounds for negative Pell. Similarly, find for every integer  $m \geq 1$ , the density of  $d \in \mathcal{D}$  for which

$$\begin{aligned} \text{rk}_{2^k} \text{Cl}^+(\mathbb{Q}(\sqrt{d})) = \text{rk}_{2^k} \text{Cl}(\mathbb{Q}(\sqrt{d})) > 0 \text{ for } 1 \leq k \leq m \text{ and} \\ \text{rk}_{2^{m+1}} \text{Cl}^+(\mathbb{Q}(\sqrt{d})) = \text{rk}_{2^{m+1}} \text{Cl}(\mathbb{Q}(\sqrt{d})) + 1. \end{aligned}$$

This gives better and better upper bounds for negative Pell.

Strategy: adapt Smith's ideas to compute these densities.

# Strategy for Stevenhagen's conjecture

## Example 1 (Definition of $2^k$ -rank)

Take

$$A = \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/8\mathbb{Z} \oplus \mathbb{Z}/3\mathbb{Z} \quad (\text{rk}_{2^k} A := \dim_{\mathbb{F}_2} 2^{k-1}A/2^k A).$$

Then  $\text{rk}_2 A = 3$ ,  $\text{rk}_4 A = \text{rk}_8 A = 1$ ,  $\text{rk}_{2^k} A = 0$  for  $k > 3$ .

Find for every integer  $m \geq 1$ , the density of  $d \in \mathcal{D}$  for which

$$\begin{aligned} \text{rk}_{2^k} \text{Cl}^+(\mathbb{Q}(\sqrt{d})) = \text{rk}_{2^k} \text{Cl}(\mathbb{Q}(\sqrt{d})) > 0 \text{ for } 1 \leq k \leq m \text{ and} \\ \text{rk}_{2^{m+1}} \text{Cl}^+(\mathbb{Q}(\sqrt{d})) = 0. \end{aligned}$$

This gives better and better lower bounds for negative Pell. Similarly, find for every integer  $m \geq 1$ , the density of  $d \in \mathcal{D}$  for which

$$\begin{aligned} \text{rk}_{2^k} \text{Cl}^+(\mathbb{Q}(\sqrt{d})) = \text{rk}_{2^k} \text{Cl}(\mathbb{Q}(\sqrt{d})) > 0 \text{ for } 1 \leq k \leq m \text{ and} \\ \text{rk}_{2^{m+1}} \text{Cl}^+(\mathbb{Q}(\sqrt{d})) = \text{rk}_{2^{m+1}} \text{Cl}(\mathbb{Q}(\sqrt{d})) + 1. \end{aligned}$$

This gives better and better upper bounds for negative Pell.

Strategy: adapt Smith's ideas to compute these densities.



# Duality of abelian groups

For a finite abelian group  $A$ , define

$$A^\vee := \text{Hom}(A, \mathbb{C}^*).$$

There is a natural pairing

$$\text{Art}_1 : A[2] \times A^\vee[2] \rightarrow \{\pm 1\}, \quad (a, \chi) \mapsto \chi(a).$$

Left kernel of  $\text{Art}_1$  is  $2A[4]$  and right kernel is  $2A^\vee[4]$ .

# Duality of abelian groups

For a finite abelian group  $A$ , define

$$A^\vee := \text{Hom}(A, \mathbb{C}^*).$$

There is a natural pairing

$$\text{Art}_1 : A[2] \times A^\vee[2] \rightarrow \{\pm 1\}, \quad (a, \chi) \mapsto \chi(a).$$

Left kernel of  $\text{Art}_1$  is  $2A[4]$  and right kernel is  $2A^\vee[4]$ .

So 4-rank is determined by the pairing  $\text{Art}_1$ . We start by describing  $\text{Cl}^{+, \vee}(K)[2]$ .

# The dual class group

## Theorem 4 (Class field theory)

*We have an isomorphism*

$$\mathrm{Cl}^+(K) \cong \mathrm{Gal}(H^+(K)/K)$$

*given by sending a prime ideal  $\mathfrak{p}$  to  $\mathrm{Art}(\mathfrak{p})$ . Furthermore, if  $K$  is Galois, this isomorphism respects the natural Galois action of  $\mathrm{Gal}(K/\mathbb{Q})$  on both sides.*

# The dual class group

## Theorem 4 (Class field theory)

*We have an isomorphism*

$$\text{Cl}^+(K) \cong \text{Gal}(H^+(K)/K)$$

*given by sending a prime ideal  $\mathfrak{p}$  to  $\text{Art}(\mathfrak{p})$ . Furthermore, if  $K$  is Galois, this isomorphism respects the natural Galois action of  $\text{Gal}(K/\mathbb{Q})$  on both sides.*

From this we get a bijection

$$\text{Cl}^{+,v}(K)[2] \leftrightarrow \{\text{quadratic unramified extensions of } K\}.$$

# The dual class group

## Theorem 4 (Class field theory)

*We have an isomorphism*

$$\text{Cl}^+(K) \cong \text{Gal}(H^+(K)/K)$$

*given by sending a prime ideal  $\mathfrak{p}$  to  $\text{Art}(\mathfrak{p})$ . Furthermore, if  $K$  is Galois, this isomorphism respects the natural Galois action of  $\text{Gal}(K/\mathbb{Q})$  on both sides.*

From this we get a bijection

$$\text{Cl}^{+, \vee}(K)[2] \leftrightarrow \{\text{quadratic unramified extensions of } K\}.$$

If  $K$  is quadratic with odd discriminant, then  $\text{Cl}^{+, \vee}(K)[2]$  is generated by the quadratic characters  $\chi_{p^*}$ , where  $p^*$  satisfies  $|p^*| = |p|$  and  $p^* \equiv 1 \pmod{4}$ .

# The dual class group

## Theorem 4 (Class field theory)

We have an isomorphism

$$\mathrm{Cl}^+(K) \cong \mathrm{Gal}(H^+(K)/K)$$

given by sending a prime ideal  $\mathfrak{p}$  to  $\mathrm{Art}(\mathfrak{p})$ . Furthermore, if  $K$  is Galois, this isomorphism respects the natural Galois action of  $\mathrm{Gal}(K/\mathbb{Q})$  on both sides.

From this we get a bijection

$$\mathrm{Cl}^{+, \vee}(K)[2] \leftrightarrow \{\text{quadratic unramified extensions of } K\}.$$

If  $K$  is quadratic with odd discriminant, then  $\mathrm{Cl}^{+, \vee}(K)[2]$  is generated by the quadratic characters  $\chi_{p^*}$ , where  $p^*$  satisfies  $|p^*| = |p|$  and  $p^* \equiv 1 \pmod{4}$ . In particular

$$d \in \mathcal{D} \iff \mathrm{rk}_2 \mathrm{Cl}^+(\mathbb{Q}(\sqrt{d})) = \mathrm{rk}_2 \mathrm{Cl}(\mathbb{Q}(\sqrt{d})) \iff (\sqrt{d}) \in 2\mathrm{Cl}^+(\mathbb{Q}(\sqrt{d}))[4].$$

# The Artin pairing

Under the earlier identifications, we have that

$$\text{Art}_1 : \text{Cl}^+(K)[2] \times \text{Cl}^{+,\vee}(K)[2] \rightarrow \{\pm 1\}, \quad (\mathfrak{p}, \chi) \mapsto \chi(\text{Art } \mathfrak{p}).$$

# The Artin pairing

Under the earlier identifications, we have that

$$\text{Art}_1 : \text{Cl}^+(K)[2] \times \text{Cl}^{+,\vee}(K)[2] \rightarrow \{\pm 1\}, \quad (\mathfrak{p}, \chi) \mapsto \chi(\text{Art } \mathfrak{p}).$$

Let  $p_1, \dots, p_t$  be the prime divisors of  $d$ . The Rédei matrix is

$$\begin{array}{ccccc} & \chi_{p_1^*} & \chi_{p_2^*} & \cdots & \chi_{p_t^*} \\ p_1 & * & \left(\frac{p_2^*}{p_1}\right) & \cdots & \left(\frac{p_t^*}{p_1}\right) \\ p_2 & \left(\frac{p_1^*}{p_2}\right) & * & \cdots & \left(\frac{p_t^*}{p_2}\right) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ p_t & \left(\frac{p_1^*}{p_t}\right) & \left(\frac{p_2^*}{p_t}\right) & \cdots & * \end{array}.$$



# The Artin pairing

Under the earlier identifications, we have that

$$\text{Art}_1 : \text{Cl}^+(K)[2] \times \text{Cl}^{+,\vee}(K)[2] \rightarrow \{\pm 1\}, \quad (\mathfrak{p}, \chi) \mapsto \chi(\text{Art } \mathfrak{p}).$$

Let  $\rho_1, \dots, \rho_t$  be the prime divisors of  $d$ . The Rédei matrix is

$$\begin{array}{ccccc} & \chi_{\rho_1^*} & \chi_{\rho_2^*} & \cdots & \chi_{\rho_t^*} \\ \rho_1 & * & \left(\frac{\rho_2^*}{\rho_1}\right) & \cdots & \left(\frac{\rho_t^*}{\rho_1}\right) \\ \rho_2 & \left(\frac{\rho_1^*}{\rho_2}\right) & * & \cdots & \left(\frac{\rho_t^*}{\rho_2}\right) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \rho_t & \left(\frac{\rho_1^*}{\rho_t}\right) & \left(\frac{\rho_2^*}{\rho_t}\right) & \cdots & * \end{array}.$$

Left kernel surjects on  $2\text{Cl}^+(K)[4]$  with 1-dimensional kernel.

## Interlude: Stevenhagen's conjecture

For  $d \in \mathcal{D}$ , recall that  $(\sqrt{d}) \in 2\text{Cl}^+(\mathbb{Q}(\sqrt{d}))[4]$ .

## Interlude: Stevenhagen's conjecture

For  $d \in \mathcal{D}$ , recall that  $(\sqrt{d}) \in 2\text{Cl}^+(\mathbb{Q}(\sqrt{d}))[4]$ .

Heuristic assumption: every non-zero element in the left kernel of the Rédei matrix is equally likely to be trivial.

# Interlude: Stevenhagen's conjecture

For  $d \in \mathcal{D}$ , recall that  $(\sqrt{d}) \in 2\text{Cl}^+(\mathbb{Q}(\sqrt{d}))[4]$ .

Heuristic assumption: every non-zero element in the left kernel of the Rédei matrix is equally likely to be trivial.

## Conjecture 1 (Stevenhagen's conjecture)

*We have*

$$\lim_{X \rightarrow \infty} \frac{\#\{d \leq X : d \in \mathcal{D}^-\}}{\#\{d \leq X : d \in \mathcal{D}\}} = \sum_{j=0}^{\infty} \frac{\mathbb{P}(\text{4-rank of } d \in \mathcal{D} \text{ equals } j)}{2^{j+1} - 1}.$$

# Interlude: Stevenhagen's conjecture

For  $d \in \mathcal{D}$ , recall that  $(\sqrt{d}) \in 2\text{Cl}^+(\mathbb{Q}(\sqrt{d}))[4]$ .

Heuristic assumption: every non-zero element in the left kernel of the Rédei matrix is equally likely to be trivial.

## Conjecture 1 (Stevenhagen's conjecture)

We have

$$\lim_{X \rightarrow \infty} \frac{\#\{d \leq X : d \in \mathcal{D}^-\}}{\#\{d \leq X : d \in \mathcal{D}\}} = \sum_{j=0}^{\infty} \frac{\mathbb{P}(\text{4-rank of } d \in \mathcal{D} \text{ equals } j)}{2^{j+1} - 1}.$$

Furthermore,

$$\mathbb{P}(\text{4-rank of } d \in \mathcal{D} \text{ equals } j) = \lim_{t \rightarrow \infty} \mathbb{P}(t \times t \text{ sym. matrix has ker. of dim. } j).$$

## The second Artin pairing

There is a natural pairing

$$\text{Art}_2 : 2A[4] \times 2A^\vee[4] \rightarrow \{\pm 1\}, \quad (a, \chi) \mapsto \psi(a), \quad 2\psi = \chi.$$

Left kernel is  $4A[8]$  and right kernel is  $4A^\vee[8]$ .

# The second Artin pairing

There is a natural pairing

$$\text{Art}_2 : 2A[4] \times 2A^\vee[4] \rightarrow \{\pm 1\}, \quad (a, \chi) \mapsto \psi(a), \quad 2\psi = \chi.$$

Left kernel is  $4A[8]$  and right kernel is  $4A^\vee[8]$ .

As before, class field theory gives that this pairing becomes

$$(\mathfrak{p}, \chi) \mapsto \psi(\text{Art } \mathfrak{p}), \quad 2\psi = \chi.$$

First goal: understand cyclic degree 4 unramified extensions of  $\mathbb{Q}(\sqrt{d})$ .

# The second Artin pairing

There is a natural pairing

$$\text{Art}_2 : 2A[4] \times 2A^\vee[4] \rightarrow \{\pm 1\}, \quad (a, \chi) \mapsto \psi(a), \quad 2\psi = \chi.$$

Left kernel is  $4A[8]$  and right kernel is  $4A^\vee[8]$ .

As before, class field theory gives that this pairing becomes

$$(\mathfrak{p}, \chi) \mapsto \psi(\text{Art } \mathfrak{p}), \quad 2\psi = \chi.$$

First goal: understand cyclic degree 4 unramified extensions of  $\mathbb{Q}(\sqrt{d})$ .

Fact: a degree 4 unramified, abelian extension of  $\mathbb{Q}(\sqrt{d})$  is Galois over  $\mathbb{Q}$  with Galois group  $D_4$ .



# The second Artin pairing

There is a natural pairing

$$\text{Art}_2 : 2A[4] \times 2A^\vee[4] \rightarrow \{\pm 1\}, \quad (a, \chi) \mapsto \psi(a), \quad 2\psi = \chi.$$

Left kernel is  $4A[8]$  and right kernel is  $4A^\vee[8]$ .

As before, class field theory gives that this pairing becomes

$$(\mathfrak{p}, \chi) \mapsto \psi(\text{Art } \mathfrak{p}), \quad 2\psi = \chi.$$

First goal: understand cyclic degree 4 unramified extensions of  $\mathbb{Q}(\sqrt{d})$ .

Fact: a degree 4 unramified, abelian extension of  $\mathbb{Q}(\sqrt{d})$  is Galois over  $\mathbb{Q}$  with Galois group  $D_4$ .

Such extensions are of the shape  $\mathbb{Q}(\sqrt{d}, \sqrt{a}, \sqrt{\alpha})$ , where

$$x^2 = ay^2 + \frac{d}{a}z^2 \text{ with } x, y, z \in \mathbb{Z} \text{ and } \gcd(x, y, z) = 1, \quad \alpha := x + y\sqrt{a}.$$

# Reflection principles

In the literature there are many known results that compare different class groups. For example, we have

$$\mathrm{rk}_3\mathrm{Cl}(\mathbb{Q}(\sqrt{d})) \leq \mathrm{rk}_3\mathrm{Cl}(\mathbb{Q}(\sqrt{-3d})) \leq 1 + \mathrm{rk}_3\mathrm{Cl}(\mathbb{Q}(\sqrt{d})),$$

which is known as Scholz's reflection principle.

# Reflection principles

In the literature there are many known results that compare different class groups. For example, we have

$$\mathrm{rk}_3\mathrm{Cl}(\mathbb{Q}(\sqrt{d})) \leq \mathrm{rk}_3\mathrm{Cl}(\mathbb{Q}(\sqrt{-3d})) \leq 1 + \mathrm{rk}_3\mathrm{Cl}(\mathbb{Q}(\sqrt{d})),$$

which is known as Scholz's reflection principle.

The main algebraic result in Smith's work is in fact a reflection principle that compares  $\mathrm{Art}_m$  of  $2^m$  quadratic fields.

# Reflection principles

In the literature there are many known results that compare different class groups. For example, we have

$$\mathrm{rk}_3\mathrm{Cl}(\mathbb{Q}(\sqrt{d})) \leq \mathrm{rk}_3\mathrm{Cl}(\mathbb{Q}(\sqrt{-3d})) \leq 1 + \mathrm{rk}_3\mathrm{Cl}(\mathbb{Q}(\sqrt{d})),$$

which is known as Scholz's reflection principle.

The main algebraic result in Smith's work is in fact a reflection principle that compares  $\mathrm{Art}_m$  of  $2^m$  quadratic fields.

How can we find such reflection principles?

# Reflection principles

In the literature there are many known results that compare different class groups. For example, we have

$$\mathrm{rk}_3\mathrm{Cl}(\mathbb{Q}(\sqrt{d})) \leq \mathrm{rk}_3\mathrm{Cl}(\mathbb{Q}(\sqrt{-3d})) \leq 1 + \mathrm{rk}_3\mathrm{Cl}(\mathbb{Q}(\sqrt{d})),$$

which is known as Scholz's reflection principle.

The main algebraic result in Smith's work is in fact a reflection principle that compares  $\mathrm{Art}_m$  of  $2^m$  quadratic fields.

How can we find such reflection principles?

Smith's idea is to look for situations where the compositum of various Hilbert class fields is in some sense *small*.

# Intersections of Hilbert class fields

Take primes  $p_1, p_2, q_1, q_2$ . Now suppose that we have a degree 4 unramified, abelian extension of  $\mathbb{Q}(\sqrt{dp_iq_j})$  each lifting the character  $\chi_a$ .

# Intersections of Hilbert class fields

Take primes  $p_1, p_2, q_1, q_2$ . Now suppose that we have a degree 4 unramified, abelian extension of  $\mathbb{Q}(\sqrt{dp_iq_j})$  each lifting the character  $\chi_a$ .

Recall that we then get  $\alpha_{i,j} \in \mathbb{Q}(\sqrt{a})$  with

$$\text{Norm}_{\mathbb{Q}(\sqrt{a})/\mathbb{Q}}(\alpha_{i,j}) = \frac{dp_iq_j}{a} z_{i,j}^2.$$

# Intersections of Hilbert class fields

Take primes  $p_1, p_2, q_1, q_2$ . Now suppose that we have a degree 4 unramified, abelian extension of  $\mathbb{Q}(\sqrt{dp_iq_j})$  each lifting the character  $\chi_a$ .

Recall that we then get  $\alpha_{i,j} \in \mathbb{Q}(\sqrt{a})$  with

$$\text{Norm}_{\mathbb{Q}(\sqrt{a})/\mathbb{Q}}(\alpha_{i,j}) = \frac{dp_iq_j}{a} z_{i,j}^2.$$

Then we see that the norm of  $\alpha_{1,1}\alpha_{1,2}\alpha_{2,1}\alpha_{2,2}$  is a square.



# Intersections of Hilbert class fields

Take primes  $p_1, p_2, q_1, q_2$ . Now suppose that we have a degree 4 unramified, abelian extension of  $\mathbb{Q}(\sqrt{dp_iq_j})$  each lifting the character  $\chi_a$ .

Recall that we then get  $\alpha_{i,j} \in \mathbb{Q}(\sqrt{a})$  with

$$\text{Norm}_{\mathbb{Q}(\sqrt{a})/\mathbb{Q}}(\alpha_{i,j}) = \frac{dp_iq_j}{a} z_{i,j}^2.$$

Then we see that the norm of  $\alpha_{1,1}\alpha_{1,2}\alpha_{2,1}\alpha_{2,2}$  is a square.

In other words, part of  $H_2(\mathbb{Q}(\sqrt{dp_2q_2}))$  is contained in the other  $H_2(\mathbb{Q}(\sqrt{dp_iq_j}))$ .

# Intersections of Hilbert class fields

Take primes  $p_1, p_2, q_1, q_2$ . Now suppose that we have a degree 4 unramified, abelian extension of  $\mathbb{Q}(\sqrt{dp_iq_j})$  each lifting the character  $\chi_a$ .

Recall that we then get  $\alpha_{i,j} \in \mathbb{Q}(\sqrt{a})$  with

$$\text{Norm}_{\mathbb{Q}(\sqrt{a})/\mathbb{Q}}(\alpha_{i,j}) = \frac{dp_iq_j}{a} z_{i,j}^2.$$

Then we see that the norm of  $\alpha_{1,1}\alpha_{1,2}\alpha_{2,1}\alpha_{2,2}$  is a square.

In other words, part of  $H_2(\mathbb{Q}(\sqrt{dp_2q_2}))$  is contained in the other  $H_2(\mathbb{Q}(\sqrt{dp_1q_1}))$ . This implies

$$\text{Art}_{2,dp_1q_1}(b, \chi_a) + \text{Art}_{2,dp_1q_2}(b, \chi_a) + \text{Art}_{2,dp_2q_1}(b, \chi_a) + \text{Art}_{2,dp_2q_2}(b, \chi_a) = 0$$

for  $b \in 2\text{Cl}(\mathbb{Q}(\sqrt{dp_iq_j}))[4]$  a fixed divisor of  $d$ .

## Another reflection principle

With similar techniques, Smith proves another reflection principle

$$\begin{aligned} & \text{Art}_{2,dp_1q_1}(b, \chi_{ap_1}) + \text{Art}_{2,dp_1q_2}(b, \chi_{ap_1}) + \\ & \text{Art}_{2,dp_2q_1}(b, \chi_{ap_2}) + \text{Art}_{2,dp_2q_2}(b, \chi_{ap_2}) = \sum_{r|b} \text{Frob}_{K_{p_1p_2, q_1q_2}/\mathbb{Q}}(r). \end{aligned}$$

## Another reflection principle

With similar techniques, Smith proves another reflection principle

$$\begin{aligned} & \text{Art}_{2,dp_1q_1}(b, \chi_{ap_1}) + \text{Art}_{2,dp_1q_2}(b, \chi_{ap_1}) + \\ & \text{Art}_{2,dp_2q_1}(b, \chi_{ap_2}) + \text{Art}_{2,dp_2q_2}(b, \chi_{ap_2}) = \sum_{r|b} \text{Frob}_{K_{p_1p_2, q_1q_2}/\mathbb{Q}}(r). \end{aligned}$$

However, in the Pell setting, the first Rédei matrix is symmetric.

## Another reflection principle

With similar techniques, Smith proves another reflection principle

$$\begin{aligned} & \text{Art}_{2,dp_1q_1}(b, \chi_{ap_1}) + \text{Art}_{2,dp_1q_2}(b, \chi_{ap_1}) + \\ & \text{Art}_{2,dp_2q_1}(b, \chi_{ap_2}) + \text{Art}_{2,dp_2q_2}(b, \chi_{ap_2}) = \sum_{r|b} \text{Frob}_{K_{p_1p_2, q_1q_2}/\mathbb{Q}}(r). \end{aligned}$$

However, in the Pell setting, the first Rédei matrix is symmetric.

That means we have to compute the Artin pairing between  $a$  and  $\chi_a$ .

## Another reflection principle

With similar techniques, Smith proves another reflection principle

$$\begin{aligned} & \text{Art}_{2,dp_1q_1}(b, \chi_{ap_1}) + \text{Art}_{2,dp_1q_2}(b, \chi_{ap_1}) + \\ & \text{Art}_{2,dp_2q_1}(b, \chi_{ap_2}) + \text{Art}_{2,dp_2q_2}(b, \chi_{ap_2}) = \sum_{r|b} \text{Frob}_{K_{p_1p_2, q_1q_2}/\mathbb{Q}}(r). \end{aligned}$$

However, in the Pell setting, the first Rédei matrix is symmetric.

That means we have to compute the Artin pairing between  $a$  and  $\chi_a$ .

We also have to compute the Artin pairing between  $dp_iq_j$  and  $\chi_a$ .

## Another reflection principle

With similar techniques, Smith proves another reflection principle

$$\begin{aligned} & \text{Art}_{2,dp_1q_1}(b, \chi_{ap_1}) + \text{Art}_{2,dp_1q_2}(b, \chi_{ap_1}) + \\ & \text{Art}_{2,dp_2q_1}(b, \chi_{ap_2}) + \text{Art}_{2,dp_2q_2}(b, \chi_{ap_2}) = \sum_{r|b} \text{Frob}_{K_{p_1p_2, q_1q_2}/\mathbb{Q}}(r). \end{aligned}$$

However, in the Pell setting, the first Rédei matrix is symmetric.

That means we have to compute the Artin pairing between  $a$  and  $\chi_a$ .

We also have to compute the Artin pairing between  $dp_iq_j$  and  $\chi_a$ .

The above reflection principle is useless in both cases.

## Another reflection principle

With similar techniques, Smith proves another reflection principle

$$\begin{aligned} & \text{Art}_{2,dp_1q_1}(b, \chi_{ap_1}) + \text{Art}_{2,dp_1q_2}(b, \chi_{ap_1}) + \\ & \text{Art}_{2,dp_2q_1}(b, \chi_{ap_2}) + \text{Art}_{2,dp_2q_2}(b, \chi_{ap_2}) = \sum_{r|b} \text{Frob}_{K_{p_1p_2, q_1q_2}/\mathbb{Q}}(r). \end{aligned}$$

However, in the Pell setting, the first Rédei matrix is symmetric.

That means we have to compute the Artin pairing between  $a$  and  $\chi_a$ .

We also have to compute the Artin pairing between  $dp_iq_j$  and  $\chi_a$ .

The above reflection principle is useless in both cases.

We develop two new reflection principles. Unlike Smith's work, they make essential use of Hilbert reciprocity in multiquadratic fields.



## Bonus slide: new reflection principles

For the Artin pairing with  $dp_i q_j$  we have (following Smith's ideas)

$$\begin{aligned} & \text{Art}_{2, dp_1 q_1}(dp_1 q_1, \chi_{ap_1}) + \text{Art}_{2, dp_1 q_2}(dp_1 q_2, \chi_{ap_1}) + \\ & \text{Art}_{2, dp_2 q_1}(dp_2 q_1, \chi_{ap_2}) + \text{Art}_{2, dp_2 q_2}(dp_2 q_2, \chi_{ap_2}) = \text{Frob}_{K_{p_1 p_2, q_1 q_2} / \mathbb{Q}}(\infty). \end{aligned}$$

## Bonus slide: new reflection principles

For the Artin pairing with  $dp_i q_j$  we have (following Smith's ideas)

$$\begin{aligned} & \text{Art}_{2, dp_1 q_1}(dp_1 q_1, \chi_{ap_1}) + \text{Art}_{2, dp_1 q_2}(dp_1 q_2, \chi_{ap_1}) + \\ & \text{Art}_{2, dp_2 q_1}(dp_2 q_1, \chi_{ap_2}) + \text{Art}_{2, dp_2 q_2}(dp_2 q_2, \chi_{ap_2}) = \text{Frob}_{K_{p_1 p_2, q_1 q_2}/\mathbb{Q}}(\infty). \end{aligned}$$

Our reciprocity law shows that

$$\text{Frob}_{K_{p_1 p_2, q_1 q_2}/\mathbb{Q}}(\infty) = \text{Frob}_{K_{p_1 p_2, -1}/\mathbb{Q}}(q_1) + \text{Frob}_{K_{p_1 p_2, -1}/\mathbb{Q}}(q_2).$$

## Bonus slide: new reflection principles

For the Artin pairing with  $dp_i q_j$  we have (following Smith's ideas)

$$\begin{aligned} & \text{Art}_{2, dp_1 q_1}(dp_1 q_1, \chi_{ap_1}) + \text{Art}_{2, dp_1 q_2}(dp_1 q_2, \chi_{ap_1}) + \\ & \text{Art}_{2, dp_2 q_1}(dp_2 q_1, \chi_{ap_2}) + \text{Art}_{2, dp_2 q_2}(dp_2 q_2, \chi_{ap_2}) = \text{Frob}_{K_{p_1 p_2, q_1 q_2}/\mathbb{Q}}(\infty). \end{aligned}$$

Our reciprocity law shows that

$$\text{Frob}_{K_{p_1 p_2, q_1 q_2}/\mathbb{Q}}(\infty) = \text{Frob}_{K_{p_1 p_2, -1}/\mathbb{Q}}(q_1) + \text{Frob}_{K_{p_1 p_2, -1}/\mathbb{Q}}(q_2).$$

For the pairing between  $a$  and  $\chi_a$  we also develop a new reflection principle.

# Summary

- ▶ Steinhilber gave a conjecture for the density of  $d \in \mathcal{D}$  for which the negative Pell equation is soluble.

# Summary

- ▶ Steinhilber gave a conjecture for the density of  $d \in \mathcal{D}$  for which the negative Pell equation is soluble.
- ▶ Carlo Pagano and I recently proved Steinhilber's conjecture using techniques related to the Cohen–Lenstra–Gerth conjecture.

# Summary

- ▶ Steinhagen gave a conjecture for the density of  $d \in \mathcal{D}$  for which the negative Pell equation is soluble.
- ▶ Carlo Pagano and I recently proved Steinhagen's conjecture using techniques related to the Cohen–Lenstra–Gerth conjecture.
- ▶ The key innovations are two new reflection principles based on Hilbert reciprocity in multiquadratic fields.

- ▶ Steinhagen gave a conjecture for the density of  $d \in \mathcal{D}$  for which the negative Pell equation is soluble.
- ▶ Carlo Pagano and I recently proved Steinhagen's conjecture using techniques related to the Cohen–Lenstra–Gerth conjecture.
- ▶ The key innovations are two new reflection principles based on Hilbert reciprocity in multiquadratic fields.

Thank you for your attention!