

VANISHING HOMOLOGY OF PROJECTIVE HYPERSURFACES WITH 1-DIMENSIONAL SINGULARITIES

DIRK SIERSMA AND MIHAI TIBĂR

ABSTRACT. We introduce the vanishing homology of projective hypersurfaces. In case of one-dimensional singular locus, we prove its concentration in two dimensions and find formulas for the ranks of the nontrivial homology groups.

1. INTRODUCTION AND RESULTS

The homology of a projective hypersurface $V \subset \mathbb{P}^{n+1}$ is known for smooth V whereas only few results are available in the singular setting. The classical Lefschetz Hyperplane Theorem (LHT) yields that the inclusion of spaces induces an isomorphism:

$$(1.1) \quad H_k(V, \mathbb{Z}) \xrightarrow{\cong} H_k(\mathbb{P}^{n+1}, \mathbb{Z})$$

for $j < n$ and an epimorphism for $j = n$, independently on the singular locus $\text{Sing } V$. Since V is a CW-complex of dimension $2n$, the remaining task is to find the homology groups $H_k(V, \mathbb{Z})$ for $j \geq n$.

In case of smooth V all homology groups appear to be free and by Poincaré duality¹: $H_k(V, \mathbb{Z}) \cong H_k(\mathbb{P}^n, \mathbb{Z})$ if $k \neq n$ and the rank of $H_n(V, \mathbb{Z})$ follows from the Euler characteristic computation $\chi(V) = n + 2 - \frac{1}{d}[1 + (-1)^{n+1}(d-1)^{n+2}]$. Smooth projective complete intersections have been studied by Libgober and Wood [LW].

In case of isolated singularities, Dimca proved a significant result [Di1], [Di3, Thm. 4.3] that we shall comment in §2 in relation to a similar result provided by our method (Proposition 2.2), but whenever the singular locus has dimension ≥ 1 there seems to be no result yet.

Our paper focuses on the first unknown case, $\dim \text{Sing } V = 1$. We approach the singular surface V by comparing its integer homology to that of a smooth hypersurface of the same degree, as an intermediate step towards computing the homology of singular hypersurfaces. Inspired by the study of vanishing cycles, we introduce the "vanishing homology" of V as follows:

Definition 1.1. Let $f = 0$ be the defining equation of $V \subset \mathbb{P}^{n+1}$ as a reduced hypersurface, where $d = \deg f$. Consider the following one-parameter smoothing of degree d , $V_\varepsilon := \{f_\varepsilon = f + \varepsilon h_d = 0\}$, where h_d denotes a general homogeneous polynomial of degree d . Let:

$$\mathbb{V}_\Delta := \{(x, \varepsilon) \in \mathbb{P}^{n+1} \times \Delta \mid f + \varepsilon h_d = 0\}$$

2000 *Mathematics Subject Classification.* 32S30, 58K60, 55R55, 32S50.

¹see [Di1] for details

denote the total space of the pencil, where $V_0 := V \subset \mathbb{P}^{n+1} \times \{0\}$ and Δ is a small enough disk centered at $0 \in \mathbb{C}$ such that V_ε is nonsingular for all $\varepsilon \in \Delta^*$. Let $A := \{f = h_d = 0\}$ be the axis of the pencil and let $\pi : \mathbb{V}_\Delta \rightarrow \Delta$ denote the projection. We define:

$$H_*^Y(V) := H_*(\mathbb{V}_\Delta, V_\varepsilon; \mathbb{Z})$$

and call it the *vanishing homology of V* .

The genericity of h_d ensures the existence of small enough disks Δ as in the above definition, see e.g. [ST2, Prop. 2.2]. Note that \mathbb{V}_Δ retracts to V , thus the vanishing homology compares V to the smooth hypersurface V_ε of the same degree. Since all smooth hypersurfaces of fixed degree are homeomorphic, the vanishing homology does not depend on the particular smoothing of degree d , thus it is an invariant of V .

We obtain in this paper that the vanishing homology $H_*^Y(V)$ is concentrated in dimensions $n+1$ and $n+2$ only (**Theorem 4.1**). By the exact sequence of the pair $(\mathbb{V}_\Delta, V_\varepsilon)$, the concentration of the vanishing homology implies the isomorphisms:

$$H_k(V, \mathbb{Z}) \simeq H_k(\mathbb{P}^n, \mathbb{Z}) \text{ for } k \neq n, n+1, n+2.$$

Our main results in §6 are formulas for the ranks of the remaining groups $H_{n+1}^Y(V)$ and $H_{n+2}^Y(V)$. They depend on the information about local isolated or special non-isolated singularities, the properties of the singular curve $\text{Sing } V$, the transversal singularity types and the monodromies along loops in the transversal local systems. The singular locus $\text{Sing } V$ has a finite set R of isolated points and finitely many curve branches. Each such branch Σ_i of $\text{Sing } V$ has a generic transversal type (of transversal Milnor fiber F_i^{tr} and Milnor number denoted by μ_i^{tr}) and the axis A cuts it at a finite set of general points P_i . It also contains a finite set Q_i of points with non-generic transversal type, which we call *special points*, and we denote by \mathcal{A}_q the local Milnor fibre at $q \in Q$. At each point $q \in Q_i$ there are finitely many locally irreducible branches of the germ (Σ_i, q) , we denote by $\gamma_{i,q}$ their number and let $\gamma_i := \sum_{q \in Q_i} \gamma_{i,q}$ (see §4.1 for the notations). We then derive the following results.

Theorem 6.1: the $(n+2)$ th vanishing Betti number is bounded by the sum of all Milnor numbers of transversal singularities, taken over all irreducible 1-dimensional components of $\text{Sing } V$, and each special singular point on $\text{Sing } V$ with non-trivial transversal monodromy decreases this Betti-number.

Corollary 6.5 (see also Example 7.3): if for each irreducible 1-dimensional component Σ_i of $\text{Sing } V$ we have at least one local special singularity with rank zero $(n-1)$ th homology group, then the vanishing homology of V is free, concentrated in dimension $n+1$ only, and the corresponding Betti number is given by the formula:

$$b_{n+1}(\mathbb{V}_\Delta, V_\varepsilon) = \sum_i (\nu_i + \gamma_i + 2g_i - 2)\mu_i^{\text{tr}} + (-1)^n \sum_{q \in Q} \tilde{\chi}(\mathcal{A}_q) + \sum_{r \in R} \mu_r,$$

where $Q := \cup_i Q_i$, $\nu_i := \#P_i$, μ_r is the Milnor number of the isolated singularity germ (V, r) , and g_i is the genus of Σ_i (see §4.5 for the meaning of the genus in case of singular Σ_i) and $\tilde{\chi}$ denotes the reduced Euler characteristic.

In the proofs we use several devices, among which the detailed construction of a CW-complex model of the pair $(\mathbb{V}_\Delta, V_\varepsilon)$ done in §4.4 and §4.5; the Euler characteristic computation given in [ST2, Theorem 5.3]; the full strength of the results on local 1-dimensional

singularities found by Siersma [Si1], [Si2], [Si3], [Si4], cf also [Yo], [Ti3], which involve the study of the local system of transversal Milnor fibers along the singular locus.

We provide several examples in §7. In certain cases we can prove the freeness of the $(n + 1)$ th vanishing homology group. We also show an example where the homology of V over \mathbb{Q} can be computed via our formulas for the vanishing homology.

Acknowledgment. We thank the Centre International de Rencontres Mathématiques which hosted us within the framework “recherche en binôme” in Luminy (November 2013), where part of this project was accomplished.

2. VANISHING HOMOLOGY IN CASE OF ISOLATED SINGULARITIES

Throughout this paper we use only homology over \mathbb{Z} . Let $V := \{f = 0\} \subset \mathbb{P}^{n+1}$ be a hypersurface of degree d with singular locus consisting of a finite set of points R . Since V has only isolated singularities, the genericity of the axis $A = \{f = h_d = 0\}$ of the pencil $\pi : \mathbb{V}_\Delta \rightarrow \Delta$ just means that A avoids R . It turns out (see also [ST2, §5]) that \mathbb{V}_Δ is non-singular and that the projection π has isolated singularities precisely at the points of R . For small enough balls B_r , at each point $r \in R$, the homotopy retraction within the fibration π yields the isomorphism:

$$(2.1) \quad H_*(\mathbb{V}_\Delta, V_\varepsilon) \simeq \bigoplus_{r \in R} H_*(B_r, B_r \cap V_\varepsilon)$$

where $B_r \cap V_\varepsilon$ is the Milnor fiber of the isolated hypersurface singularity germ (V, r) . The relative homology $H_*(B_r, B_r \cap V_\varepsilon)$ is concentrated in $n + 1$ and $H_{n+1}(B_r, B_r \cap V_\varepsilon)$ is isomorphic to the Milnor lattice \mathbb{L}_r of the hypersurface germ (V, r) , thus isomorphic to \mathbb{Z}^{μ_r} , where μ_r is the Milnor number of (V, r) . We get the following conclusion:

Lemma 2.1. *If $\text{Sing } V \leq 0$ then:*

$$H_k^Y(V) = 0 \text{ if } k \neq n + 1,$$

$$H_{n+1}^Y(V) = \bigoplus_{r \in R} \mathbb{L}_r.$$

□

From the long exact sequence of the pair $(\mathbb{V}_\Delta, V_\varepsilon)$ we also obtain the 5-terms exact sequence:

$$0 \rightarrow H_{n+1}(V_\varepsilon) \rightarrow H_{n+1}(V) \rightarrow \bigoplus_{r \in R} \mathbb{L}_r \xrightarrow{\Phi_n} \mathbb{L} \rightarrow H_n(V) \rightarrow 0$$

where $\mathbb{L} := H_n(V_\varepsilon)$ is the intersection lattice of the middle homology of the smooth hypersurface of degree d and the map Φ_n is identified to the boundary map $H_{n+1}(\mathbb{V}_\Delta, V_\varepsilon) \rightarrow H_n(V_\varepsilon)$. We get the integer homology of V as follows:

- Proposition 2.2.** (a) $H_k(V) \simeq H_k(\mathbb{P}^n)$ for $k \neq n, n + 1$,
 (b) $H_{n+1}(V) \simeq H_{n+1}(\mathbb{P}^n) \oplus \ker \Phi_n$,
 (c) $H_n(V) \simeq \text{coker } \Phi_n$.

□

This is striking similar to Dimca's result [Di1, Theorem 2.1], [Di3, Theorem 5.4.3], although formulated and proved in different terms. As Dimca observed in [Di1], we also point out here that the relation between vanishing homology and absolute homology is encoded by the morphism Φ_n , which is difficult to identify from the equation of f .

3. LOCAL THEORY OF 1-DIMENSIONAL SINGULAR LOCUS

We shall need several facts from the local theory of singularities with a 1-dimensional singular set. We recall them here, following [Si3], see also the survey [Si4].

We consider a holomorphic function germ $f : (\mathbb{C}^{n+1}, 0) \rightarrow (\mathbb{C}, 0)$ with singular locus Σ of dimension 1. Let $\Sigma = \Sigma_1 \cup \dots \cup \Sigma_r$ be the decomposition into irreducible curve components. Let F be the local Milnor fiber of f . The homology $\tilde{H}_*(F)$ is concentrated in dimensions $n-1$ and n , namely $H_n(F) = \mathbb{Z}^{\mu_n}$, which is free, and $H_{n-1}(F)$ which can have torsion.

There is a well-defined local system on $\Sigma_i \setminus \{0\}$ having as fiber the homology of the transversal Milnor fiber $\tilde{H}_{n-1}(F_i^{\text{th}})$, i.e., F_i^{th} is the Milnor fiber of the restriction of f to the transversal hyperplane section at some $x \in \Sigma_i \setminus \{0\}$, which is an isolated singularity whose equisingularity class is independent of the point x . Thus $\tilde{H}_*(F_i^{\text{th}})$ is concentrated in dimension $n-1$. On this group there acts the *local system monodromy* (also called *vertical monodromy*):

$$A_i : \tilde{H}_{n-1}(F_i^{\text{th}}) \rightarrow \tilde{H}_{n-1}(F_i^{\text{th}}).$$

As explained in [Si3], one considers a tubular neighborhood $\mathcal{N} := \sqcup_{i=1}^r \mathcal{N}_i$ of the link of Σ and decomposes the boundary ∂F of the Milnor fiber as $\partial F = \partial_1 F \cup \partial_2 F$, where $\partial_2 F := \partial F \cap \mathcal{N}$. Then $\partial_2 F = \bigsqcup_{i=1}^r \partial_2 F_i$, where $\partial_2 F_i := \Sigma_i \cap \mathcal{N}_i$.

The homology groups of $\partial_2 F$ are related to the local system monodromies A_i in the following way. Each boundary component $\partial_2 F_i$ is fibered over the link of Σ_i with fiber F_i^{th} . The Wang sequence of this fibration yields the following non-trivial part, for $n \geq 3$:

$$(3.1) \quad 0 \rightarrow H_n(\partial_2 F_i) \rightarrow H_{n-1}(F_i^{\text{th}}) \xrightarrow{A_i - I} H_{n-1}(F_i^{\text{th}}) \rightarrow H_{n-1}(\partial_2 F_i) \rightarrow 0$$

In this sequence the following two homology groups play a crucial role: $H_n(\partial_2 F) = \bigoplus_{i=1}^r \text{Ker}(A_i - I)$ and $H_{n-1}(\partial_2 F) \cong \bigoplus_{i=1}^r \text{Coker}(A_i - I)$. The first group is free, the second can have torsion, and they are isomorphic up to torsion. For $n = 2$ there is an adapted interpretation of this sequence, cf [Si3, Section 6].

What we will actually need in the following is a relative version of this Wang sequence. Let E_i^{th} be the transversal Milnor neighborhood containing the transversal fiber F_i^{th} and let $\partial_2 E_i$ denote the total space of its fibration above the link of Σ_i . Therefore E_i^{th} is contractible and $\partial_2 E_i$ may be identified with the tubular neighborhood \mathcal{N}_i which retracts to the link of Σ_i . We then have:

Lemma 3.1. *For $n \geq 2$*

$$0 \rightarrow H_{n+1}(\partial_2 E_i, \partial_2 F_i) \rightarrow H_n(E_i^{\text{th}}, F_i^{\text{th}}) \xrightarrow{A_i - I} H_n(E_i^{\text{th}}, F_i^{\text{th}}) \rightarrow H_{n-1}(\partial_2 E_i, \partial_2 F_i) \rightarrow 0$$

is an exact sequence, and

$$\begin{aligned} H_{n+1}(\partial_2 E, \partial_2 F) &= \bigoplus_{i=1}^r \text{Ker}(A_i - I) \\ H_n(\partial_2 E, \partial_2 F) &\cong \bigoplus_{i=1}^r \text{Coker}(A_i - I) \end{aligned}$$

Proof. For $n > 2$ the statement follows immediately from the above Wang sequence (3.1) and the definitions of E_i^n and $\partial_2 E_i$. One observes that $n = 2$ is no longer a special case like it was in the absolute setting (see the remark after (3.1)). \square

The non-trivial part of the long exact sequence of the pair $(F, \partial_2 F)$ is the following 6-terms piece. More precisely, we need the following result:

Proposition 3.2. [Si3] *The sequence*

$0 \rightarrow H_{n+1}(F, \partial_2 F) \rightarrow H_n(\partial_2 F) \rightarrow H_n(F) \rightarrow H_n(F, \partial_2 F) \rightarrow H_{n-1}(\partial_2 F) \rightarrow H_{n-1}(F) \rightarrow 0$
is exact. Moreover

$$H_{n+1}(F, \partial_2 F) \cong H_{n-1}(F)^{\text{free}} \text{ and } H_n(F, \partial_2 F) \cong H_n(F) \oplus H_{n-1}(F)^{\text{torsion}}.$$

\square

Note that $H_n(\partial_2 F) = \bigoplus_{i=1}^r \text{Ker}(A_i - I)$ and $H_{n-1}(\partial_2 F) \cong \bigoplus_{i=1}^r \text{Coker}(A_i - I)$ play again a crucial role.

4. THE VANISHING NEIGHBOURHOOD OF THE PROJECTIVE HYPERSURFACE

We give here the necessary constructions and lemmas that we shall use in the proof of the announced vanishing theorem:

Theorem 4.1. *If $\dim \text{Sing } V \leq 1$ then $H_j^Y(V) = 0$ for all $j \neq n + 1, n + 2$.*

Let $V := \{f = 0\} \subset \mathbb{P}^{n+1}$ denote a hypersurface of degree d with singular locus $\hat{\Sigma}$ of dimension one, more precisely $\hat{\Sigma}$ consists of a union $\Sigma := \cup_i \Sigma_i \cup R$ of irreducible projective curves Σ_i and of a finite set of points R .

We recall that we have denoted by $A = \{f = h_d = 0\}$ the axis of the pencil $\pi : \mathbb{V}_\Delta \rightarrow \Delta$ defined in the Introduction. One considers the polar locus of the map $(h_d, f) : \mathbb{C}^{n+2} \rightarrow \mathbb{C}^2$ and since this is a homogeneous set one takes its image in \mathbb{P}^{n+1} which will be denoted by $\Gamma(h_d, f)$. Let us recall from [ST2] the meaning of ‘‘general’’ for h_d in this setting. By using the Veronese embedding of degree d we find a Zariski open set \mathcal{O} of linear functions in the target such that whenever $g \in \mathcal{O}$ then its pull-back is a general homogeneous polynomial h_d defining a hypersurface $H := \{h_d = 0\}$ which is transversal to V in the stratified sense, i.e. after endowing V with some Whitney stratification, of which the strata are as follows: the isolated singular points $\{\{r\}, r \in R\}$ of V and the point-strata $\{\{q\}, q \in Q\}$ in Σ , the components of $\Sigma \setminus Q$ and the open stratum $V \setminus \hat{\Sigma}$. Such h_d will be called *general*. This definition implies that A intersects $\hat{\Sigma}$ at general points, in particular does not contain any points of $Q \cup R$. It was shown in [ST2, Lemma 5.1] that the space \mathbb{V}_Δ has isolated singularities: $\text{Sing } \mathbb{V}_\Delta = (A \cap \Sigma) \times \{0\}$, and that $\pi : \mathbb{V}_\Delta \rightarrow \Delta$ is a map with 1-dimensional singular locus $\text{Sing}(\pi) = \hat{\Sigma} \times \{0\}$. One of the key preliminary results

is the following supplement to [ST2, Lemma 5.2], which extends the proof in *loc.cit.* from Euler characteristic to homology ²:

Lemma 4.2. *If h_d is general then $\Gamma_p(h_d, f) = \emptyset$ at any point $p \in A \times \{0\}$. In particular, for a small enough ball B_p centered at p , the local relative homology is trivial, i.e.:*

$$H_*(B_p, B_p \cap V_\varepsilon) = 0.$$

Proof. The notation B_p stands for the intersection of \mathbb{V}_Δ with a small ball in some chosen affine chart $\mathbb{C}^{n+1} \times \Delta$ of the ambient space $\mathbb{P}^{n+1} \times \Delta$. In particular B_p is of dimension $n + 1$. Consider the map $(\pi, h_d) : B_p \rightarrow \Delta \times \Delta'$. Consider the germ of the polar locus of this map at p , denoted by $\Gamma(\pi, \hat{h}_d)$, where \hat{h}_d is the de-homogenization of h_d in the chosen chart. It follows from the definition of the polar locus that some point $(x, \varepsilon) \in \mathbb{V}_\Delta$, where $\varepsilon = -f(x)/h_d(x)$, is contained in $\Gamma(\pi, h_d) \setminus (\{f = 0\} \cup \{h_d = 0\})$ if and only if $x \in \Gamma(f, h_d) \setminus (\{f = 0\} \cup \{h_d = 0\})$. By the first statement, $\Gamma(f, h_d)$ is empty at p . The absence of the polar locus implies that $B_p \cap V_\varepsilon$ is homotopy equivalent (by deformation retraction) to the space $B_p \cap V_\varepsilon \cap \{h_d = 0\}$. The latter is the slice by $\varepsilon = \text{constant}$ of the space $\mathbb{V}_\Delta \cap \{h_d = 0\} = \{f = 0\} \times \Delta$, which is a product space. Since this is homeomorphic to the complex link of this space and a product space has contractible complex link, we deduce that $B_p \cap V_\varepsilon$ is contractible too. Since B_p is contractible itself, we get our claim. \square

4.1. Notations. Let us assume for the moment that Σ is irreducible and discuss the reducible case at the end in §5.2. Let g be its genus, in the sense of the definition given at §4.5. We use the following notations:

$P := A \cap \Sigma$ the set of axis points of Σ ; $Q :=$ the set of special points on Σ ;

$R :=$ the set of isolated singular points.

$\Sigma^* := \Sigma \setminus (P \cup Q)$ and $\mathcal{Y} :=$ small enough tubular neighborhood of Σ^* .

B_p, B_q, B_r are small enough Milnor balls within $\mathbb{V}_\Delta \subset \mathbb{P}^{n+1} \times \Delta$ at the points $p \in P, q \in Q, r \in R$ respectively, and

$B_P := \sqcup_p B_p, B_Q := \sqcup_q B_q$ and $B_R := \sqcup_r B_r$.

Let us denote the projection of the tubular neighborhood by $\pi_\Sigma : \mathcal{Y} \rightarrow \Sigma^*$.

Let $\nu := \#P$ be the number of axis points. At any special point $q \in Q$, let S_q be the index set of locally irreducible branch of the germ (Σ, q) , and let $\gamma := \sum_{q \in Q} \#S_q$.

By homotopy retraction and by excision we have:

$$(4.1) \quad H_*(\mathbb{V}_\Delta, V_\varepsilon) \simeq H_*(\mathcal{Y} \cup B_P \cup B_Q, V_\varepsilon \cap \mathcal{Y} \cup B_P \cup B_Q) \oplus \bigoplus_{r \in R} H_*(B_r, V_\varepsilon \cap B_r).$$

We introduce the following shorter notations:

$$\mathcal{X} := B_P \sqcup B_Q, \mathcal{A} := V_\varepsilon \cap \mathcal{X}, \mathcal{B} := V_\varepsilon \cap \mathcal{Y}, \mathcal{Z} := \mathcal{X} \cap \mathcal{Y}, \mathcal{C} := \mathcal{A} \cap \mathcal{B}$$

$$(\mathcal{X}_p, \mathcal{A}_p) := (B_p, V_\varepsilon \cap B_p), (\mathcal{X}_q, \mathcal{A}_q) := (B_q, V_\varepsilon \cap B_q).$$

²A related result was obtained in [PP]. Like in case of [PP], the proof actually works for any singular locus $\text{Sing } V$ and any general pencil.

In the new notations, the first direct summand of (4.1) is $H_*(\mathcal{X} \cup \mathcal{Y}, \mathcal{A} \cup \mathcal{B})$, thus (4.1) writes as follows:

$$(4.2) \quad H_*(\mathbb{V}_\Delta, V_\varepsilon) \simeq H_*(\mathcal{X} \cup \mathcal{Y}, \mathcal{A} \cup \mathcal{B}) \oplus \bigoplus_{r \in R} H_*(B_r, V_\varepsilon \cap B_r).$$

Note that each direct summand $H_*(B_r, V_\varepsilon \cap B_r)$ is concentrated in dimension $n + 1$ since it identifies to the Milnor lattice of the isolated singularities germs (V_0, r) , where μ_r denotes its Milnor number. This aspect was treated in §1 in case of isolated singularities. We shall therefore deal from now on with the first term in the direct sum of (4.1).

We next consider the relative Mayer-Vietoris long exact sequence:

$$(4.3) \quad \cdots \rightarrow H_*(\mathcal{Z}, \mathcal{C}) \rightarrow H_*(\mathcal{X}, \mathcal{A}) \oplus H_*(\mathcal{Y}, \mathcal{B}) \rightarrow H_*(\mathcal{X} \cup \mathcal{Y}, \mathcal{A} \cup \mathcal{B}) \xrightarrow{\partial_s} \cdots$$

of the pair $(\mathcal{X} \cup \mathcal{Y}, \mathcal{A} \cup \mathcal{B})$ and we compute in the following each term of it.

4.2. The homology of $(\mathcal{X}, \mathcal{A})$. One has the direct sum decomposition $H_*(\mathcal{X}, \mathcal{A}) \simeq \bigoplus_p H_*(\mathcal{X}_p, \mathcal{A}_p) \oplus \bigoplus_q H_*(\mathcal{X}_q, \mathcal{A}_q)$ since \mathcal{X} is a disjoint union. The triviality $H_*(\mathcal{X}_p, \mathcal{A}_p) = 0$ follows by Lemma 4.2. The pairs $(\mathcal{X}_q, \mathcal{A}_q)$ are local Milnor data of the germs (V, q) with 1-dimensional singular locus and therefore the relative homology $H_*(\mathcal{X}_q, \mathcal{A}_q)$ is concentrated in dimensions n and $n + 1$.

4.3. The homology of $(\mathcal{Z}, \mathcal{C})$. The pair $(\mathcal{Z}, \mathcal{C})$ is a disjoint union of pairs localized at points $p \in P$ and $q \in Q$. For axis points $p \in P$ we have similarly a unique pair $(\mathcal{Z}_p, \mathcal{C}_p)$ as bundle over the link of Σ at p with fiber the transversal data $(E_p^\natural, F_p^\natural)$, in the notations of §3. For the non-axis points $q \in Q$ we have one contribution for each *locally irreducible branch of the germ* (Σ, q) . Let S_q be the index set of all these branches at $q \in Q$. We get the following decomposition:

$$(4.4) \quad H_*(\mathcal{Z}, \mathcal{C}) \simeq \bigoplus_{p \in P} H_*(\mathcal{Z}_p, \mathcal{C}_p) \oplus \bigoplus_{q \in Q} \bigoplus_{s \in S_q} H_*(\mathcal{Z}_s, \mathcal{C}_s).$$

More precisely, one such local pair $(\mathcal{Z}_s, \mathcal{C}_s)$ is the bundle over the corresponding component of the link of the curve germ Σ at q having as fiber the local transversal Milnor data $(E_s^\natural, F_s^\natural)$. In the notations of §3, we thus have: $\partial_2 \mathcal{A}_q = \sqcup_{s \in S_q} \mathcal{C}_s$.

The relative homology groups in the above decomposition (4.4) depend on the *vertical monodromy* via the Wang sequence of Lemma 3.1, as follows:

$$(4.5) \quad 0 \rightarrow H_{n+1}(\mathcal{Z}_s, \mathcal{C}_s) \rightarrow H_n(E^\natural, F^\natural) \xrightarrow{A_s - I} H_n(E^\natural, F^\natural) \rightarrow H_n(\mathcal{Z}_s, \mathcal{C}_s) \rightarrow 0.$$

Note that here the transversal data is independent of the points q or the index s since Σ^* is connected and therefore the transversal fiber is uniquely defined. However the vertical monodromies A_s depend on $s \in S_q$. From the above and from Lemma 4.2 we get:

Lemma 4.3. *At points $q \in Q$, for each $s \in S_q$ one has:*

$$\begin{aligned} H_k(\mathcal{Z}_s, \mathcal{C}_s) &= 0 & k \neq n, n + 1, \\ H_{n+1}(\mathcal{Z}_s, \mathcal{C}_s) &\cong \ker(A_s - I), & H_n(\mathcal{Z}_s, \mathcal{C}_s) &\cong \operatorname{coker}(A_s - I). \end{aligned}$$

At axis points $p \in P$ and more generally, at any point p such that $A_p = I$, one has:

$$\begin{aligned} H_k(\mathcal{Z}_p, \mathcal{C}_p) &= 0 & k \neq n, n + 1, \\ H_{n+1}(\mathcal{Z}_p, \mathcal{C}_p) &\cong H_n(\mathcal{Z}_p, \mathcal{C}_p) \cong H_n(E^\natural, F^\natural) = \mathbb{Z}^{\mu^\natural}. \end{aligned}$$

Proof. The first statement follows from the Wang sequence (4.5) and since $H_k(E^\natural, F^\natural)$ is concentrated in $k = n$. The last statement follows because at the axis points $p \in P$ the local system (similar to (4.5)) is trivial by Lemma 4.2 and therefore the vertical monodromy A_p is the identity. \square

We conclude that $H_*(\mathcal{Z}, \mathcal{C})$ is concentrated in dimensions n and $n + 1$ only.

4.4. The CW-complex structure of $(\mathcal{Z}, \mathcal{C})$. The pair $(\mathcal{Z}_s, \mathcal{C}_s)$ has moreover the following structure of a relative CW-complex, up to homotopy. Each bundle over some circle link can be obtained from a trivial bundle over an interval by identifying the fibers above the end points via the geometric vertical monodromy A_s . In order to obtain \mathcal{Z}_s from \mathcal{C}_s one can start by first attaching n -cells $c_1, \dots, c_{\mu^\natural}$ to the fiber F^\natural in order to kill the μ^\natural generators of $H_{n-1}(F^\natural)$ at the identified ends, and next by attaching $(n + 1)$ -cells $e_1, \dots, e_{\mu^\natural}$ to the preceding n -skeleton. The attaching of some $(n + 1)$ -cell is as follows: consider some n -cell a of the n -skeleton and take the cylinder $I \times a$ as an $(n + 1)$ -cell. Fix an orientation of the circle link, attach the base $\{0\} \times a$ over a , then follow the circle bundle in the fixed orientation by the monodromy A_s and attach the end $\{1\} \times a$ over $A_s(a)$. At the level of the cell complex, the boundary map of this attaching identifies to $A_s - I : \mathbb{Z}^{\mu^\natural} \rightarrow \mathbb{Z}^{\mu^\natural}$.

4.5. The CW-complex structure of $(\mathcal{Y}, \mathcal{B})$. For technical reasons we introduce one more puncture on Σ . Let us therefore define the total set of punctures $T := P \sqcup Q \sqcup \{y\}$, where y is a general point of Σ , then redefine $\Sigma^* := \Sigma \setminus T$ by considering the new puncture y . Moreover we use notations $(\mathcal{X}_y, \mathcal{A}_y)$ and $(\mathcal{Z}_y, \mathcal{C}_y)$.

Let $n : \tilde{\Sigma} \rightarrow \Sigma$ be the normalization map. Then we have the isomorphism $\Sigma^* = \Sigma \setminus T \simeq \tilde{\Sigma} \setminus n^{-1}(T)$. We choose generators of $\pi_1(\Sigma^*, z)$ for some base point $z \in \Sigma^*$ as follows: first the $2g$ loops (called *genus loops* in the following) which are generators of $\pi_1(\tilde{\Sigma}, n^{-1}(z))$, where g denotes the genus of the normalization $\tilde{\Sigma}$, and next by choosing one loop for each puncture of P and of Q . The total set of loops is indexed by the set $T' = T \setminus \{y\}$. Let us denote by W the set of indices for the union of T' with the genus loops, and therefore $\#W = 2g + \nu + \gamma$, where $\nu := \#P$ and $\gamma := \sum_{q \in Q} \#S_q$ (recall the Notations from §4.1). By enlarging “the hole” defined by the puncture y , we retract Σ^* to the chosen bouquet configuration of non-intersecting loops, denoted by Γ . The number of loops is $2g + \nu + \gamma$. Note that $\nu > 0$ since there must be at least d “axis points”.

The pair $(\mathcal{Y}, \mathcal{B})$ is then homotopy equivalent (by retraction) to the pair $(\pi_\Sigma^{-1}(\Gamma), \mathcal{B} \cap \pi_\Sigma^{-1}(\Gamma))$. We endow the latter with the structure of a relative CW-complex as we did with $(\mathcal{Z}, \mathcal{C})$ at §4.4, namely for each loop the similar CW-complex structure as we have defined above for some pair $(\mathcal{Z}_s, \mathcal{C}_s)$, see Figure 1.

The difference is that the pairs $(\mathcal{Z}_s, \mathcal{C}_s)$ are disjoint whereas in Σ^* the loops meet at a single point z . We thus take as reference the transversal fiber $F^\natural = \mathcal{B} \cap \pi_\Sigma^{-1}(z)$ above this point, namely we attach the n -cells (thimbles) only once to this single fiber in order to kill the μ^\natural generators of $H_{n-1}(F^\natural)$. The $(n + 1)$ -cells of $(\mathcal{Y}, \mathcal{B})$ correspond to the fiber bundles over the loops in the bouquet model of Σ^* . Over each loop, one attaches a number of μ^\natural $(n + 1)$ -cells to the fixed n -skeleton described before, more precisely one $(n + 1)$ -cell over

5. CONCENTRATION OF THE VANISHING HOMOLOGY. PROOF OF THEOREM 4.1

Lemma 4.3, §4.2 and Lemma 4.4 show that the terms $H_*(\mathcal{X}, \mathcal{A})$, $H_*(\mathcal{Y}, \mathcal{B})$ and $H_*(\mathcal{Z}, \mathcal{C})$ of the Mayer-Vietoris sequence (4.3) are concentrated only in dimensions n and $n + 1$, which fact implies the following result:

Proposition 5.1. *The relative Mayer Vietoris sequence (4.3) is trivial except for the following 7-terms sequence:*

$$(5.1) \quad \begin{aligned} 0 &\rightarrow H_{n+2}(\mathcal{X} \cup \mathcal{Y}, \mathcal{A} \cup \mathcal{B}) \rightarrow \\ &\rightarrow H_{n+1}(\mathcal{Z}, \mathcal{C}) \rightarrow H_{n+1}(\mathcal{X}, \mathcal{A}) \oplus H_{n+1}(\mathcal{Y}, \mathcal{B}) \rightarrow H_{n+1}(\mathcal{X} \cup \mathcal{Y}, \mathcal{A} \cup \mathcal{B}) \rightarrow \\ &\rightarrow H_n(\mathcal{Z}, \mathcal{C}) \xrightarrow{j} H_n(\mathcal{X}, \mathcal{A}) \oplus H_n(\mathcal{Y}, \mathcal{B}) \rightarrow H_n(\mathcal{X} \cup \mathcal{Y}, \mathcal{A} \cup \mathcal{B}) \rightarrow 0. \end{aligned}$$

□

From Proposition 5.1 and from (4.2) it follows that the vanishing homology $H_*(\mathbb{V}_\Delta, V_\varepsilon)$ is concentrated in dimensions $n, n + 1, n + 2$.

We pursue by showing that $H_n(\mathbb{V}_\Delta, V_\varepsilon) = 0$, i.e. that the last term of (5.1) is zero. We need the relative version of the exact sequence of Proposition 3.2, which appears to have an important overlap with our relative Mayer-Vietoris sequence.

Proposition 5.2. *For some point $q \in Q$, the sequence*

$$\begin{aligned} 0 &\rightarrow H_{n+1}(\mathcal{A}_q, \partial_2 \mathcal{A}_q) \rightarrow \bigoplus_{s \in S_q} H_{n+1}(\mathcal{Z}_s, \mathcal{C}_s) \rightarrow H_{n+1}(\mathcal{X}_q, \mathcal{A}_q) \rightarrow \\ &\rightarrow H_n(\mathcal{A}_q, \partial_2 \mathcal{A}_q) \rightarrow \bigoplus_{s \in S_q} H_n(\mathcal{Z}_s, \mathcal{C}_s) \rightarrow H_n(\mathcal{X}_q, \mathcal{A}_q) \rightarrow 0 \end{aligned}$$

is exact for $n \geq 2$. Moreover we have:

$$H_{n+1}(\mathcal{A}_q, \partial_2 \mathcal{A}_q) \cong H_{n-1}(\mathcal{A}_q)^{free} \text{ and } H_n(\mathcal{A}_q, \partial_2 \mathcal{A}_q) \cong H_n(\mathcal{A}_q) \oplus H_{n-1}(\mathcal{A}_q)^{torsion}$$

Proof. Note that we have the following coincidence of objects which have different notations in the projective setting of this section and in the local setting of §3: $\mathcal{A}_q := F$, $\partial_2 \mathcal{A}_q := \partial_2 F$.

We also have the isomorphisms $H_{*+1}(\mathcal{X}_q, \mathcal{A}_q) = \tilde{H}_*(\mathcal{A}_q)$ since \mathcal{X}_q is contractible, then $H_*(\partial_2 \mathcal{A}_q) = \bigoplus_{s \in S_q} H_*(\mathcal{C}_s)$ by definition, and $H_k(\mathcal{C}_s) = H_{k+1}(\mathcal{Z}_s, \mathcal{C}_s)$ for $k > 2$, since \mathcal{Z}_s contracts to a circle. We use Proposition 3.2 and check that, like in Lemma 3.1 on another (but similar) relative situation, the case $n = 2$ does not give any problem for the exactness of the above sequence. □

5.1. Surjectivity of j . We focus on the following map which occurs in the 7-term exact sequence (5.1):

$$(5.2) \quad j = j_1 \oplus j_2 : H_n(\mathcal{Z}, \mathcal{C}) \rightarrow H_n(\mathcal{X}, \mathcal{A}) \oplus H_n(\mathcal{Y}, \mathcal{B}).$$

5.1.1. *The first component $j_1 : H_n(\mathcal{Z}, \mathcal{C}) \rightarrow H_n(\mathcal{X}, \mathcal{A})$.*

Note that, as shown above, we have the following direct sum decompositions of the source and the target:

$$\begin{aligned} H_n(\mathcal{Z}, \mathcal{C}) &= \bigoplus_{p \in P} H_n(\mathcal{Z}_p, \mathcal{C}_p) \oplus_{q \in Q} \bigoplus_{s \in S_q} H_n(\mathcal{Z}_s, \mathcal{C}_s) \oplus H_n(\mathcal{Z}_y, \mathcal{C}_y), \\ H_n(\mathcal{X}, \mathcal{A}) &= \bigoplus_{q \in Q} H_n(\mathcal{X}_q, \mathcal{A}_q) \oplus H_n(\mathcal{X}_y, \mathcal{A}_y). \end{aligned}$$

The terms corresponding to the points $p \in P$ are mapped by j_1 to zero since $H_n(\mathcal{X}_p, \mathcal{A}_p) = 0$ by Lemma 4.2. Next, as shown in Proposition 5.2, at the special points $q \in Q$ we have

surjections: $\bigoplus_{s \in S_q} H_n(\mathcal{Z}_s, \mathcal{C}_s) \rightarrow H_n(\mathcal{X}_q, \mathcal{A}_q)$ and moreover $H_n(\mathcal{Z}_y, \mathcal{C}_y) \rightarrow H_n(\mathcal{X}_y, \mathcal{A}_y)$ is an isomorphism. We conclude to the surjectivity of the morphism j_1 .

5.1.2. *The second component $j_2 : H_n(\mathcal{Z}, \mathcal{C}) \rightarrow H_n(\mathcal{Y}, \mathcal{B})$.*

Both sides are described with a relative CW-complex as explained in §4.5. At the level of n -cells there are μ^{th} n -cell generators for each $p \in P$, and the same for each $s \in S_q$ and any $q \in Q$. Each of these generators is mapped bijectively to the single cluster of n -cell generators attached to the reference fiber F^{th} (which is the fiber above the common point of the loops, see also Figure 1). We have the same boundary map for each axis point $p \in P$ in the source and in the target of j_2 and therefore, at the level of the n -homology, the restriction $j_{2|} : H_n(\mathcal{Z}_p, \mathcal{C}_p) \rightarrow H_n(\mathcal{Y}, \mathcal{B})$ is surjective. Since we have at least one axis point on Σ and $\bigoplus_{p \in P} H_n(\mathcal{Z}_p, \mathcal{C}_p) \subset \ker j_1$, this shows that the restriction $j_{2|} : \bigoplus_{p \in P} H_n(\mathcal{Z}_p, \mathcal{C}_p) \rightarrow H_n(\mathcal{Y}, \mathcal{B})$ is surjective too. We have thus proven the surjectivity of j and in particular the following statement:

Proposition 5.3. *$H_n(\mathbb{V}_\Delta, V_\varepsilon) = 0$ and in particular the relative Mayer Vietoris sequence (5.1) reduces to the 6-terms sequence:*

$$(5.3) \quad \begin{aligned} 0 &\rightarrow H_{n+2}(\mathcal{X} \cup \mathcal{Y}, \mathcal{A} \cup \mathcal{B}) \rightarrow H_{n+1}(\mathcal{Z}, \mathcal{C}) \rightarrow H_{n+1}(\mathcal{X}, \mathcal{A}) \oplus H_{n+1}(\mathcal{Y}, \mathcal{B}) \\ &\rightarrow H_{n+1}(\mathcal{X} \cup \mathcal{Y}, \mathcal{A} \cup \mathcal{B}) \rightarrow H_n(\mathcal{Z}, \mathcal{C}) \xrightarrow{j} H_n(\mathcal{X}, \mathcal{A}) \oplus H_n(\mathcal{Y}, \mathcal{B}) \rightarrow 0 \end{aligned}$$

This shows that the relative homology $H_*(\mathbb{V}_\Delta, V_\varepsilon)$ is concentrated at the levels $n + 1$ and $n + 2$, and thus finishes the proof of Theorem 4.1 in case of irreducible Σ .

5.2. **Reducible Σ .** Let $\Sigma = \Sigma_1 \cup \dots \cup \Sigma_\rho$ be the decomposition into irreducible components. The proof of Theorem 4.1 in the reducible case remains the same modulo the following small changes and additional notations:

- (a) For each i one considers the set Q_i of special singular points of Σ_i . The points of intersection $\Sigma_{i_1} \cap \Sigma_{i_2}$ for $i_1 \neq i_2$ are considered as special points of both sets Q_{i_1} and Q_{i_2} , and therefore the union $Q := \bigcup_i Q_i$ is not disjoint. For some $q \in \Sigma_{i_1} \cap \Sigma_{i_2}$, the set of indices S_q runs over all the local irreducible components of the curve germ (Σ, q) . Nevertheless, when we are counting the local irreducible branches at some point $q \in Q_i$ on a specified component Σ_i then the set S_q will tacitly mean only those local branches of Σ_i at q .
- (b) The pair $(\mathcal{Y}, \mathcal{B})$ is a disjoint union and its homology decomposes accordingly, namely $H_*(\mathcal{Y}, \mathcal{B}) = \bigoplus_{1 \leq i \leq \rho} H_*(\mathcal{Y}_i, \mathcal{B}_i)$.
- (c) For each component Σ_i one has its transversal Milnor fiber denoted by F_i^{th} and its transversal Milnor number μ_i^{th} .

6. BETTI NUMBERS OF HYPERSURFACES WITH 1-DIMENSIONAL SINGULAR LOCUS

By Theorem 4.1, the vanishing homology of a hypersurface $V \subset \mathbb{P}^{n+1}$ with 1-dimensional singularities is concentrated in dimensions $n + 1$ and $n + 2$. We show that its $(n + 2)$ th vanishing homology group depends on the local data of the special points Q and on genus loop monodromies along the singular branches. We study this dependance in more detail, we find a formula for the rank of the free group $H_{n+2}(\mathbb{V}_\Delta, V_\varepsilon)$, and discover mild conditions which insure the vanishing of this group.

We continue to use the notations of §4. Let us especially recall the notations from §4.5 adapted here to the general setting of a reducible singular locus $\Sigma = \cup_{i=1}^{\rho} \Sigma_i$. For any $1 \leq i \leq \rho$, $\Sigma_i^* = \Sigma_i \setminus (P_i \sqcup Q_i \sqcup \{y_i\})$ retracts to a bouquet Γ_i of $2g_i + \nu_i + \gamma_i$ circles, where g_i denotes the genus of the normalization $\tilde{\Sigma}_i$, where $\nu_i := \#P_i$ is the number of axis points $A \cap \Sigma_i$, where $\gamma_i := \sum_{q \in Q_i} \#S_q$ and Q_i denotes the set of special points of Σ_i , the set S_q is indexing the local branches of Σ_i at q , and where $y_i \in \Sigma_i$ is some point not in the set $P_i \cup Q_i$. We denote by G_i the set of genus loops of Γ_i .

By Proposition 5.1 we have $H_{n+2}(\mathbb{V}_{\Delta}, V_{\varepsilon}) = \ker j = \ker[j_1 \oplus j_2]$, where

$$j_1 \oplus j_2 : H_{n+1}(\mathcal{Z}, \mathcal{C}) \rightarrow H_{n+1}(\mathcal{X}, \mathcal{A}) \oplus H_{n+1}(\mathcal{Y}, \mathcal{B}).$$

The main idea in this section is to embed $H_{n+2}(\mathbb{V}_{\Delta}, V_{\varepsilon})$ into the module $\mathbb{D} = \oplus_{i=1}^{\rho} \mathbb{D}_i$, where \mathbb{D}_i is the image of the diagonal map:

$$\Delta_*^i : H_n(E_i^{\text{rh}}, F_i^{\text{rh}}) \rightarrow \oplus_{q \in Q_i} \oplus_{s \in S_q} H_n(E_i^{\text{rh}}, F_i^{\text{rh}}), \quad a \mapsto (a, a, \dots, a).$$

The source and the target of $j_1 \oplus j_2$ have a direct sum decomposition at level $n+1$, like has been discussed at §5.1 for the n -th homology groups³:

$$(6.1) \quad j_1 \oplus j_2 : \oplus_{p \in P} H_{n+1}(\mathcal{Z}_p, \mathcal{C}_p) \oplus_{q \in Q} \oplus_{s \in S_q} H_{n+1}(\mathcal{Z}_s, \mathcal{C}_s) \oplus_{i=1}^{\rho} H_{n+1}(\mathcal{Z}_{y_i}, \mathcal{C}_{y_i}) \rightarrow \\ \oplus_{q \in Q} H_{n+1}(\mathcal{X}_q, \mathcal{A}_q) \bigoplus H_{n+1}(\mathcal{Y}, \mathcal{B}).$$

By Lemma 4.3 we have $H_{n+1}(\mathcal{Z}_v, \mathcal{C}_v) = \ker(A_v - I)$, where:

$$A_v - I : H_n(E_i^{\text{rh}}, F_i^{\text{rh}}) \rightarrow H_n(E_i^{\text{rh}}, F_i^{\text{rh}})$$

is the vertical monodromy at some point $v \in P_i$, or $v \in S_q$ and $q \in Q_i$, or $v = y_i$. The left hand side of (6.1) consists therefore of local contributions of the form $\ker(A_v - I) \subset H_n(E_i^{\text{rh}}, F_i^{\text{rh}}) \simeq H_{n-1}(F_i^{\text{rh}}) \simeq \mathbb{Z}^{\mu_i^{\text{rh}}}$.

We have studied j_1 in §5.1.1 at the level n . For the $(n+1)$ th homology groups, the restriction of j_1 to the first summand in (6.1) is zero since its image is in $\oplus_{p \in P} H_{n+1}(\mathcal{X}_p, \mathcal{A}_p)$ which is zero by Lemma 4.2. The image by j_1 of $\oplus_{i=1}^{\rho} H_{n+1}(\mathcal{Z}_{y_i}, \mathcal{C}_{y_i})$ is also zero by considering the local 6-term sequence from proposition 3.2. The restriction of j_1 to the remaining summand is the direct sum $\oplus_{q \in Q} j_{1,q}$ of the maps:

$$j_{1,q} : \bigoplus_{s \in S_q} H_{n+1}(\mathcal{Z}_s, \mathcal{C}_s) \rightarrow H_{n+1}(\mathcal{X}_q, \mathcal{A}_q).$$

The kernel of some $j_{1,q}$ is equal to $H_{n+1}(\mathcal{A}_q, \partial_2 \mathcal{A}_q)$, where \mathcal{A}_q is the local Milnor fiber of the hypersurface germ (V, q) , $q \in Q$, and is identified in Proposition 5.2 to the free part of $H_{n-1}(\mathcal{A}_q)$. Since $H_{n+1}(\mathcal{A}_q, \partial_2 \mathcal{A}_q)$ is contained in $\oplus_{i=1}^{\rho} \oplus_{Q_i \ni q} \oplus_{s \in S_q} H_{n+1}(\mathcal{Z}_s, \mathcal{C}_s)$, it turns out that the intersection $(\oplus_{i=1}^{\rho} \mathbb{D}_i) \cap \oplus_{q \in Q} H_{n+1}(\mathcal{A}_q, \partial_2 \mathcal{A}_q)$ is well defined.

Theorem 6.1. *In the above notations we have:*

$$H_{n+2}^{\vee}(V) = (\mathbb{D} \cap \oplus_{q \in Q} H_{n+1}(\mathcal{A}_q, \partial_2 \mathcal{A}_q)) \cap \oplus_{i=1}^{\rho} \Delta_*^i \left(\bigcap_{j \in G_i} \ker(A_j - I) \right)$$

where $A_j : H_n(E_i^{\text{rh}}, F_i^{\text{rh}}) \rightarrow H_n(E_i^{\text{rh}}, F_i^{\text{rh}})$ denotes the monodromy along the loop of Γ_i indexed by $j \in G_i$.

³we remind from §5.2 that the notation S_q depends of whether the point q is considered in Q or in Q_i , namely it takes either the local branches of Σ at q , or the local branches of Σ_i at q , accordingly.

In particular $H_{n+2}^\vee(V)$ is free and its rank is bounded as follows⁴:

$$\text{rank } H_{n+2}^\vee(V) \leq \sum_{i=1}^{\rho} \min_{s \in S_q, q \in Q_i, j \in G_i} \{ \dim \ker(A_s - I), \dim \ker(A_j - I) \} \leq \sum_{i=1}^{\rho} \mu_i^{\text{th}}.$$

Proof. In order to handle the map j_2 , we recall the relative CW-complex structure of $(\mathcal{Y}, \mathcal{B})$ given in §4.5. On each component Γ_i we have identified the set of points T_i which consists of the axis points P_i , the special points Q_i , and one general point y_i . The punctured Σ_i^* retracts to a configuration Γ_i of $2g_i + \nu_i + \gamma_i$ loops indexed by the set W_i , based at some point z_i , where $2g_i$ of them are “genus loops” and the other loops are projections by the normalization map $n_i : \tilde{\Sigma}_i \rightarrow \Sigma_i$ of loops around all the punctures of $\tilde{\Sigma}_i \setminus n_i^{-1}(P_i \sqcup Q_i)$. Notice that $\#T_i - 1 \geq \nu_i > 0$.

Let $\Gamma := \sqcup_i \Gamma_i$. Consider the spaces $\mathcal{Y}_\Gamma := \pi_\Sigma^{-1}(\Gamma)$ and $\mathcal{B}_\Gamma := \mathcal{B} \cap \mathcal{Y}_\Gamma$. We have the homotopy equivalence of pairs $(\mathcal{Y}, \mathcal{B}) \simeq (\mathcal{Y}_\Gamma, \mathcal{B}_\Gamma)$ which has been discussed at §4.5 and use the CW-complex model for $(\mathcal{Y}_\Gamma, \mathcal{B}_\Gamma)$. We also have the decomposition $(\mathcal{Y}, \mathcal{B}) = \sqcup_{i=1}^{\rho} (\mathcal{Y}_i, \mathcal{B}_i)$ according to the components Γ_i .

In our representation, the map j_2 splits into the direct sum of the following maps, for $i \in \{1, \dots, \rho\}$:

$$j_{2,i} : \oplus_{p \in P_i} H_{n+1}(\mathcal{Z}_p, \mathcal{C}_p) \oplus_{q \in Q_i} \oplus_{s \in S_q} H_{n+1}(\mathcal{Z}_s, \mathcal{C}_s) \oplus H_{n+1}(\mathcal{Z}_{y_i}, \mathcal{C}_{y_i}) \rightarrow H_{n+1}(\mathcal{Y}_i, \mathcal{B}_i).$$

By Lemma 4.4, the map $j_{2,i}$ restricts to an embedding of the direct sum $\oplus_{p \in P_i} H_{n+1}(\mathcal{Z}_p, \mathcal{C}_p) \oplus_{q \in Q_i} \oplus_{s \in S_q} H_{n+1}(\mathcal{Z}_s, \mathcal{C}_s)$ into $H_{n+1}(\mathcal{Y}_i, \mathcal{B}_i)$. Note that $H_{n+1}(\mathcal{Z}_v, \mathcal{C}_v) = \ker(A_v - I) \subset H_n(E_i^{\text{th}}, F_i^{\text{th}}) \simeq H_{n-1}(F_i^{\text{th}})$ for any point $v \in P_i$ or $v \in S_q$ and $q \in Q_i$. The kernel $\ker j_{2,i}$ is therefore determined by the relations induced by the image of the remaining direct summand $H_{n+1}(\mathcal{Z}_{y_i}, \mathcal{C}_{y_i})$ into $H_{n+1}(\mathcal{Y}_i, \mathcal{B}_i)$.

More precisely, each $(n+1)$ -cycle generator w of $H_{n+1}(\mathcal{Z}_{y_i}, \mathcal{C}_{y_i}) \simeq H_n(E_i^{\text{th}}, F_i^{\text{th}}) \simeq H_{n-1}(F_i^{\text{th}})$ induces one single relation. Namely $j_2(w)$ is a $(n+1)$ -cycle above the loop around the point y_i , and since this loop is homotopy equivalent to a certain composition of other loops of Γ_i , it follows that $j_2(w)$ is precisely homologous to the corresponding sum of cycles above the loops in Γ_i . Our scope is to find all such sums which contain as terms only elements from the images $j_2(H_{n+1}(\mathcal{Z}_p, \mathcal{C}_p))$ for $p \in P_i$ and $j_2(H_{n+1}(\mathcal{Z}_s, \mathcal{C}_s))$ for $s \in S_q$ and $q \in Q_i$. We have the following facts:

- 1). By Lemma 4.3 and §4.4, such images are in the kernels of $A - I$ where A is the vertical monodromy of the loop corresponding to $p \in P_i$ or to $s \in S_q$ and $q \in Q_i$. Therefore the expression of $j_2(w)$ contains the sum of those generators of $j_2(H_{n+1}(\mathcal{Z}_p, \mathcal{C}_p))$ and of $j_2(H_{n+1}(\mathcal{Z}_s, \mathcal{C}_s))$ which correspond to the same representative $w \in H_{n-1}(F_i^{\text{th}})$, for any $p \in P_i$ and any $s \in S_q$ and $q \in Q_i$. This implies that $w \in \cap_{s \in S_q, q \in Q_i} \ker(A_s - I)$. Note that the points $p \in P_i$ are superfluous in this intersection since $A_p = I$ for all such points.
- 2). Let us consider a pair γ_1 and γ_2 of genus loops (whenever $g_i > 0$) and let us denote by B_1 and B_2 the local system monodromy along these loops. The relation produced by $j_2(w)$ contains in principle the following relative cycle along the wedge $\gamma_1 \vee \gamma_2$: it starts from the representative $a_w \in H_{n-1}(F_i^{\text{th}})$ of w , moves in the local system along γ_1 arriving as $B_1(a_w)$ after one loop at the fiber over the base point z , next moved along γ_2 to $B_2 B_1(a_w)$, then in the opposite direction along γ_1 to $B_1^{-1} B_2 B_1(a_w)$ and finally in the opposite direction

⁴note that no multiplicities but only transversal types are involved in the rank formula.

along γ_2 to $B_2^{-1}B_1^{-1}B_2B_1(a_w)$. Our condition tells that the relation produced by $j_2(w)$ does not involve $(n+1)$ -cycles along the genus loops since $\text{Im } j_2 \cap \bigoplus_{j \in G_i} H_{n+1}(E_j^{\text{th}}, F^{\text{th}}) = 0$, by Lemma 4.4 and (4.6). Therefore the relative cycles along γ_1 and along γ_2 must cancel, which fact amounts to the following two pairs of equalities:

$$\begin{aligned} B_1^{-1}B_2B_1(a_w) &= a_w & \text{and} & & B_2B_1(a_w) &= B_1(a_w), \\ B_2^{-1}B_1^{-1}B_2B_1(a_w) &= B_1(a_w) & \text{and} & & B_1^{-1}B_2B_1(a_w) &= B_2B_1(a_w). \end{aligned}$$

These equalities are cyclic, thus the 8 above terms appear to be equal. In particular we get $B_1(a_w) = a_w$ and $B_2(a_w) = (a_w)$ for any $w \in \bigcap_{s \in S_q, q \in Q_i} \ker(A_s - I)$. We conclude to the same equalities for any pair of genus loops.

Altogether we obtain the following diagonal presentation of $\ker j_{2,i}$:

$$\ker j_{2,i} = \left\{ (a_w, a_w, \dots, a_w) \in \bigoplus_{q \in Q_i} \bigoplus_{s \in S_q} H_{n+1}(\mathcal{Z}_s, \mathcal{C}_s) \oplus H_{n+1}(\mathcal{Z}_{y_i}, \mathcal{C}_{y_i}) \mid w \in \bigcap_{s \in S_q, q \in Q_i} \ker(A_s - I) \cap \bigcap_{j \in G_i} \ker(A_j - I) \right\} \subset \mathbb{D}_i.$$

Since $H_{n+2}(\mathbb{V}_\Delta, V_\varepsilon) \subset \ker j_2 = \bigoplus_{i=1}^\rho \Delta_*^i \left(\bigcap_{s \in S_q, q \in Q_i} \ker(A_s - I) \cap \bigcap_{j \in G_i} \ker(A_j - I) \right)$ we get in particular the claimed inequality for the Betti number $b_{n+2}(\mathbb{V}_\Delta, V_\varepsilon)$. The freeness of $H_{n+2}(\mathbb{V}_\Delta, V_\varepsilon)$ follows from the fact that $\ker j_2$ is free (as the image of the intersection of free \mathbb{Z} -submodules).

We also obtain the desired expression of $H_{n+2}(\mathbb{V}_\Delta, V_\varepsilon) = \ker(j_1 \oplus j_2) = \ker j_1 \cap \ker j_2$ by intersecting $\ker j_2$ with the diagonal expression of $\ker j_1$ given just before the statement of Theorem 6.1. \square

REMARK 6.2. Irreducible Σ .

In case Σ is irreducible, the equality of Theorem 6.1 reads:

$$H_{n+2}^\vee(V) = \bigoplus_{q \in Q} H_{n+1}(\mathcal{A}_q, \partial_2 \mathcal{A}_q) \cap \bigcap_{j \in G} \ker(A_j - I).$$

In particular, if there are no special points on Σ and the monodromy along every the genus loop is the identity, then $H_{n+2}^\vee(V) \simeq H_{n-1}(F^{\text{th}})$. This situation can be seen in the example $V := \{xy = 0\} \subset \mathbb{P}^3$ for which $H_4^\vee(V) \simeq \mathbb{Z}$ and $\text{rank } H_3^\vee(V) = 1$.

REMARK 6.3. $(n+1)$ th vanishing Betti number.

It appears that $H_{n+2}^\vee(V)$ does not depend neither on the axis points, nor on the isolated singular points of V . However $H_{n+1}^\vee(V)$ depends on those elements since the Euler number does, after [ST2, Theorem 5.3]:

$$(6.2) \quad \chi(\mathbb{V}_\Delta, V_\varepsilon) = (-1)^{n+1} \sum_{i=1}^\rho (2g_i + \nu_i + \gamma_i - 2) \mu_i^{\text{th}} - \sum_{q \in Q} \tilde{\chi}(\mathcal{A}_q) + (-1)^{n+1} \sum_{r \in R} \mu_r.$$

where $\tilde{\chi}(\mathcal{A}_q) = \chi(\mathcal{A}_q) - 1$.

Theorem 6.1 is useful when we have information about the transversal monodromies, namely about the eigenspaces corresponding to the eigenvalue 1. We immediately derive:

Corollary 6.4. *If, for every $i \in \{1, \dots, \rho\}$, at least one of the transversal monodromies along the loops $\Gamma_i \subset \Sigma_i$ has no eigenvalue 1, then $H_{n+2}^\vee(V) = 0$. \square*

We may also apply Theorem 6.1 when we have enough information about local Milnor fibers of special points, like in the following case (see also Example 7.3):

Corollary 6.5. *Assume that for any $i \in \{1, \dots, \rho\}$ there is some special point $q_i \in Q$ such that the $(n-1)^{\text{th}}$ homology group of the local Milnor fiber \mathcal{A}_{q_i} of the hypersurface germ (V, q_i) has rank zero. Then:*

$$H_{n+2}^\vee(V) = 0$$

and the single non-zero vanishing Betti number $b_{n+1}^\vee(V)$ is given by the formula:

$$(6.3) \quad \text{rank } H_{n+1}^\vee(V) = \sum_i (\nu_i + \gamma_i + 2g_i - 2)\mu_i^\natural + (-1)^n \sum_{q \in Q} \tilde{\chi}(\mathcal{A}_q) + \sum_{r \in R} \mu_r.$$

Proof. Let (w_1, \dots, w_ρ) be an element of the reference space $\oplus_{i=1}^\rho H_n(E_i^\natural, F_i^\natural) \cong \oplus_{i=1}^\rho \mathbb{Z}\mu_i^\natural$. By the diagonal map this corresponds to elements $w_i \in H_{n+1}(\mathcal{Z}_s, \mathcal{C}_s)$ for $s \in S_q$ and $q \in \Sigma_i$. By the discussion introducing Theorem 6.1 the kernel of some component $j_{1,q} : \oplus_{s \in S_q} H_{n+1}(\mathcal{Z}_s, \mathcal{C}_s) \rightarrow H_{n+1}(\mathcal{X}_q, \mathcal{A}_q)$ is equal to $H_{n+1}(\mathcal{A}_q, \partial_2 \mathcal{A}_q)$ which in turn is identified to the free part of $H_{n-1}(\mathcal{A}_q)$. The rank zero condition implies that $w_i = 0$ for i such that $q \in \Sigma_i$, thus all w_i are zero.

As for the rank of $H_{n+1}(\mathbb{V}_\Delta, V_\varepsilon)$, the formula follows from the Euler characteristic computation (6.2). \square

REMARK 6.6. In case of an irreducible singular set Σ , Corollary 6.5 tells that one singular point $q \in Q$ with a $(n-1)$ th Betti number of the Milnor fiber equal to zero is sufficient for the vanishing of $H_{n+2}^\vee(V)$.

7. COMPUTATIONS OF BETTI NUMBERS

7.1. Vanishing Betti numbers. As direct application of Theorem 6.1, we provide explicit computations of the ranks of the vanishing homology of some projective hypersurfaces.

EXAMPLE 7.1. [some cubic hypersurfaces]

If $V := \{x^2z + y^2w = 0\} \subset \mathbb{P}^3$ then $\text{Sing } V$ is a projective line and its generic transversal type is A_1 . There are three axis points and two special points q with local singularity type D_∞ . The hypersurface singularity germ D_∞ is an *isolated line singularity* in the terminology of [Sil]. Its Milnor fiber F is homotopy equivalent to the sphere S^2 , the transversal monodromy is $-\text{id}$. From Theorem 6.1 it follows that $H_4^\vee(V) \simeq H_1(F) = 0$ and applying Corollary 6.5 we get that $\text{rank } H_3^\vee(V) = 5$.

For $V := \{x^2z + y^2w + t^3 = 0\} \subset \mathbb{P}^4$, $\text{Sing } V$ is again a projective line but its generic transversal type is A_2 , with three axis points and two special points for both of which the local Milnor fiber is homotopy equivalent to $S^3 \vee S^3$. Then Theorem 6.1 yields the isomorphism $H_5^\vee(V) \simeq H_2(F) = 0$ and by Corollary 6.5 we get $\text{rank } H_4^\vee(V) = 10$. This construction can be iterated, for instance $V := \{x^2z + y^2w + t_1^3 + t_2^3 = 0\} \subset \mathbb{P}^5$ has $H_6^\vee(V) = 0$ and $\text{rank } H_5^\vee(V) = 20$.

EXAMPLE 7.2. [including an isolated singular point]

Let $V = \{y^2(x+y-1)(x-y+1) + z^4 = 0\} \subset \mathbb{P}^3$. We have $\text{Sing } V$ is the disjoint union of Σ , a projective line $\{y = z = 0\}$ with transversal type A_3 and a point $R = \{(0 : 1 : 0 : 0)\}$

of type A_3 . There are two special points: $Q = \{(1 : 0 : 0 : 0), (-1 : 0 : 0 : 0)\}$, each of them with Milnor fiber $S^2 \vee S^2 \vee S^2$. It follows that $H_4^Y(V) = 0$ and $\text{rank } H_3^Y(V) = 21$.

EXAMPLE 7.3. [singular locus with two disjoint curve components]

Let $V := \{f = x^2z^2 + x^2w^2 + y^2z^2 + 2y^2w^2 = 0\} \subset \mathbb{P}^3$, which is defined by an element f of the ideal $(x, y)^2 \cap (z, w)^2$. Then $\text{Sing } V = \Sigma = \Sigma_1 \cup \Sigma_2$, where $\Sigma_1 = \{x = y = 0\}$ and $\Sigma_2 = \{z = w = 0\}$. It turns out that the generic transversal type at both of the line components of the singular locus is A_1 and that there are exactly four D_∞ -points on each of these two components. We are in the situation of Corollary 6.5, hence $H_4^Y(V) = 0$ and $\text{rank } H_3^Y(V) = 20$.

7.2. Computation of vanishing homology groups. Using the full details of the proof of Theorem 6.1, we may compute not only the rank of the vanishing homology groups, but in several examples even the vanishing homology group $H_{n+1}^Y(V)$ itself, as follows.

The main ingredient is the map $j^{[k]} = j_1^{[k]} \oplus j_2^{[k]} : H_k(\mathcal{Z}, \mathcal{C}) \rightarrow H_k(\mathcal{X}, \mathcal{A}) \oplus H_k(\mathcal{Y}, \mathcal{B})$, which was denoted by j in (5.2). Like in (6.1), we use the direct sum splitting into axis, special and auxiliary contributions. From the Mayer-Vietoris 6-term sequence (5.3) we derive the short exact sequence:

$$0 \rightarrow \text{coker } j^{[n+1]} \rightarrow H_{n+1}(\mathbb{V}_\Delta, V_\varepsilon) \rightarrow \ker j^{[n]} \rightarrow 0$$

and the strategy will be to work with $j^{[n+1]}$ and $j^{[n]}$ at the level of generators.

EXAMPLE 7.4. Let $V := \{x^2z + y^3 + xyw = 0\} \subset \mathbb{P}^3$. Then $\text{Sing } V$ is a projective line with generic transversal type A_1 , 3 axis points, and a single special point q of local singularity type $J_{2,\infty}$. The latter is an isolated line singularity germ, cf [Si1], with Milnor fiber a bouquet of 4 spheres S^2 and where the transversal monodromy is the identity. By Theorem 6.1 and Corollary 6.5 we get $H_4^Y(V) \simeq H_1(F) = 0$ and $\text{rank } H_3^Y(V) = 6$. We next can show (but skip the details) that there is an isomorphism $H_3^Y(V) \simeq \mathbb{Z}^6$. Note that Dimca [Di2] observed that V has the rational homology of \mathbb{P}^2 .

EXAMPLE 7.5. $V := \{xyz = 0\} \subset \mathbb{P}^3$ of degree $d = 3$. Then V is reducible with 3 components, $\text{Sing } V$ is the union of 3 projective lines intersecting at a single point $[0 : 0 : 0 : 1]$, and the transversal type along each of them is A_1 . Following the proof of Theorem 6.1, we get $\ker j_2^{[3]} = \bigoplus_{i=1}^3 H_1(F_i^{\text{th}}) \simeq \mathbb{Z}^3$ and $\ker j_1^{[3]} \simeq H_1(F)$ where F denotes the Milnor fiber of the non-isolated singularity of V at the single special point $[0 : 0 : 0 : 1]$, which is homotopically equivalent to $S^1 \times S^1$. We thus get $H_4^Y(V) \simeq H_1(F) \simeq \mathbb{Z}^2$.

The axis A of our pencil has degree 9 and intersects each of the components of V at 3 general points. Hence $\nu_i = 3$, $\gamma_i = 1$ for any $i = 1, 2, 3$.

Applying formula (6.2) we get that the vanishing Euler characteristic is -5 , and that $\text{rank } H_3^Y(V) = 7$. We can moreover show the freeness of this group; we skip the details.

7.3. The surfaces case. Several examples in the previous subsections are surfaces and the computation of H_4 and the rank of H_3 could be simplified by counting the number of irreducible components of Σ . Indeed in case of surfaces $V \subset P^3$ we have:

$$(7.1) \quad H_4^Y(V) \simeq \mathbb{Z}^{r-1},$$

where r is the number of irreducible components of V .

We have included these examples anyhow as applications of our method.

Combining (7.1) with Theorem 6.1 yields several consequences on the singular set and its generic transversal types. We mention here only one:

Corollary 7.6. $r - 1 \leq \sum_{i=1}^{\rho} \mu_i^{\natural}$. □

7.4. Absolute homology of projective hypersurfaces. If $\dim \text{Sing } V \leq 1$ then from Theorem 4.1 and the long exact sequence of the pair $(\mathbb{V}_{\Delta}, V_{\varepsilon})$ one gets the isomorphisms:

$$H_k(V) \simeq H_k(V_{\varepsilon}) = H_k(\mathbb{P}^n) \text{ for } k \neq n, n+1, n+2.$$

This corresponds to Kato's result [Ka] in cohomology.⁵

In the remaining dimensions we have an 8-term exact sequence:

$$(7.2) \quad \begin{aligned} 0 \rightarrow H_{n+2}(V_{\varepsilon}) \rightarrow H_{n+2}(\mathbb{V}_{\Delta}) \rightarrow H_{n+2}(\mathbb{V}_{\Delta}, V_{\varepsilon}) \xrightarrow{\Phi_{n+1}} H_{n+1}(V_{\varepsilon}) \\ \rightarrow H_{n+1}(\mathbb{V}_{\Delta}) \rightarrow H_{n+1}(\mathbb{V}_{\Delta}, V_{\varepsilon}) \xrightarrow{\Phi_n} H_n(V_{\varepsilon}) \rightarrow H_n(\mathbb{V}_{\Delta}) \rightarrow 0 \end{aligned}$$

from which we obtain:

Proposition 7.7.

- (a) $b_{n+2}(V) \leq 1 + \sum_{i=1}^{\rho} \mu_i^{\natural}$
- (b) $b_n(V) \leq \dim \mathbb{L}$,

where $\mathbb{L} := H_n(V_{\varepsilon})$ is the intersection lattice of the middle homology of the smooth hypersurface V_{ε} of degree d . In case n is even, this moreover yields:

- (c) $H_{n+2}(V) \simeq \mathbb{Z} \oplus H_{n+2}^{\vee}(V)$,
- (d) $H_{n+1}(V) \simeq \ker \Phi_n$,
- (e) $H_n(V) \simeq \text{coker } \Phi_n$.

This can be regarded as a natural extension of Proposition 2.2 to 1-dimensional singularities, thus extending also Dimca's corresponding result for isolated singularities that was discussed in §2. Like in the isolated singularities setting, one has to deal with the difficulty of identifying Φ_n from the equation of f .

EXAMPLE 7.8. Let $V := \{f(x, y) + f(z, w) = 0\} \subset \mathbb{P}^3$, where $f(x, y) = y^2 \prod_{i=1}^3 (x - \alpha_i y)$, with $\alpha_i \neq 0$ pairwise different. Its singular set is the smooth line given by $y = w = 0$, with generic transversal type $y^2 + w^3$. There are two special points $[0 : 0 : 1 : 0]$ and $[1 : 0 : 0 : 0]$, each with Milnor fiber a bouquet of spheres S^2 . By Corollary 6.5 we get $H_4^{\vee}(V) = 0$ and from the Euler characteristic formula (6.2) (by computing its ingredients) we get $H_4^{\vee}(V) = 0$. One can compute the eigenvalues of the monodromies for all types of singular points; they are all different from 1. By using Randell's criterion [Ra, Proposition 3.6] one can show that V is a \mathbb{Q} -homology manifold. Since a homology manifold satisfies Poincaré duality, it follows e.g. that $H_3(V; \mathbb{Q}) \cong H_1(V; \mathbb{Q}) \cong H_1(\mathbb{P}^n; \mathbb{Q}) = 0$ and $H_4(V; \mathbb{Q}) \cong H_0(V; \mathbb{Q}) \cong H_0(\mathbb{P}^n; \mathbb{Q}) \cong \mathbb{Q}$. By computations and the exact sequence (7.2) we get also $H_2(V; \mathbb{Q})$ since $\text{rank } H_2(V) = \text{rank } H_2(V_{\varepsilon}) - \text{rank } H_3(\mathbb{V}_{\Delta}, V_{\varepsilon}) = 53 - 38 = 15$.

⁵Dimca states such a result [Di4, Theorem 4.1] referring to [Di3, p. 144] for Kato's proof in cohomology [Ka].

REFERENCES

- [Di1] A. Dimca, *On the homology and cohomology of complete intersections with isolated singularities*. Compositio Math. 58 (1986), no. 3, 321-339.
- [Di2] A. Dimca, *On the Milnor fibrations of weighted homogeneous polynomials*. Compositio Math. 76 (1990), 19-47.
- [Di3] A. Dimca, *Singularities and topology of hypersurfaces*. Universitext. Springer-Verlag, New York, 1992.
- [Di4] A. Dimca, *Singularities and their deformations: how they change the shape and view of objects*. Algebraic geometry and geometric modeling, 87101, Math. Vis., Springer, Berlin, 2006.
- [Ka] M. Kato, *Topology of k -regular spaces and algebraic sets*. Manifolds–Tokyo 1973 (Proc. Internat. Conf. on Manifolds and Related Topics in Topology), pp. 153–159. Univ. Tokyo Press, Tokyo, 1975.
- [Lê] Lê D.T., *Calcul du nombre de cycles évanouissants d’une hypersurface complexe*. Ann. Inst. Fourier (Grenoble) 23 (1973), no. 4, 261–270.
- [LW] A.S. Libgober, J.W. Wood, *On the topological structure of even-dimensional complete intersections*. Trans. Amer. Math. Soc. 267 (1981), no. 2, 637–660.
- [PP] A. Parusiński, P. Pragacz, *Characteristic classes of hypersurfaces and characteristic cycles*. J. Algebraic Geom. 10 (2001), no. 1, 63–79.
- [Ra] R.Randell, *On the topology of non-isolated singularities*. In: Geometric Topology (Proc. 1977 Georgia Topology Conference), Ac. Press, New York(1979) 445–473.
- [Si1] D. Siersma, *Isolated line singularities*. Singularities, Part 2 (Arcata, Calif., 1981), 485–496, Proc. Sympos. Pure Math., 40, Amer. Math. Soc., Providence, RI, 1983.
- [Si2] D. Siersma, *Quasihomogeneous singularities with transversal type A_1* . Singularities (Iowa City, IA, 1986), 261–294, Contemp. Math., 90, Amer. Math. Soc., Providence, RI, 1989.
- [Si3] D. Siersma, *Variation mappings on singularities with a 1-dimensional critical locus*. Topology 30 (1991), no. 3, 445–469.
- [Si4] D. Siersma, *The vanishing topology of non isolated singularities*. New developments in singularity theory (Cambridge, 2000), 447–472, NATO Sci. Ser. II Math. Phys. Chem., 21, Kluwer Acad. Publ., Dordrecht, 2001.
- [ST1] D. Siersma, M. Tibăr, *Deformations of polynomials, boundary singularities and monodromy*, Mosc. Math. J. 3 (2) (2003), 1–19.
- [ST2] D. Siersma, M. Tibăr, *Betti numbers of polynomials*, Mosc. Math. J. 11 (2011), no. 3, 599–615.
- [Ti1] M. Tibăr, *Connectivity via nongeneric pencils*. Internat. J. Math. 13 (2002), no. 2, 111–123.
- [Ti2] M. Tibăr, *Polynomials and vanishing cycles*. Cambridge Tracts in Mathematics, 170. Cambridge University Press, Cambridge, 2007. xii+253 pp.
- [Ti3] M. Tibăr, *The vanishing neighbourhood of non-isolated singularities*. Israel J. Math. 157 (2007), 309–322.
- [Th] R. Thom, *Ensembles et morphismes stratifiés*, Bull. Amer. Math. Soc., 75 (1969), 249-312.
- [Yo] I.N. Yomdin, *Complex surfaces with a one-dimensional set of singularities*. Sibirsk. Mat. Z. 15 (1974), 1061–1082, 1181.

INSTITUTE OF MATHEMATICS, UTRECHT UNIVERSITY, PO BOX 80010, 3508 TA UTRECHT THE NETHERLANDS.

E-mail address: D.Siersma@uu.nl

MATHÉMATIQUES, UMR 8524 CNRS, UNIVERSITÉ LILLE 1, 59655 VILLENEUVE D’ASCQ, FRANCE.

E-mail address: tibar@math.univ-lille1.fr